



# Weekly Assessment of CVP and SWP Delta Operations on ESA-listed Species

June 11, 2023

## Executive Summary

### Operational Conditions

See Weekly Fish and Water Operation Outlook document for June 11 – June 17.

### Winter-run Chinook Salmon

Loss of natural winter-run Chinook Salmon (by length at date, LAD) has not occurred in the past week at either facility, and loss of genetically confirmed winter-run Chinook Salmon has not occurred at either facility since 3/21/2024. 0% of juvenile natural winter-run Chinook Salmon from brood year (BY) 2023 are estimated to be present in the Delta. The 50% single year loss threshold for LAD winter-run Chinook Salmon was exceeded on 2/25/2024. The 75% single year loss threshold for LAD winter-run Chinook Salmon was exceeded on 3/7/2024. The 100% single year loss threshold for LAD winter-run Chinook Salmon was exceeded on 3/20/2024. Loss of natural LAD winter-run Chinook Salmon at the CVP and SWP fish collection facilities is unlikely to occur over the next week due to seasonal timing. Greater than 95% of winter-run Chinook Salmon are estimated to have migrated past Chipps Island and 4 of the seven non-consecutive days of temperatures greater than 22.2 C have been observed at both Mossdale and Prisoners Point as of 6/9

### Spring-run Chinook Salmon

Loss of natural YOY LAD spring-run Chinook Salmon has occurred in the past week at the State and Federal fish salvage facilities. 0 % of juvenile natural YOY spring-run Chinook Salmon was estimated in the Delta. CV spring-run Chinook Salmon fry are rearing and migrating downstream. Spring-run Chinook Salmon LAD loss has decreased at the salvage facilities due to seasonal timing. Due to the wetter conditions this season and cool water temperatures, it is likely that spring-run Chinook Salmon will out-migrate later this year than in a year with drier and warmer hydrological conditions. Therefore spring-run loss may continue at a similar trend through May (see Table A1). Loss of natural spring-run Chinook Salmon at the CVP and SWP fish collection facilities is unlikely to occur over the next week. Greater than 95% of spring-run Chinook Salmon are estimated to have migrated past Chipps Island and 4 of the seven non-consecutive days of temperatures greater than 22.2 C have been observed at both Mossdale and Prisoners Point as of 6/9.

## Central Valley Steelhead

Loss of natural CCV steelhead has occurred in the past week at the State and Federal fish salvage facilities. Loss of CCV steelhead at the CVP and SWP fish collection facilities may occur over the next week. 0-8 % of CCV steelhead were estimated in the Delta. The 50% single year loss threshold for the December 1 to March 31 period was exceeded on 2/11/2024 the 75% single year loss threshold was exceeded on 2/22/2024, and the 100% single year loss threshold was exceeded on 2/23/2024. The March – December ITL was exceeded on 3/20/2024. Steelhead loss as of 5/23/2024 for the April 1 – June 16 period is at 63% of the April-June ITL. The 50% single year loss threshold for the April 1 to June 15 period was exceeded on 4/9/2024, the 75% threshold was exceeded on 4/15/2024, and the 100% threshold was exceeded on 4/26/2024. Less than 95% of steelhead are estimated to have migrated past Chipps Island and 4 of the seven non-consecutive days of temperatures greater than 71.6 F have been observed at Mossdale as of 6/9 OMR management for CCV steelhead protections will offramp on June 15th.

## DCC gates recommendation

The DCC gates are currently closed but will open on 6/14 for recreation and salinity.

## Delta Smelt

Based on recent detection data and distribution patterns over the past decade, adult Delta Smelt spawning is ongoing. Twenty-three confirmed larval Delta Smelt have been detected in Suisun Bay, Suisun Marsh, the Lower Sacramento, the Sacramento River Deep Water Shipping Channel, Cache Slough/Liberty Island, and the South Delta since 5/6/2024. While there is a greater chance of entraining additional DS this week at more negative OMRI (up to -5,000 cfs) and negative QWEST (up to -2,000 cfs), risk of entrainment throughout the Delta remains low for OMRI of -3,500 to -5,000 cfs for all life stages based on low turbidity/ Secchi depth >1m, and the majority of detections outside of the Central and South Delta.

## Monitoring Teams summary

Salmon Monitoring Team had no issues to raise to WOMT

## Operational and Regulatory Conditions

See current Weekly Fish and Water Operation Outlook document.

## Biology, Distribution, and Evaluation Winter-run Chinook salmon, Spring-run Chinook salmon, Central Valley Steelhead

### Population Status

#### *Winter-run Chinook Salmon*

- Delta Life Stages:

- Juveniles, Adults
- Brood Year 2023 Productivity:
  - Natural winter-run Chinook salmon: The Final WR JPE for BY 2023 is 234,896.
  - Mean cumulative weekly passage of winter-run Chinook salmon through January 14th at Red Bluff Diversion Dam (RBDD) for the last 21 years of passage data is 99.8% (one SD of 0.2%). By 3/24/2024, 1,110,528 winter-run Chinook salmon were estimated to have passed RBDD compared to the total cumulative passage for WY 2023 of 240,059 winter-run Chinook salmon.
  - Hatchery winter-run Chinook salmon:
    - One release of Livingston Stone National Fish Hatchery CWT tagged winter-run Chinook salmon occurred on 12/28/23.
    - The second release of LSNFH CWT tagged winter-run Chinook salmon occurred on 1/19/24.
    - A third release of LSNFH CWT tagged winter-run Chinook salmon occurred on 2/16/24. A subset of these fish were acoustically tagged;
    - 47,937 Coleman NFH Complex brood year 2023 winter Chinook Salmon on March 4, 2024. The release took place on the North Fork Battle Creek at Wilson Hill Bridge near Manton, CA. 100% marked (with an adipose-fin and a left pelvic-fin clip and CWT) and has an overall estimated average fork length of 85 mm.
    - 84,994 Coleman NFH Complex brood year 2023 winter Chinook Salmon on March 11, 2024. The release took place on the North Fork Battle Creek at Wilson Hill Bridge near Manton, CA. 100% marked (with an adipose-fin and a left pelvic-fin clip and CWT) and has an overall estimated average fork length of 85 mm.
    - None of the winter-run from the above two Coleman NFH hatchery release groups have been detected in salvage.

### ***Spring-run Chinook Salmon***

- Delta Life Stages:
  - Young-of-year (YOY) and Yearlings
- Brood Year 2023 Productivity:
  - Natural spring-run Chinook salmon: No JPE has been established for spring-run Chinook salmon.

- Hatchery spring-run Chinook salmon surrogates associated with the Proposed Action (PA 4.10.5.10.2 Additional Real-Time OMR Restrictions and Performance Objectives):
  - Approximately 60,764 brood year 2023 late-fall Chinook salmon were released on December 22, 2023 into Battle Creek at the Coleman NFH. This group is 100% marked with an adipose-fin clip and coded-wire tagged.
  - Approximately 71,049 Coleman NFH brood year 2023 late-fall Chinook Salmon were released on December 29, 2023 into Battle Creek at Coleman NFH. This group is 100% marked (with an adipose-fin clip and CWT) and has an overall estimated average fork length of 145 mm.
  - Approximately 67,018 Coleman NFH brood year 2023 late-fall Chinook Salmon were released on January 11, 2024 into Battle Creek at Coleman NFH. This group is 100% marked (with an adipose-fin clip and CWT) and has an overall estimated average fork length of 145 mm.
- Loss has occurred for the above release group 1, release group 2, and release group 3 of the late-fall hatchery yearling spring-run surrogates released between December 22nd and January 11th.
- A release of 699,854 brood-year 2023 spring-run Chinook salmon from the Feather River hatchery at Gridley Boat Launch and Boyd's Pump Boat Launch occurred on March 15, 2024. This group was 100% marked (with an adipose fin clip and CWT) and had an overall estimated average fork length of 74 mm.
- Approximately 712,177 Coleman NFH brood year 2023 fall-run Chinook Salmon were released on March 21, 2024 into Battle Creek at Coleman NFH. This group is 25% marked (with an adipose-fin clip and CWT) and has an overall estimated average fork length of 75 mm.
- A release of 700,626 brood-year 2023 spring-run Chinook salmon from the Feather River hatchery at Gridley Boat Launch and Boyd's Pump Boat Launch occurred on March 29, 2024. This group was 100% marked (with an adipose fin clip and CWT) and had an overall estimated average fork length of 84 mm.
- A release of 210,351 brood-year 2023 fall-run were released from Nimbus Hatchery on 4/19/24 serving as surrogates for young-of-year spring-run migrants. This group was 25% marked (with an adipose fin clip and CWT) and had an overall estimated average fork length of 90 mm.
- A release of 560,304 brood-year 2023 spring-run Chinook salmon from the Feather River hatchery at Gridley Boat Launch and Boyd's Pump Boat Launch occurred on April 23, 2024. This group was 100% marked (with an adipose fin clip and CWT) and had an overall estimated average fork length of 94 mm.

- A release of 106,531 brood year 2023 fall-run Chinook salmon from Coleman National Fish Hatchery at Butte City Boat Ramp serving as surrogates for young-of-year spring-run migrants. This group was 25% marked (with an adipose fin clip and CWT) and had an overall fork length of 75 mm.
- Of the six above young-of-year spring-run surrogate groups released between March 15 and May 1, none have been detected in salvage.

### ***Central Valley Steelhead***

- Delta Life Stages:
  - Spawning Adults, Kelts, Juveniles
- Brood Year 2023 Productivity:
  - Spawner abundance: There is limited information about the adult steelhead population. It is estimated to be small, contributing to the limited productivity of the population.
  - The majority of wild and hatchery steelhead observed at the salvage facilities are between 200 and 300 mm fork length (Figure 8). This is similar to the size distribution of age-1 clipped hatchery steelhead. However, wild steelhead in the 200-300 mm size range are likely to be comprised of both age-1 and age-2 individuals. While the size distribution of steelhead at salvage is similar to the sizes of age-1 *O. mykiss* at the American River, this is the typical size of age-2 steelhead from the Stanislaus River, Mokelumne River, Clear Creek, and Mill Creek (Satterthwaite et al. 2010, Eschenroeder et al. 2022, Lisa Elliott – unpublished data).
  - Natural steelhead: No JPE has been established for steelhead. Data are limited.
  - Hatchery steelhead: Reclamation’s Proposed Action has no hatchery steelhead triggers, but see Table 7 for information on hatchery steelhead release and annual loss numbers.
  - A release of 240 hatchery steelhead occurred on 3/15/24 at Lighthouse Marina from the Mokelumne hatchery. These fish were acoustically tagged and tracked through CalFishTrack as the Mokelumne Hatchery Steelhead Export Effects Study. The goal of the release is to evaluate the effect of steelhead movement through the Delta at -500 cfs OMR. As of 5/6, 98 of the 240 steelhead released have reached Benicia Bridge East Span and 1 has been observed in salvage.
  - Three hundred and thirty-five 6-year Steelhead study hatchery fish were released on 3/20/24 into the Lower San Joaquin River. Of these 27 were detected at the SWP intake canal and 10 in the CVP. A further 57 fish have reached the Benicia Bridge.

- As of 4/10/24, 179 wild steelhead have been acoustically tagged and released in the Stanislaus River, with 2 of those fish so far being detected at Benicia and 4 being detected at the salvage facilities.
- As of 4/19/24, 335 hatchery steelhead were released into the Lower San Joaquin River. Of these 12 were detected at the SWP intake canal and 9 in the CVP. A further 37 fish have reached the Benicia Bridge.
- As of 4/26/24, 157 wild steelhead have been acoustically tagged and released in Mill and Deer Creeks in the Sacramento River. Sixty-nine of those fish have been detected at Benicia, and none have been detected at the salvage facilities.
- As of 5/9/24, 236 hatchery steelhead were released into the Lower San Joaquin River. Of these 5 has been detected at the SWP intake canal and 8 in the CVP. Twenty-one have reached the Benicia Bridge.

## Distribution

### *Winter-run Chinook Salmon*

#### Current Distribution:

- For winter-run Chinook Salmon observations reported to SaMT since previous meeting, see Table 1.
- For SaMT distribution estimates, see Table 2.
- As of 5/24/2024, 16 genetic natural winter-run Chinook Salmon have been identified at the SWP facility. As of 5/24/2024, 7 genetic natural winter-run Chinook Salmon have been identified at the CVP facility.
- The 2024 IOP was implemented on 3/29/2024.
- Acoustically tagged hatchery winter-run Chinook salmon are no longer being observed passing receivers at Benicia Bridge and none have been detected in salvage.

#### Historical Trends

- For historical winter-run Chinook salmon trends in salvage, see Table 3. See also Attachment 1.
- Loss of natural LAD and genetic winter-run Chinook salmon at the CVP and SWP fish collection facilities is likely to remain zero over the next week based on seasonal timing and previous years salvage data.
- Genetic data since 1996 has indicated that there have been zero observations of genetic winter-run Chinook salmon within the Delta past May 1st (Sacramento trawl, Chipp Island trawl, collection at the salvage facilities) (see Brandes et al. 2021, Kevin Reece and Brian Mahardja unpublished data).

## Forecasted Distribution within Central Valley and Delta regions

- The STARS model projects route-specific proportion of entrainment, survival, and travel times (Table 5). This model does not estimate entrainment into the lower Sacramento River sloughs (i.e., Three-Mile Slough).

### *Spring-run Chinook salmon*

#### Current Distribution

- For spring-run Chinook salmon observations reported to SaMT since previous meeting, see Table 1.
- For SaMT distribution estimates, see Table 2.

#### Historical Trends

- For historical spring-run Chinook salmon trends in salvage, see Table 3. Based on historic trends in salvage YOY spring-run Chinook salmon loss may continue over the next week.

## Forecasted Distribution within Central Valley and Delta regions

- YOY spring-run Chinook are migrating out of the Delta.

### *Central Valley Steelhead*

#### Current Distribution

- For CCV Steelhead observations reported to SaMT since previous meeting, see Table 1.
- For SaMT distribution estimates, see Table 2.
- As of 6/10/2024, 757 unclipped CCV steelhead have been observed at the CVP or SWP facilities.
- Loss of CCV Steelhead exceeded the 50%,75%, 100%, annual loss thresholds and the ITL for the Dec. 1 to March 31 period on 2/11/2024, 2/22/2024, 2/23/2024, and 3/20/2024, respectively.
- Loss of CCV Steelhead exceeded the 50% annual loss thresholds for the April 1 to June 15 period on 4/9/2024, the 75% threshold on 4/15/2024, and the 100% threshold on 4/26/2024.
- Steelhead have been observed at the salvage facilities in the last 7 days.
- Steelhead have recently been observed in the Chipps Island Trawl which supports out-migration is occurring.
- Acoustically tagged steelhead from the Stanislaus, Mokelumne, and San Joaquin River have been observed passing receivers at Benicia Bridge.

## Historical Trends

- At Red Bluff Diversion Dam, the estimated passage of BY 2023 steelhead is 57,255, which is the second highest in the last 10 brood years.
- For historical CCV steelhead trends in salvage, see Table 3. The estimated number of steelhead salvaged for water year 2024 as of 6/10/24 is 2355 and the highest in the last 10 water years. Based on historic trends in salvage, juvenile CCV steelhead loss may occur over the next week.
- Forecasted Distribution within Central Valley and Delta regions
- The entrainment tool estimates CCV steelhead loss to continue (Table 6, Fig. 1).

Table 1. Fish observation reported since the previous SaMT meeting. Only data reported prior to Tuesday’s meeting is included. See Operations Outlook for notes on interruptions in any surveys. WR represent winter-run, SR represent spring-run, FR represent fall-run, LF represent late-fall run, All represent chinook with no run designation, SH represent steelhead, and GS represent green sturgeon.

Clipped	Sample	Dates	WR	SR	FR	LF	All	SH	GS
N	Butte	06/03 - 06/09	0	17	0	0	0	3	0
N	Knights Landing RST	06/05 - 06/07	0	0	0	0	0	0	0
N	Lower Sacramento RST	06/01 - 06/04	0	0	0	0	0	0	0
N	Feather River (Herringer)	05/22 - 06/03	0	0	13	1	0	1	0
N	Feather River (Eye Side)	05/22 - 06/03	0	0	592	0	0	18	0
N	Yuba	05/22 - 06/03	0	0	287	21	0	28	0
N	Lower Feather RST	05/25 - 06/09	0	0	31	1	0	1	0
N	Beach Seines	05/22 - 05/22	0	2	0	0	0	0	0
N	Sacramento Trawls	06/03 - 06/07	0	0	5	0	0	0	0
N	Mossdale	06/05 - 06/11	0	1	13	0	0	0	0
N	Chipps Island Trawls	06/03 - 06/07	0	0	7	0	0	1	0
Y	Knights Landing RST	06/05 - 06/07	0	0	0	0	0	0	0
Y	Lower Sacramento RST	06/01 - 06/04	0	0	0	0	0	0	0
Y	Feather River (Herringer)	05/22 - 06/03	0	0	0	0	0	0	0
Y	Feather River (Eye Side)	05/22 - 06/03	0	0	0	2	0	0	0
Y	Yuba	05/22 - 06/03	0	0	0	0	0	0	0
Y	Lower Feather RST	05/25 - 06/09	0	0	0	0	0	2	0
Y	Mossdale	06/05 - 06/11	0	0	0	0	0	0	0



Table 2. Salmonid distribution estimates

Location	Yet to Enter Delta (%)	In the Delta (%)	Exited Delta past Chipps Island (%)
Young-of-year (YOY) winter-run Chinook salmon	Current: 0% Last Week: 0%	Current: 0% Last Week: 0-1%	Current: 100% Last Week: 99-100%
YOY spring-run Chinook salmon	Current: 0% Last Week: 1-2%	Current: 0-1% Last Week: 1-3%	Current: 99-100% Last Week: 95-98%
YOY hatchery winter-run Chinook salmon	Current: 0% Last Week: 0%	Current: 0 % Last Week: 0%	Current: 100% Last Week: 100%
Natural origin steelhead	Current: 0-2% Last Week: 1-5%	Current: 0-8 % Last Week: 9-15%	Current: 90-100% Last Week: 80-90%

Table 3. Historic migration and salvage patterns. Last updated 06/11/2024.

Species	Red Bluff Diversion Dam	Tisdale Rst	Knights Landing Rst	SacTrawl Sherwood Catch Index	Chipps Island Trawl Catch Index	Salvage
Chinook, Winter-run, Unclipped	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) WY: 2014 - 2023
Chinook, Spring-run, Unclipped	100.0%(99.9% ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	100.0%(100.0 % ,100.0%) BY: 2014 - 2022	99.5%(98.8%,100.3%) WY: 2014 - 2023
Steelhead, Unclipped (January-December)	39.3%(20.8%, 57.8%) BY: 2014 - 2023	86.7%(65.8%, 107.5%) BY: 2014 - 2023	81.1%(64.1%, 98.1%) BY: 2014 - 2023	100.0%(100.0 % ,100.0%) BY: 2014 - 2023	96.0%(91.2%, 100.7%) BY: 2014 - 2023	N/A
Steelhead, Unclipped (December-March)	N/A	N/A	N/A	N/A	N/A	100.0%(100.0 % ,100.0%) WY: 2015 - 2024
Steelhead, Unclipped (April-June)	N/A	N/A	N/A	N/A	N/A	98.0%(95.4%,100.6%) WY: 2014 - 2023

Table 4. Mean daily flow and percent change (Wilkins Slough, Deer Creek, Mill Creek; cfs from CDEC) and temperature and percent change (Knights Landing; °F from RST).

Date	Mill Creek (MLM): mean daily flow (cfs)	Mill Creek (MLM): flow percent change	Mill Creek (MLM): Alert	Deer Creek (DCV): mean daily flow (cfs)	Deer Creek (DCV): flow percent change	Deer Creek (DCV): Alert	Wilkins Slough (WLK): mean daily flow (cfs)	Knights Landing RST: water temperature (f)	Alert Triggered
6/9/2024	407.7	-3.6%	Flow>95cfs	241.4	-3.7%	Flow>95cfs	5,780.2	N/A	N/A
6/8/2024	422.8	-2.6%	Flow>95cfs	250.5	-4.8%	Flow>95cfs	5,826.4	N/A	N/A
6/7/2024	434.0	-1.2%	Flow>95cfs	263.0	-5.5%	Flow>95cfs	6,041.8	N/A	N/A
6/6/2024	439.2	2.1%	Flow>95cfs	278.2	-4.8%	Flow>95cfs	6,191.4	N/A	N/A
6/5/2024	430.2	4.0%	Flow>95cfs	292.2	-5.5%	Flow>95cfs	6,390.5	N/A	N/A
6/4/2024	413.8	-4.9%	Flow>95cfs	309.3	-0.2%	Flow>95cfs	6,441.0	N/A	N/A
6/3/2024	435.2	1.8%	Flow>95cfs	309.9	-1.7%	Flow>95cfs	6,553.5	N/A	N/A

Table 5. STARS model simulations for route-specific, travel times, and survival. Travel time is calculated in days.

Stock	Date	Route	Median Travel Time	Survival	Routing Probability
Winter Chinook	2024-06-02	Overall	6.52	0.00	N/A
Winter Chinook	2024-06-09	Overall	6.73	0.00	N/A
Winter Chinook	2024-06-09	Sacramento River	6.29	0.00	0.60
Winter Chinook	2024-06-09	Yolo Bypass	10.13	0.30	0.00
Winter Chinook	2024-06-09	Sutter Slough	6.40	0.00	0.14
Winter Chinook	2024-06-09	Steamboat Slough	6.14	0.00	0.14
Winter Chinook	2024-06-09	Interior Delta	9.75	0.00	0.13
Late-fall Chinook	2024-06-09	Overall	10.74	0.45	N/A
Late-fall Chinook	2024-06-09	Delta Cross Channel	N/A	N/A	0.00
Late-fall Chinook	2024-06-09	Georgiana Slough	15.40	0.21	0.25
Late-fall Chinook	2024-06-09	Sacramento River	9.25	0.58	0.47

The entrainment tool estimates a median and maximum loss of winter-run Chinook Salmon and juvenile CCV Steelhead each week (Table 6a).

Table 6a-c. WY 2024 loss and salvage predictor data: Environmental details, current and forecast. Unless values (i.e. OMR and export) are explicitly stated in Table 6c, covariate values are the same as values listed in 6b. Model results from 6/11/2024.

- a) WY 2024 loss and salvage predictor data: Predicted weekly loss of winter-run Chinook salmon and steelhead at CVP and SWP facilities estimated from model developed in Tillotson et al. (2022).

Parameter	Modeled Current Week	Modeled Next Week
Predicted Steelhead, Median %	0	0
Predicted Steelhead, High %	123	17
Predicted Chinook Winter Run, Median %	0	0
Predicted Chinook Winter Run, High %	0	0

- b) Environmental details, current and forecast

Parameter	Data	Forecast
Temperature (Mallard Island, C)	21.1	21.1
Precipitation (5-d running sum, inches)	0	0
Old and Middle River Flows (cfs)	-5654	-5654
Sacramento River Flow (Freeport, cfs)	16064	16064
DCC Gates	closed	closed
San Joaquin River Flow (Vernalis, cfs)	3416	3416
Export	5421	5421

- c) Salvage predictor estimates for natural origin Central Valley Steelhead from model developed in Tillotson et al. (2022)<sup>1</sup> across multiple OMR flow scenarios represented within a range of percentiles. The model with expected discharge values (Discharge Range) in the Sacramento and San Joaquin Rivers through 6/14. Only the top two scenarios are feasible within the current operational outlook.

OMR	Exports	San Joaquin Flow	Sacramento Flow	25th Percentile Weekly Loss	Median Weekly Loss	75th Percentile Weekly Loss	Median Daily Loss
-832	1500	3087	14789	0	0	17.32	0
0	600	3087	14789	0	8.16	25.98	1.17

<sup>1</sup> Tillotson, M.D., J. Hassrick, A.L. Collins, & C. Phillis. 2022. Machine Learning Forecasts to Reduce Risk of Entrainment Loss of Endangered Salmonids at Large-Scale Water Diversions in the Sacramento-San Joaquin Delta, California. San Francisco Estuary & Watershed Science 20(2):1-21

OMR	Exports	San Joaquin Flow	Sacramento Flow	25th Percentile Weekly Loss	Median Weekly Loss	75th Percentile Weekly Loss	Median Daily Loss
-500	1125	3087	14789	0	8.16	25.98	1.17
-1000	1700	3087	14789	0	0	25.98	0
-1500	2225	3087	14789	0	0	9.34	0
-2000	2800	3087	14789	0	0	6.915	0
-2500	3300	3087	14789	0	0	5.44	0
-3000	3900	3087	14789	0	0	8.16	0
-3500	4400	3087	14789	0	0	13.42	0
-4000	5000	3087	14789	0	0	25.98	0
-4500	5500	3087	14789	0	0	17.32	0
-5000	6000	3087	14789	0	0	17.32	0

Table 7. Annual loss of clipped juvenile steelhead at the salvage facilities and total hatchery juvenile steelhead release numbers for brood years 2016 to 2023\*. From 2016 to 2024, average annual % lost to the facilities was 0.153%. Note that release locations and dates, which vary by year, were not considered for this calculation. Hatchery release numbers were acquired from: CDFW hatchery releases- Calfish.org and USFWS hatchery releases data provided by Kevin Offill, 3/13/2024. Water facility loss data acquired from: SacPAS and reflects Water Year 2017 - 3/12/2024. \* BY23 includes all fish released prior to 3/14/2024 (data is preliminary)

Brood Year	Total Hatchery Steelhead Release Number (BY)	Loss of clipped steelhead at the facilities (WY)	% Total Hatchery Number Lost to the Facilities	Water Year
2016	1,019,501	164.29	0.016	2017
2017	811,379	2,462.90	0.304	2018
2018	1,264,939	5,777.70	0.457	2019
2019	1,084,899	659.44	0.061	2020
2020	1,853,751	341.69	0.018	2021
2021	1,676,701	639.79	0.038	2022
2022	1,623,483	3,650.30	0.225	2023
2023*	1,517,998	1,581.93	0.104	2024

“Today, the Central Valley O. mykiss population complex is dominated by hatchery-origin fish” (NMFS 2003; Lindley et al. 2007). Best available information indicates that natural-origin O. mykiss comprise only 6% to 16% of the O. mykiss in the northern Sacramento-San Joaquin Delta and San Francisco Estuary. While diversifying the portfolio of release practices is expected to benefit stock complex resiliency and stability, it is also important to consider the potential genetic

and ecological impacts of releasing hatchery-origin pre-smolts on natural-origin *O. mykiss* and other life stages of hatchery-origin *O. mykiss*." See Eric R. Huber, Rachael E. Ryan, Rachel C. Johnson, Anna M. Sturrock, Robert A. Lusardi, and Stephanie M. Carlson. Seventy years of diminishing biocomplexity of California Central Valley hatchery steelhead, *Oncorhynchus mykiss*. *Canadian Journal of Fisheries and Aquatic Sciences*.





Figure 1. Predicted weekly loss of winter-run Chinook salmon and steelhead at the CVP and SWP facilities.

Figure 1 is two-line graphs of the predicted weekly loss of steelhead and winter-run Chinook salmon for water year 2024 beginning on December 1, 2023. The first line graph shows the cumulative loss of winter-run Chinook salmon comparing the predicted loss of 4100 and the observed loss of 4100. The second line graph shows the cumulative loss of Steelhead salmon comparing the median predicted loss of about 5000 and the observed loss of about 5000.





Figure 2. Cumulative natural winter-run Chinook salmon (Top) and natural Steelhead (Bottom) loss for the year (blue) and 2009 – 2018 historic cumulative loss (gray, different symbols). Historic daily mean plotted in black circles.

Figure 2 is two line graphs showing the natural winter-run Chinook salmon and natural steelhead loss and the 2009-2018 cumulative loss.

The first line graph shows the WY 2024 natural winter-run Chinook salmon cumulative loss to date at 4205.13, with a cumulative loss and historic daily mean of just over 4200.

The second graph shows the WY 2024 cumulative loss from December 1, 2023 until March 31, 2024 at 3374.55. The WY 2024 cumulative loss for the period of April 1, 2024 until June 15, 2024 is 1768.56.






Figure 3. LAD winter-run historic cumulative loss from salvage facilities from water year (WY) 1994-2024. The solid blue line represents 2024 and to ensure readability, only cumulative loss curves that have exceeded current 2024 cumulative loss were labeled by WY (updated through 3/21). The 3 overlapping years are 1999, 2002, and 2007.

Figure 3 is a line graph that shows 2024 cumulative loss at around 4000 and highlights years that have exceeded this number which include 1995, 2000, 2001, 2003, 2004, and 2011. Cumulative loss is broken down into wet and dry as well as pre-2009 BiOp and following the 2009 and 2019 BiOps.




Figure 4. (Top-right) Cumulative natural LAD winter-run, (Top-left) natural genetic winter-run, and (Bottom) natural Steelhead with relevant loss thresholds. Genetic loss thresholds are purely for informative purposes to SaMT and WOMT, and are based on an unimplemented proposed action in which a 100% loss threshold is equal to 0.5004% of the JPE and 75% and 50% of that threshold respectively. The unimplemented PA and subsequent 0.5004% of the JPE threshold are a product of the reinitiation of consultation on the long-term operation of the Central Valley Project and State Water Project and may represent annual thresholds if the proposed action resulting from the consultation is implemented.

Figure 4 is three line graphs, the first of which shows cumulative natural LAD winter-run loss, the second shows natural genetic winter-run loss, and the third shows natural steelhead loss.

The first line graph shows the WY 2024 cumulative natural LAD winter-run loss at 4205.05 or 89.5 percent of the single year incidental take limit.

The second line graph shows the WY 2024 winter-run genetic loss at 127.45 or 10.8 percent of the annual loss threshold.

The third line graph shows the WY 2024 cumulative loss at 3374.81 at the end of the early period. Since then the loss has reached 1910.78 or 62.9% percent of the annual loss threshold.




Figure 5. Central Valley Steelhead historic cumulative loss from salvage facilities from water year (WY) 1994-2024. The solid blue line represents 2024 and to ensure readability, only cumulative loss curves that have exceeded current 2024 cumulative loss were labeled by WY (updated through 3/21/24).

Figure 5 is a line graph that shows current and historic cumulative steelhead loss prior to the 2009 BiOP and following the 2009 and 2019 BiOPs, broken down by wet and dry.




Figure 6. Fork length distribution of wild and hatchery steelhead collected at the CVP and SWP salvage facilities in WY 2024. Data included steelhead through 3/17/2024.

Figure 6 is a bar chart which shows the fork length distribution of wild and hatchery steelhead collected at the CVP and SWP facilities in WY 2024. The data is broken down into fish collected with 'clipped' and 'unclipped' adipose fins, with unclipped having a greater fork length and being collected in higher numbers.

## Evaluation

1. After January 1, are more than 5% of juveniles from one or more salmonid species present in the Delta?
  - a. Greater than 5% of Central Valley Steelhead are estimated to be present in the Delta.
2. Does the operational outlook's ranges impact fish movement and change the potential distribution of fish?
  - a. Potential effects within the 7 days (near-term) in the operations outlook.



loss threshold was exceeded on 2/22/2024, the 100% loss threshold was exceeded on 2/23/2024, and the incidental take limit was exceeded on 3/20/2024.

- iv. A means of calculating a juvenile production estimate currently does not exist for steelhead, limiting the ability of SaMT members to assess the impact of loss on the basin-wide or genetically distinct populations of *Oncorhynchus mykiss*. Information on steelhead abundance and distribution in the Delta is also limited, as the existing fish surveys do not target steelhead and may be ineffective at catching these larger-sized fish (yearling or older). Although the proportion of wild steelhead population lost at the salvage facilities is unknown due to lack of WY 2024 population size, the proportion of hatchery or tagged steelhead observed at the facilities relative to release or tagged fish numbers can potentially offer insight into the proportional entrainment of wild steelhead. Annual loss of clipped steelhead at the salvage facilities represent between 0.02 to 0.46% of total hatchery release numbers between water year 2017 and 2024 (Table 7).
  - v. Based on telemetry studies from this water year [CalFish Track Central Valley Enhanced Acoustic Tagging Project](#), a substantial proportion of *Oncorhynchus mykiss* from the San Joaquin River and Stanislaus River have been contacted in the vicinity of the salvage facilities, whereas none from the Sacramento River (Mill Creek and Deer Creek) have been found in the vicinity of the salvage facilities. Moreover, only one of the 240 tagged Mokelumne River hatchery *Oncorhynchus mykiss* released into the central Delta have been detected at the salvage facilities. Telemetry data from this year suggests that the majority of salvaged Steelhead were of San Joaquin origin.
  - vi. Salmonid species are expected to have increased survival when flows are high (Michel et al. 2021) and acoustic telemetry studies indicated that survival is correlated with increased flows at Vernalis for San-Joaquin origin Steelhead (Buchanan et al. 2021), potentially influencing the number of fish observed at the salvage facilities.
4. If an annual loss threshold has been exceeded, do continued OMR restrictions benefit fish movement and survival based on real-time information?
    - a. Winter-run Chinook salmon
      - i. The 100% annual loss threshold for winter-run Chinook salmon was exceeded in WY 2024 on 3/20/2024. However, winter-run presence

in June is highly unlikely according to historical data and seasonal timing of winter-run.

b. Central Valley Steelhead

- i. The 100% annual loss threshold for steelhead (December 1 – March 31) was exceeded for WY 2024. Steelhead loss exceeded the 50% annual loss threshold on 4/9/2024, the 75% single year loss threshold on 4/15/2024, and the 100% single year loss threshold on 4/26/2024 for the April 1 – June 15 period.
- ii. It is uncertain if additional OMR restrictions benefit fish movement and survival. Buchanan et al.'s (2021) study indicated that high flows increase through-Delta survival of steelhead, with little to no association between Delta export and survival. However, this study was conducted in 2011-2016 when OMR was restricted to be no more negative than -5,000 cfs, and therefore does not necessarily suggest that OMR flow more negative than -5,000 cfs cannot be associated with lower survival.
- iii. STARS model output indicates higher survival for Sacramento River-origin salmon in the Delta relative to the past several years due to hydrological conditions ([San Francisco Bay Delta Survival, Travel time, and Routing Simulation \(STARS\)](#)). Steelhead appear to be outmigrating at larger sizes than Chinook salmon based on salvage data (~200-300 mm fork length) and therefore may have even higher survival than Chinook salmon. Although the December-March period is expected to reflect the Delta outmigration timing of Sacramento-basin steelhead, real-time data from [CalFish](#) Track indicate that the majority of steelhead entrained at the salvage facilities have been and continue to be from the San Joaquin basin.
- iv. Mokelumne hatchery steelhead were released on 3/15/2024 at the Lighthouse Resort in the Delta to evaluate the effects of export on steelhead. To date, the majority of detections after release have occurred downstream at Benicia.
- v. Changes in water exports do not appear to significantly affect routing and survival of San Joaquin-origin steelhead (Buchanan et al. 2021), and there seems to be no export level that would lead to zero loss of steelhead this week (Table 6c). Nevertheless, reduction of export or a more positive OMR should generally lead to a reduction in salvage and loss (Table 6c, Figure 7). Agencies in SaMT acknowledge that any decrease in loss of steelhead would benefit the species.



5. If OMR is more negative than -5,000 cfs, are there changes in spawning, rearing, foraging, sheltering, or migration behavior beyond those anticipated to occur under OMR management at -5,000 cfs?
  - a. OMRI is expected to range between -3,500 and -5000 cfs for the next week. Under OMR flows more negative than -5,000 cfs the SaMT expects impacts to rearing, foraging, sheltering, or migration of salmonids present in the south Delta. Salmonid presence in the south Delta is difficult to assess because of limited observations and there is uncertainty in how much of the population might be impacted.

## Salmonid References

- Blankenship, S.M., J. Israel, E. Buttermore, and K. Reece. 2021. Knights Landing, California Department of Fish and Wildlife, Genetic Determination of Population of Origin 2017 through 2019 ver 1. Environmental Data Initiative.  
<https://doi.org/10.6073/pasta/85fbc988c0b1362e84c318e69c7a939e>
- Brandes PL, Pyper B, Banks M, Jacobson D, Garrison T, Cramer S. 2021. Comparison of Length-at-Date Criteria and Genetic Run Assignments for Juvenile Chinook Salmon Caught at Sacramento and Chipps Island in the Sacramento-San Joaquin Delta of California. *San Fr Estuary Watershed Sci.* 19(3).  
<https://doi.org/10.15447/SFEWS.2021V19ISS3ART2>
- Buchanan RA, Buttermore E, Israel J. 2021. Outmigration survival of a threatened steelhead population through a tidal estuary. *Can J Fish Aquat Sci.* 78(12):1869–1886.  
<https://doi.org/10.1139/cjfas-2020-0467>
- Buttermore, E., J. Israel, K. Reece, and S.M. Blankenship. 2021a. Sacramento trawl, Delta Juvenile Fish Monitoring Program, Genetic Determination of Population of Origin 2017-2021 ver 1. Environmental Data Initiative.  
<https://doi.org/10.6073/pasta/41983026f39bc11c329a18079dbca295>
- Buttermore, E., J. Israel, K. Reece, and S.M. Blankenship. 2021b. Chipps Island trawl, Delta Juvenile Fish Monitoring Program, Genetic Determination of Population of Origin 2017-2021 ver 1. Environmental Data Initiative.  
<https://doi.org/10.6073/pasta/f93fed9aa841ffa971aeded3872e0917>
- Buttermore, E., J. Israel, K. Reece, and S.M. Blankenship. 2021c. Central Valley Project, Genetic Determination of Population of Origin 2011-2021 ver 2. Environmental Data Initiative.  
<https://doi.org/10.6073/pasta/3561ee1976e9b6d2b023d1b7dce2450a>
- Eschenroeder J, Peterson M, Hellmair M, Pilger TJ, Demko D, Fuller A. 2022. Counting the Parts to Understand the Whole: Rethinking Monitoring of Steelhead in California’s Central Valley. *San Fr Estuary Watershed Sci.* 20(1).  
<https://doi.org/10.15447/sfews.2022v20iss1art2>

- Harvey BN, Jacobson DP, Banks MA. 2014. Quantifying the Uncertainty of a Juvenile Chinook Salmon Race Identification Method for a Mixed-Race Stock. *North Am J Fish Manag.* 34(6):1177–1186. <https://doi.org/10.1080/02755947.2014.951804>
- Interagency Ecological Program (IEP), L. Vance, and N. Kwan. 2023. Interagency Ecological Program: Fish catch and water quality data from the Sacramento River floodplain and tidal slough, collected by the Yolo Bypass Fish Monitoring Program, 1998-2023. ver 4. Environmental Data Initiative. <https://doi.org/10.6073/pasta/e2d248fcfaa8a1668b602d11984a5a2c>
- Johnson RC, Windell S, Brandes PL, Conrad JL, Ferguson J, Goertler PAL, Harvey BN, Heublein J, Israel JA, Kratville DW, et al. 2017. Science Advancements Key to Increasing Management Value of Life Stage Monitoring Networks for Endangered Sacramento River Winter-Run Chinook Salmon in California. *San Fr Estuary Watershed Sci.* 15(3). <https://doi.org/10.15447/sfew.2017v15iss3art1> A
- Michel CJ, Notch JJ, Cordoleani F, Ammann AJ, Danner EM. 2021. Nonlinear survival of imperiled fish informs managed flows in a highly modified river. *Ecosphere.* 12(5). <https://doi.org/10.1002/ecs2.3498>
- Reece, K., E. Buttermore, J. Israel, and S.M. Blankenship. 2021. State Water Project, Genetic Determination of Population of Origin 2011-2021 ver 1. Environmental Data Initiative. <https://doi.org/10.6073/pasta/f7793f0b265c99e1fe4a8875bca5682e>
- Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, Mangel M. 2010. State-dependent life history models in a changing (and regulated) environment: Steelhead in the California Central Valley. *Evol Appl.* 3(3):221–243. <https://doi.org/10.1111/j.1752-4571.2009.00103.x>
- Tillotson M, Hassrick J, Collins A, Phillis C. 2022. Machine Learning Forecasts to Reduce Risk of Entrainment Loss of Endangered Salmonids at Large-Scale Water Diversions in the Sacramento–San Joaquin Delta, California. *San Fr Estuary Watershed Sci.* 20(2):0–21. <https://doi.org/10.15447/sfew.2022v20iss2art3>

## Biology Distribution and Evaluation of Green Sturgeon

### Population Status

- Delta Life Stages:
  - Adults and Juveniles

### Distribution

#### *Current Distribution*

- Adults: Most abundant during spring spawning migration period of March through May, and post spawning out-migration periods May through June; October through January

depending on first winter storm event resulting in significant Sacramento River flow increases. Adult presence year-round to a lesser extent mainly in San Pablo Bay.

- Juveniles: Age-1 through Age-3 juveniles present year-round and widely distributed. Juveniles tagged with acoustic tags in the main channel Sacramento River near Sherman Island detected in the Sacramento River as far upstream as the Cache Slough complex, in the San Joaquin River at the Antioch Bridge, in Threemile, Horseshoe Bend, and Montezuma Sloughs. Seasonal abundance at the primary sampling site (near Sherman Island) appears to be highest during summer in based on capture and telemetry data. Residence time at the primary sampling site for individual fish ranges from one day to over one year but telemetry data show outmigration from the primary sampling site to the Pacific Ocean ranges from 27 to 552 days. Recent capture data shows diurnal depth preference in the main channel of the Sacramento River. No recent documentation of shallow water habitat presence or foraging.

### *Historical Trends*

- Juvenile and adult green sturgeon are historically present in the San Joaquin and Sacramento rivers and Delta.

### *Forecasted Distribution within Central Valley and Delta regions*

- Juvenile and adult green sturgeon are likely present in the San Joaquin and Sacramento rivers and Delta during the next week.

## Evaluation

1. Is there likely to be salvage that may exceed the annual loss limit?
  - a. Green sturgeon salvage is 0 fish (as of 6/10/2024). The agencies in the SaMT assessed the likelihood of salvage occurring in the next week is unlikely to occur.

## Biology, Distribution, and Evaluation of Delta Smelt

### Population Status

- Delta Smelt Life Stages:
  - Adults, Larvae, Juveniles
- Brood Year 2023:
- Abundance estimate:

- The most recent non-zero abundance estimate for postlarval/juvenile Delta Smelt is from the week of May 6, 2024 and is 3,558,101 (95% CI 1,380,606 to 7,590,408).
- Biological Conditions:
  - Delta Smelt are spawning and larval and juvenile Delta Smelt are present. Adult Delta Smelt have not been detected since 3/26/2024. Larval and juvenile Delta Smelt are expected to be present in the Sacramento River Deep Water Shipping Channel, Suisun Bay, Suisun Marsh, Lower Sacramento, and the South Delta based on the most recent survey detections. The Smelt Monitoring Team discussed the most recent monitoring data (TABLE 7) and considered published literature and professional judgement on the historical trends in regional distribution.

## Distribution

### *Current Distribution*

- Real time detection data are currently limited to EDSM, Chipps Island Trawl, and 20 mm Survey. Bay Study provides data as available.
- Two hundred and forty-two Delta Smelt (54 adult; 186 larvae, 2 juveniles) have been detected this water year (surveys and salvage). Forty-eight of these have been from experimental releases.
- Adult Delta Smelt have not been detected by surveys since 3/26/2024.
- Twenty-three larval Delta Smelt have been detected by surveys in Suisun Bay, Suisun Marsh, the Lower Sacramento, the Sacramento Deep Water Shipping Channel, Cache Slough/Liberty Island, and the Southern Delta starting 5/6/2024.
- One juvenile Delta Smelt was salvaged on 4/29/2024. No adult Delta Smelt have been salvaged after 2/23/2024. Cumulative seasonal salvage is 60.
- Qualitative Larval sampling began on 2/20/2024 at 0400 at the Tracy Fish Collection Facility (TFCF) and began at 1100 on 3/11/2024 at the Skinner Fish Facility (SFF).
- COA 8.5.2: Triggered 2/5/2024 by SLS survey 3 and implemented starting 2/7/2024; it was retriggered on 2/21/2024 by SLS survey 4, and on 3/4/2024 by SLS survey 5. It was not triggered by SLS survey 6 or 20 mm surveys 1-6. First larval detection was 3/22/2024.

Table 7. Summary of newly reported detections of Delta Smelt by Region and Salvage Facilities since the last assessment. Regions are those defined by EDSM sampling. Delta Smelt >58mm FL are considered adults. Subadult fish are considered by the SMT to be fish from the previous year’s cohort based on size and timing of collection. Young of year are considered juveniles and larvae. Salvage values reflect pre-expansion salvage.

Life Stage	North	South	West	Far West	Salvage
Adult	0	0	0	0	0
Subadult	0	0	0	0	0
Larvae/Juvenile	10	0	6	0	0

Table 8. Summary of recent Delta Smelt detections reported since last assessment and the total detections for the current water year. Notes reflect latest information on reported detections or completion of survey for the water year and include both larval and adult detections. Total Fish counts do not distinguish between hatchery origin and wild Delta Smelt. Table indicates new detections and previously reported detections that have undergone preliminary ID, QA/QC, and genetic confirmation. Numbers are updated as QA/QC and genetic confirmation become available. Tracy and Skinner facility counts reflect pre-expansion salvage.

Sampling Method	Frequency	New Detections	Preliminary Detections	QA/QC Detections	Genetically Confirmed to Date	Total WY2024	Notes
EDSM	Weekly	12	0	204	1	217	Ongoing. Phase 2 began on 4/1/2024 and will end 6/27/2024
SKT	Monthly	0	N/A	N/A	N/A	0	not occurring this year
SLS	Biweekly	0	N/A	N/A	N/A	0	Complete
20-mm	Biweekly	4	0	5	N/A	9	Ongoing
Summer Townet	Biweekly	0	N/A	N/A	N/A	0	Began 6/3/2024
Bay Study	Monthly	0	N/A	N/A	N/A	0	Ongoing
FMWT	Monthly	0	N/A	N/A	N/A	0	Complete
Chippis Island Trawl	Weekly	0	N/A	N/A	N/A	0	Ongoing
DJFMP	Monthly	0	N/A	1	N/A	1	Ongoing

Sampling Method	Frequency	New Detections	Preliminary Detections	QA/QC Detections	Genetically Confirmed to Date	Total WY2024	Notes
Electrofishing survey							
FCCL Brood Stock Collections	Weekly	0	N/A	N/A	N/A	0	Complete
LEPS	As available	0	N/A	N/A	N/A	0	Began 1/2/2024
FRP	Daily	0	N/A	N/A	N/A	0	Ongoing
Tracy Fish Collection Facility (CVP)	Daily	0	N/A	15	N/A	15	Ongoing
Skinner Fish Facility (SWP)	Daily	0	N/A	N/A	N/A	0	Ongoing
Total	N/A	N/A	N/A	N/A	N/A	242	Sum of all Delta Smelt observed during the water year

### ***Cultured Delta Smelt Experimental Releases***

- Experimental releases completed in Water Year 2024 include:
  - Release 1: 14,104 fish released at Sacramento River at Rio Vista (truck release)
  - Release 2: 13,089 fish released at Sacramento River at Rio Vista (hard and soft releases)
  - Release 3: 12,691 fish released at Sacramento River at Rio Vista (hard carboy and trailer release)
  - Release 4: 25,649 fish released at Sacramento River at Rio Vista (truck release)
  - Release 5: 12,778 fish released at Sacramento River at Rio Vista (hard and soft releases)
  - Release 6: 13,157 fish released at Sacramento River at Rio Vista (hard trailer and soft release)
- See additional details at: [SacPAS: Central Valley Prediction & Assessment of Salmon](#)

Table 10. Weekly summary of the origin of Delta Smelt. These identifications are considered tentative and additional genetic testing will confirm the identity of individuals. Individuals with no tags are provided alive to the FCCL as potential additions to the FCCL Broodstock.

Date	Survey	Stratum/Station	Total Caught	Ad. Clipped	VIE	No Tag
5/1/2024	20mm	Suisun Bay (Suisun Bay & Marsh)/411	4	N/A	N/A	X
5/7/2024	EDSM 20mm	Sac DW Ship Channel	7	N/A	N/A	X
5/9/2024	EDSM 20mm	Lower Sacramento	4	N/A	N/A	X
5/9/2024	EDSM 20mm	Suisun Marsh	1	N/A	N/A	X

**Historical Trends**

- Upstream migration for Delta Smelt occurs between September and December and in response to “first flush” conditions (Sommer et al. 2011, Grimaldo et al. 2009). Migration typically ranges one to four weeks after flow and turbidity increases, based on salvage data (Sommer et al. 2011).
- Historically, detections of ripe Delta Smelt began in January and peaked in February and March and the majority of Delta Smelt spawning occurs within a temperature range of 9-18°C (Damon et al. 2016).
- Based on historical monitoring data from the past few years (<https://github.com/Delta-Stewardship-Council/deltafish>), first detection of larvae in the Central and South Delta has typically occurred by mid to late March. ([https://www.cbr.washington.edu/sacramento/data/query\\_salvage\\_hrt.html](https://www.cbr.washington.edu/sacramento/data/query_salvage_hrt.html)).
- Salvage data as presented on SacPas indicates that adult Delta Smelt salvage in recent years has reached the 50th percentile at the end of February – beginning of March.
- Historically, the highest peak in salvage is in May and the second highest is in June (Grimaldo et al 2009).

**Forecasted Distribution within Central Valley and Delta regions**

- Predicting the distribution of Delta Smelt is currently difficult because detection data is limited to a few wild individuals and historic patterns may not be representative of the low population levels.
- The SMT uses turbidity as a surrogate for Delta Smelt presence and in making assessments of the likelihood of entrainment for larval Delta Smelt after spawning begins.

- The potential of experimentally released Delta Smelt to distribute from their release site is unknown at this time and SMT cannot predict their distribution beyond the original release site and subsequent recaptures. There is a high degree of uncertainty regarding the response of cultured fish to environmental cues typically applied to wild Delta Smelt.

## Abiotic Conditions

### *Turbidity*

- Sunny and hot, cooling slightly towards the end of the week. In Stockton, calm today and tomorrow with W wind 5-10 mph Wednesday night, gusting as high as 20mph. In Antioch, NW, W, WSW winds up to 16 mph, with gusts up to 24 mph through Thursday.
- Turbidity is below 12 FNU at OBI and at other stations in the central and south Delta.

Table 11. Relevant Environmental Factors to the current management actions for Delta Smelt.

Date Reported	Parameter	Value
5/27/2024-5/30/2024	20mm 12-station average Secchi Depth (m)	1.09 m
6/10/2024	Clifton Court Forebay (station CLC) 3-day average water temperature (°C)	24.1 °C

### *X2 Conditions*

- As of 6/10/2024, X2 was estimated to be 73 km.

### *Other Environmental Conditions*

- The Fish and Water Operation Outlook OMR Index values are expected to range between -3,500 to -5,000 cfs this week.
- QWEST was +2,410 cfs as of 6/10/2024 and is estimated to range between -2,000 to +2,500 cfs
- Real time tracking of environmental conditions, relevant thresholds and Delta Smelt catch data are updated daily at: [SacPAS: Central Valley Prediction & Assessment of Salmon](#)
- Water temperature was 22.5°C at Rio Vista and 22.4°C at Antioch as of 6/10/2024.
- Anticipated Vernalis Flows: 2,500 to 3,200 cfs

## Evaluation

### USBR and DWR Proposed Operations:

- Reclamation and DWR propose joint water project operations be limited to a 14-day average OMRI flow no more negative than -5,000 cfs.



## Questions and Discussions:

1. Between December 1 and January 31, has any first flush condition been exceeded?
  - a. First flush conditions were exceeded on 1/21/2024. Integrated Early Winter Pulse Protection (IEWPP) was implemented from 01/23/2024 through 2/5/2024.
2. Do DSM have a high risk of migration and dispersal into areas at high risk of future entrainment? (December 1- January 31)
  - a. No longer relevant.
3. Has a spent female been collected?
  - a. Neither a ripe nor spent female were collected before April 1, 2024.
4. If OMR of -2000 cfs does not reduce OBI turbidity below 12 NTU/FNU, what OMR target is deemed protective between -2000 and -5000 cfs?
  - a. No longer relevant.
5. If OBI is 12 NTU/FNU, what do other station locations show?
  - a. No longer relevant.
6. If OBI is 12 NTU/FNU, is a turbidity bridge avoidance action not warranted? What is the supporting information?
  - a. No longer relevant.
7. After March 15 and if QWEST is negative, are larval or juvenile DSM within the entrainment zone of the CVP and SWP pumps based on surveys?
  - a. QWEST will remain positive when the Delta Cross Channel (DCC) Gates are open and becomes negative when DCC Gates are closed. One larval DSM was detected in the South Delta on 5/13, so DSM may be in the entrainment zone, but the majority of fish have been detected outside of the South and Central Delta.
  - b. Due to following the 2024 IOP, we are also following COA 8.5.2. Temperature became suitable for spawning on 1/31/2024 ( $> 12^{\circ}\text{C}$  three-day average at SJJ; Damon et al. 2016). One larval DSM was detected in the Southern Delta on 5/13, so DSM may be in the entrainment zone, but the majority of fish have been detected outside of the South and Central Delta.

Mean Secchi depth for South Delta stations is > 1 m as of 5/27/2024-5/30/2024.

8. Based on real-time spatial distribution of Delta Smelt and currently available turbidity information, should OMR be managed to no more negative than -3,500?
  - a. No. Water temperatures became suitable for spawning on 1/31/2024, and QWEST may be positive or negative this week, depending on DCC Gates. However mean Secchi depth for South Delta stations was >1 m as of 5/27/2024-5/30/2024, thus COA 8.5.2 and PA protections were not triggered by the most recent 20mm surveys (20 mm 6). One larval DSM was detected in the Southern Delta on 5/13, but the majority of fish have been detected outside of the South and Central Delta.
9. What do hydrodynamic models, informed by EDSM or other relevant data, suggest the estimated percentage of larval and juvenile DSM that could be entrained may be?
  - a. No PTM runs were requested by the SMT in advance of the SMT meeting. Flows at Vernalis have decreased this week. QWEST will remain positive when the Delta Cross Channel (DCC) Gates are open and becomes negative when DCC Gates are closed. The Fish and Water Operation Outlook OMR Index values are expected to range between -3,500 to -5,000 cfs this week.
  - b. The majority of fish have been detected outside of the South and Central Delta, and turbidity is low in the Central and South Delta. Thus, we continue to expect low entrainment risk between OMRI range of -3,500 to -5,000 cfs.

## Delta Smelt References

- Damon, L. J., S. B. Slater, R. D. Baxter, and R. W. Fujimura. 2016. Fecundity and reproductive potential of wild female Delta smelt in the upper San Francisco Estuary, California. *California Fish and Game* 102(4):188–210.
- Hobbs, J. A., Lewis, L. S., Willmes, M., Denney, C., & Bush, E. (2019). Complex life histories discovered in a critically endangered fish. *Scientific Reports*, 9(1).  
<https://doi.org/10.1038/s41598-019-52273-8>
- Grimaldo, L. F., T. Sommer, N. Van Ark, G. Jones, E. Holland, P. B. Moyle, B. Herbold & P. Smith (2009) Factors Affecting Fish Entrainment into Massive Water Diversions in a Tidal Freshwater Estuary: Can Fish Losses be Managed? *North American Journal of Fisheries Management*, 29:5, 1253-1270, DOI: 10.1577/M08-062.1
- Gross, E. S. (2021). Modeling Delta Smelt Distribution for Hypothesized Swimming Behaviors. *San Francisco Estuary and Watershed Science*, 19(1).

- Kimmerer, W. J. (2008). Losses of Sacramento River Chinook Salmon and Delta Smelt to Entrainment in Water Diversions in the Sacramento-San Joaquin Delta. *San Francisco Estuary and Watershed Science*, 6(2).
- Polansky, L., Newman, K.B., Nobriga, M.L. et al. Spatiotemporal Models of an Estuarine Fish Species to Identify Patterns and Factors Impacting Their Distribution and Abundance. *Estuaries and Coasts* 41, 572–581 (2018). )
- Smith, W. E., Polansky, L., and M. L Nobriga. 2021. Disentangling risks to an endangered fish: using a state-space life cycle model to separate natural mortality from anthropogenic losses. *Canadian Journal of Fisheries and Aquatic Sciences*, 78: 1008-1029.
- Sommer, T., F. Mejia, M. Nobriga, and L. Grimaldo. 2011. The Spawning Migration of Delta Smelt in the Upper San Francisco Estuary. *San Francisco Estuary and Watershed Science* 9(2).

## Attachment 1:

### Winter-run Chinook Salmon Historical Data

Below are a series figures of containing historical information on Winter-run Chinook Salmon.




Figure A1. Historic juvenile production estimates (JPEs) from brood years 1992 through 2023. Method 2 (O'Farrell et al 2018) used from 2019 onward.

Figure A1 is a bar graph that shows winter-run Chinook historic juvenile production estimates from brood years 1992 through 2023. The years 2005 and 2006 show the highest estimates at roughly 3,800,000 and 3,750,000 respectively.

Genetic vs LAD Historical Loss Comparison




Figure A2. Comparison of the of the loss between annual loss calculated from genetically confirmed loss versus LAD only loss calculation. Break between 7000 and 13000 used to accommodate high loss values observed in Water Year 2001. Genetic loss through 3/21/2024.

Figure A2 is a bar graph that compares genetic and LAD historical loss from 1996 through March 21, 2024.




Figure A3. Cumulative loss of genetic winter-run as a percentage of the Juvenile Production Estimate for each Water Year. Inset graph represents all years and main graph excludes 2001 which is an order of magnitude greater than all other years. Only those years that are greater than 2024 are labeled.

Figure A3 is a line graph that shows cumulative loss of genetic winter-run as a percent of the Juvenile Production Estimate for each Water Year. The years that are greater than the percentage in 2024 are labeled, and include: 1996, 1998, 1999, 2000, 2003, 2004, 2008, 2009, 2010, 2011, 2012. An inset map shows the percentage in 2001 which is an order of magnitude greater than all other years.

## Spring-run Chinook Salmon Historical Data

Table A1. Historical loss of LAD natural-origin spring-run from WY 2010 to WY 2023. Wet years were highlighted in bold. WY 2024 is spring-run loss data through 5/10/24.

Water Year	Total Loss	50% of Total Loss:	75% of Total Loss:	90% of Total Loss:	100% of Total Loss:	Water Year Type
2010	6,082.20	5/4/2010	5/16/2010	5/27/2010	6/5/2010	Below Normal
2011	52,504.32	5/8/2011	5/16/2011	5/30/2011	6/24/2011	Wet
2012	2,394.27	4/17/2012	4/21/2012	5/2/2012	6/8/2012	Below Normal
2013	2,495.92	4/22/2013	5/1/2013	5/11/2013	5/25/2013	Dry
2014	348.72	4/9/2014	4/19/2014	4/23/2014	5/10/2014	Critically Dry
2015	70.02	4/22/2015	4/23/2015	5/4/2015	5/18/2015	Critically Dry
2016	297.79	4/27/2016	5/2/2016	5/14/2016	5/19/2016	Below Normal
2017	72,011.18	5/11/2017	5/15/2017	5/23/2017	6/29/2017	Wet
2018	18,313.05	5/8/2018	5/19/2018	6/3/2018	5/23/2018	Below Normal
2019	6,100.44	5/6/2019	5/19/2019	5/20/2019	6/25/2019	Wet
2020	4,167.11	4/21/2020	4/24/2020	4/30/2020	5/26/2020	Dry
2021	517.99	4/27/2021	5/1/2021	5/4/2021	5/12/2021	Critically Dry
2022	552.55	4/28/2022	5/2/2022	5/12/2022	5/21/2022	Critically Dry
2023	10,191.83	5/20/2023	5/29/2023	6/2/2023	7/1/2023	Wet
2024	11,648.78 (partial)	N/A	N/A	N/A	N/A	Above Normal

Table A2. Average loss of LAD natural-origin spring-run in dry/critically dry years, normal years, and wet years.

Average Loss	50% of Total Loss:	75% of Total Loss:	90% of Total Loss:	100% of Total Loss:
Average Loss in Dry/Critically Dry Years	4/22	4/27	5/5	5/18
Average Loss in Below Normal Years	4/29	5/7	5/19	5/29
Average Loss in Wet Years	5/11	5/19	5/26	6/27