



— BUREAU OF —
RECLAMATION

Weekly Fish and Water Operations Outlook

6/3/2025 – 6/9/2025

Water Project Operational Intent for Week

The D-1641 standards for CVP/SWP operations in June include X2 Delta Outflow requirements and water quality standards to protect the diversion of irrigation water from the Delta.

SWP and CVP exports are governed by SWRCB's 0.35 E/I ratio and applicable OMR constraints under ESA.

Biological Context

Water temperatures in the Delta are increasing. OMR shall not be more negative than –5,000 cfs, unless more restrictive OMR thresholds are triggered by other conditions.

Forecasted Weather

Warm and dry conditions continue this week. Periodically breezy winds combined with low humidity result in Red Flag Warnings for the Northern and Central Sacramento Valley and Coastal Range.

Tables

Table 1: Anticipated weekly operational ranges by tributary. Environmental and fish conditions updated by respective watershed groups at varying intervals that may not coincide with the weekly range of Water Operations

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Clear Creek	<ul style="list-style-type: none"> • Current Release: 800 cfs • Anticipated Weekly Range of Releases: 200 cfs to 800 cfs 	<ul style="list-style-type: none"> • Adult spring-run Chinook Salmon are migrating to the creek. • Spring, fall, and late fall-run Chinook Salmon fry are rearing and emigrating. • O. mykiss/steelhead are emerging and migrating downstream. <p>(Updated 5/13/2025)</p>
Sacramento River	<ul style="list-style-type: none"> • Shasta Storage: 4.143 MAF • Current Release: 8,500 cfs • Anticipated Weekly Range of Releases: 8,500 cfs. 	<ul style="list-style-type: none"> • Spring-run Chinook adults are migrating upstream and holding. • Winter-run Chinook adults are beginning to spawn and have been holding in the system some months now. • Late fall-run Chinook fry are rearing and emigrating downstream. • Spring and Fall-run Chinook Salmon smolts are emigrating with hatchery released fall-run Chinook. • O. mykiss/steelhead eggs incubating and hatching and fry are rearing and emigrating. • White sturgeon are currently spawning and larvae are moving downstream. • Larval Green Sturgeon are hatching (as observed by USFWS) and redistributing from spawning and incubation areas in low numbers <p>(Updated 6/2/2025)</p>

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Feather River	<ul style="list-style-type: none"> • Oroville Storage: 3.418 MAF • Current Release: 2,100 cfs • Anticipated Weekly Range of Releases: 1,800 cfs to 3,000 cfs • Daily temperature maximum: 55 +/- 4 degrees F at Fish Hatchery 	<ul style="list-style-type: none"> • Spring-run Chinook Salmon juveniles are migrating downstream. • Spring-run Chinook Salmon adults are migrating upstream and holding. • Fall-run Chinook Salmon juveniles are migrating downstream. • <i>O. mykiss</i> are emerging and migrating downstream. • Sturgeon have been detected in multiple locations throughout the river. White and Green Sturgeon are currently in spawning season. <p>(Updated 6/2/2025)</p>
American River	<ul style="list-style-type: none"> • Folsom Storage: 914 TAF • Current Release: 2,500 cfs • Anticipated Weekly Range of Releases: 2,500 cfs 	<ul style="list-style-type: none"> • Fall-run Chinook Salmon Fry are migrating downstream in low numbers. Most have already exited the system. <p>(Updated 6/2/2025)</p>
Stanislaus River	<ul style="list-style-type: none"> • New Melones Storage: 1.912 MAF • Current Release: 1,700 cfs • Anticipated Range of Weekly Releases: 1,700 cfs 	<ul style="list-style-type: none"> • Juvenile and adult <i>O. mykiss</i> are present. • <i>O. mykiss</i> are spawning and fry are emerging • Fall-run Chinook fry are migrating downstream. <p>(Updated 6/2/2025)</p>

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Delta	<ul style="list-style-type: none"> • Freeport: 13,000 to 17,000 cfs • Vernalis: 1,400 to 2,400 cfs • Delta Outflow index: 7,500 to 11,000 cfs • Combined Exports: 5,200 to 6,500 cfs • JPP: 2,700 to 3,200 cfs • CCF: 2,100 cfs to 3,300 cfs • Expected Daily OMR Index Values: -4,700 to -5,100 cfs • DCC Gates: Closed Monday through Thursday. Open Friday through Sunday • X2 = 73 km • Tides: Transition from Neap to Spring; First Quarter Moon on June 2, 2025 	<ul style="list-style-type: none"> • YOY Chinook Salmon are migrating through the Delta and exiting the system. • Adult Delta smelt were last detected by EDSM on 3/17/25 in the SDWSC. 20-mm survey has detected three larval Delta smelt in the SDWSC and Central and South Delta. The most recent detections were on 4/14/25. • A total of 124,946 individual adult Delta smelt were released in WY2025. So far, there have been 79 confirmed detections of cultured Delta smelt. • Cumulative adult Delta smelt salvage is 17. One larval Delta smelt was detected in larval sampling at TFCF on 4/16/25. • Larval white sturgeon have been detected in the Sacramento River and Miner Slough. • Larval longfin smelt have been detected in the Central and South Delta, the Sacramento River, Suisun Marsh, Suisun Bay, the Confluence, Carquinez Strait, and San Pablo Bay. • Juvenile longfin smelt have been detected in South and Central San Francisco Bay, San Pablo Bay, Suisun Marsh, and Suisun Bay. • Adult longfin smelt were last detected May 7 at Chipps Island. • Adult, juvenile, and larval LFS have been detected in salvage. Cumulative adult LFS salvage = 8 and cumulative juvenile LFS salvage = 204. <p>(Updated 6/2/2025)</p>

Table 2: WY 2025 Salmonid Current Loss and Delta Smelt Abiotic Conditions. Additional Real-Time OMR Restrictions and Performance Objectives (4.10.5.10.2, 4.10.5.10.3) and Onset of OMR Management (4.10.5.10.1). Genetic identification of salmon is not used in calculating loss, but results are included in the Assessment as they become available.

Species/run	Threshold	Current Status	Weekly Trend	Updated
Green sturgeon	WY 2025 salvage = 74	WY 2025 salvage = 1 (1.3%)	Occasional salvage possible	5/27/25
Natural winter-run Chinook Salmon (JPE= 98,982)	Incidental Take Limit= 554 Annual thresholds 50%= 277 fish 75%= 415 fish 100%= 554 fish	Loss= 28.82 (5.2% of threshold) 7-day rolling sum as of 6/02/25 = 0.00	Salvage is unlikely in the upcoming week.	6/2/25
Natural Steelhead	100% threshold = 3,000	WY 2025 loss = 600 (20% of threshold) as of 6/02/25	Salvage possible at reduced rates	6/2/25
Steelhead Weekly Loss Threshold	7-day rolling sum of steelhead salvage exceeds loss of 120 fish	No exceedances – 7 day rolling sum as of 6/02/25 = 2.72	Salvage possible at reduced rates	6/2/25
Sacramento River Hatchery winter-run Chinook salmon (JPE= 135,342)	Annual thresholds 50%= 81 fish 75%= 122 fish 100%= 162 fish	Loss = 216.58 (133%) 50%= exceeded 3/18 75%= exceeded 3/19 100%= exceeded 3/22	Salvage is unlikely in the upcoming week.	4/15/25
Battle Creek Hatchery winter-run Chinook salmon	JPE = 2,868	Loss = 0 (0%)	Released on 4/16/25.	4/21/25
Proposed Action Hatchery yearling spring-run Chinook salmon surrogates	See Table 3a	See Table 3a	See Table 3a	4/7/25
Delta Smelt	See Table 3b	See Table 3b	See Table 3b	1/06/2025
Longfin Smelt	See Table 3c	See Table 3c	See Table 3c	1/06/2025

Table 3a-d: Relevant Water Year 2023 Fish Criteria and Status for Listed Fish under the SWP Long-Term Incidental Take Permit.

Table 3a: Chinook Salmon

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Onset of OMR Management (8.3)	Jan. 1 - Jun. 30	In effect	Begins January 1 or earlier if COA 8.3.1, COA 8.3.2, or COA 8.3.3 are in effect (see Table 3b)	N/A	N/A	3/10/25	N/A
Winter-run Annual Loss (8.4.3)	July 1 - Jun. 30	In effect	-Natural-origin Winter-run Loss Threshold: 0.5% of JPE (494.47) -Hatchery-origin Winter-run Loss Threshold: 0.12% of JPE (162.41) -Battle Creek Loss threshold: 0.12% of JPE (3.44)	Confirmed Genetic WR Annual Loss = 28.82 Hatchery origin Winter-run Loss = 216.58 Battle Creek Winter-run Loss = 0	Unlikely to observe salvage of natural or hatchery - origin winter-run based on historical salvage.	6/2/25	50%, 75%, and 100% of the LSNFH loss threshold was hit on 3/18/25, 3/19/25, and 3/22/25 respectively. 117,225 BY 2024 WR was released in Battle Creek on 4/16/25.
Natural-origin Winter-run Early Season Weekly Loss Thresholds (8.2.1)	Nov. 1- Dec. 31	Not in Effect	N/A	N/A	N/A	2/4/25	N/A

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Natural-origin Winter-run Weekly Loss (8.4.4)	Jan 1 – June 30	In effect	Thresholds based on Table 4, Column E of 2024 SWP ITP: [Annual Loss Threshold (based on JPE surrogate) x 50% of Annual Loss Threshold x Winter-run in Delta (based on Column E)]	4/2/25-6/30/25 Threshold: 0	Based on salvage thru 6/1/25 7-day LAD loss: 0.00 Total loss of 7 day rolling sum (includes genetically confirmed): 0	6/2/25	No LAD or Genetically confirmed Older Juvenile has been salvaged at Delta fish facility

Spring-run Protection Action and Surrogate Annual Loss (8.4.5)	Natural-origin: Oct. – June 30 Hatchery-origin: Nov. 1 – June 30	Natural-origin=In effect Hatchery-origin: In effect	Yearling spring run surrogates: Group 1: 1,747.23 (0.25% of 698,892 fish released) Group 2: 193.39 (0.25% of 77,355 fish released) Group 3: 186.10 (0.25% of 74,725) YOY spring run surrogates: Group 1: 1,191.85 (0.25% of 476,741) Group 2: 1,189.58 (0.25% of 475,831) Group 3: 1,260.76 (0.25% of 504,304) Group 4: 1,737.93 (0.25% of 695,170) Group 5: 940.31	Yearling spring-run Surrogates: Current Loss for Group #1 through 5/18/25: 1,050.61 (60.13% of the loss threshold) Current Loss for Group #2 through 5/18/25: 72.52 (37.50% of the loss threshold) Current Loss for Group #3 through 5/18/25: 43.33 (23.28% of the loss threshold) YOY Spring-run Surrogates: Current loss for FR YOY spring-run surrogate Group 1: 16.56 (1.39 % of the loss threshold) Current loss for FR YOY spring-run surrogate Group 2: 110.14 (9.26 % of the loss threshold) Current loss FR YOY spring-run surrogate	Unlikely to see salvage in the upcoming week	Yearling Group 1, 2 & 3 and YOY Group 1, 2, 3, 4, 5, 6 updated through 6/1/25 salvage data for SWP and CVP.	No loss from any spring run surrogate group occurred last week.
--	---	--	---	--	--	---	---

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
			(0.25% of 376,122) Group 6: 920.22 (0.25% of 368,085)	Group 3: 0 Current loss CNFH YOY spring-run surrogate Group 4: 0 Current loss FR YOY spring-run surrogate Group 5: 0 Current loss FR YOY spring-run surrogate Group 6: 0			

Table 3b: Delta Smelt

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
First Flush Action (8.3.1)	Dec. 1 – last day of February	Off ramped Action triggered on Dec. 16, implemented from Dec. 19 through Jan 1, 2025	- three-day Freeport (FPT) daily flow running avg >= 25,000 AND [three-day Freeport turbidity running avg >=50 NTU OR Smelt Monitoring Team recommendation]	FPT 3-day avg. Flow = Not relevant Turbidity = Not relevant	N/A	1/6/2025	N/A
Adult Delta Smelt Entrainment Protection ("Turbidity Bridge Avoidance") (8.3.2)	After IEWPP or Dec. 20 until 3-day average temperatures at Jersey Point (SJJ) or Rio Vista (RVB) exceed 12 °C (53.6 °F)	Not active; offramped as of 2/25/25	Occurs after the Integrated Early Winter Pulse protection or December 20 (whichever comes first) until 3-day average temperature offramp at Jersey Point (SJJ) or Rio Vista (RVB) > 12 °C (53.6 °F) -OBI, OSJ, and HOL turbidity > 12 FNU -Vernalis flow > 10,000 cfs (temporary offramp); <8,000 cfs (reinstated)	OSJ Turbidity = Not relevant HOL Turbidity = Not relevant OBI Turbidity = Not relevant 3-d SJJ temp = Not relevant 3-d RVB temp = Not relevant Vernalis Flow = Not relevant	N/A	2/25/25	N/A

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Larval and Juvenile Delta smelt Protection (8.4.1)	After Adult Delta smelt Entrainment Protection ends	Active as of 2/25/25. Not triggered.	SLS/20mm Secchi depth for 12 south delta stations \leq 1m -Rio Vista flows $>55,0000$ cfs or Vernalis flows $>8,000$ cfs (temporary offramp); $<40,000$ cfs (Rio Vista) or $<5,000$ (Vernalis) action reinstated	Secchi depth = 141 cm (20-mm Survey 6) Rio Vista flows = 9,200 cfs Vernalis flows = 2,400 cfs	N/A	6/1/25	N/A

Table 3c: Longfin Smelt

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Adult LFS Protection (8.3.3)	Dec. 1 - end of February	Not active	-Cum. salvage $> (\text{Age } 1+ \text{LFS Index}/20) + 1 = 181$ fish	Cum LFS salvage greater than 60mm = Not relevant	No change expected	3/17/25	N/A

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Larval and Juvenile Longfin Smelt Entrainment Protection (8.4.2)	Jan. 1 – Jun. 30	Active; not triggered. Triggered on 1/19 and 1/28. Implemented 1/20-1/26	-7-day average QWEST < +1,500 cfs, AND LFS larvae or juveniles in most recent SLS or 20 mm survey at 809 & 812 > 50; OR cumulative salvage > 50 or 75% avg annual salvage 2009-present -Rio Vista flows > 55,000 cfs or Vernalis flows > 8,000 cfs (temporary offramp); < 40,000 cfs (Rio Vista) or < 5,000 (Vernalis) reinstated	7-day average QWEST = +2,100 cfs Larval/juvenile (>20mm) 809 + 812 catch (20-mm 6) = 0 Cumulative juvenile (>20mm) salvage = 204 Rio Vista flows = 9,200cfs Vernalis flows = 2,400 cfs	N/A	6/2/25	N/A

Table 3d: White Sturgeon

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
White Sturgeon Entrainment Protection Action (8.4.7)	Year-round	Active; not triggered Flow Conditions: Not met Survey Conditions: Met	-YOY WS detected in one of the listed north or central Delta survey stations in the last 90 days - Mean total exports for the last 90 days $\geq 14,296.76 + (-0.41) \times (90\text{-day average Vernalis flow})$	YOY WS detections= 20-mm station 707 on 4/3/25 90-Day Avg Vernalis flows = 2,409 cfs 90-Day Avg Exports = 4,918 cfs	More YOY detections possible Flow/ Exports conditions unlikely to meet criterion	5/12/25	Survey Conditions met until 7/2/2025 WY 2025 salvage = 4

Table 3e: OMR

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
OMR Storm Flex (8.5)	Start of OMR – Onramp of Larval and Juvenile DS Protection Action (8.4.1) or last day of February (whichever occurs first)	Off-ramped as of 2/25/25	-Delta is in excess -QWEST is > +1,500 cfs -X2 is < 81 km - Daily average turbidity at OSJ, HOL, and OBI are <12 FNU -Higher level of outflow available for diversion due to storm flows -Measurable amount of precipitation has occurred -None of COA's are controlling operations (8.2.1, 8.3.2, 8.3.3, 8.4.2, 8.4.3, 8.4.4, 8.4.5, 8.4.7) -Cumulative loss at CVP and SWP of yearling CNFH LFR Chinook salmon (as yearling CHNSR surrogates) is < 0.5% with any of the release groups	QWEST=Not relevant X2 = Not relevant Turbidity = Not relevant No relevant salmon loss threshold exceedances (see Tables 2a & 3a) No COA's are controlling operations	N/A	2/25/25	N/A

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
End of OMR Management (8.6)	Jun. 1 – Jun. 30	In effect	<p>Smelt: -Daily mean water temperature at Clifton Court Forebay (CLC) is > or equal to 25 C for 3 consecutive days</p> <p>Salmonids: -Daily mean water temperature is > 22.2 C at Mossdale and Prisoners Point for 7 days (can be non-consecutive).</p>	<p>Smelt: CLC temperatures (°C) 5/31-6/2: 23.5, 23.8, 23.6 N/A Salmonids: 1 Day at Prisoners Point (6/1)</p>	N/A	6/2/2025	N/A
Spring Outflow (COA 8.12.1)	April 1 – May 31	Not in effect	<p>Critical year: ratio of Vernalis flow to SWP and CVP combined exports shall be 1 to 1. Dry year: ratio of Vernalis flow to SWP and CVP combined exports shall be 2 to 1. Below Normal year: ratio of Vernalis flow to SWP and CVP combined exports shall be 3 to 1. Above Normal/Wet year: ratio of Vernalis flow to SWP and CVP combined exports shall be 4 to 1</p>	<p>High flow offramp (Delta Outflow greater than 44,500 cfs) was in effect from 4/1/25 - 4/9/25</p>	N/A	4/22/25	N/A

Table 4: Fish monitoring gear efficiency and disruptions. Status Categories: [1] Active (ongoing sampling), [2] Partial Interruption (some sampling interruptions), [3] Interrupted (sampling fully suspended), [4] Not Active (sampling not scheduled)

Monitoring survey	Region	Notes (as of 6/2/2025)	Status
SWP regular counts, CWT reading	Delta	Active	1
SWP larval sampling	Delta	Inactive since 5/30	4
CVP regular counts, CWT reading	Delta	Active	1
CVP larval sampling	Delta	Inactive since 5/30	4
Smelt Larval Survey	Delta	Not Active	4
LES	Delta	Not Active	4
20mm Survey	Delta	Active	1
Fall Mid-water Trawl	Delta	Not Active	4
Summer Townet Survey	Delta	Active (as of 6/2/25)	1
Bay Study	Delta	Active	1
DJFMP- Chipps and Sacramento Trawls	Delta	Active	1
DJFMP- Seines	Delta	Active	1
EDSM	Delta	Not Active	4
EMP	Delta	Active	1
Mossdale	Delta	Active (CDFW)	1
USGS Flow monitoring	Delta	Active	1
Red Bluff Diversion Dam Rotary Screw Trap (RST)	Sacramento River	Active	1
Knights Landing RST	Sacramento River	Not Active	4
Tisdale RST	Sacramento River	Not Active	4
GCID RST	Sacramento River	Not Active	4
Mill Creek RST	Mill Creek	Active	1
Deer Creek RST	Deer Creek	Not Active	4
Yuba River (Hallwood) RST	Yuba River	Active	1
Butte Creek Carcass Surveys	Butte Creek	Not Active	4
Butte Creek RST	Butte Creek	Not Active	4
Redd dewatering and stranding surveys	Sacramento River	Active	1

Monitoring survey	Region	Notes (as of 6/2/2025)	Status
Sacramento Carcass and Redd Surveys (winter-run Chinook Salmon)	Sacramento River	Active	1
Lower Sacramento RST	Sacramento River	Not Active	4
Feather River (upper DWR) RST	Sacramento River	Not Active	4
Feather River (lower CDFW) RST	Sacramento River	Active	1
Feather River Carcass Survey (fall-run Chinook Salmon)	Sacramento River	Not Active	4
Sonar, telemetry (sturgeon)	Feather River	Active	1
Egg mats (sturgeon)	Feather River	Active	1
SJRRP CDFW Field Monitoring	San Joaquin River	Active	1
SJRRP USFWS and USBR Field Monitoring	San Joaquin River	Active	1
Stanislaus Fish Weir	San Joaquin River	Active	1
Stanislaus River Carcass Survey (steelhead)	San Joaquin River	Active	1
American River Carcass Survey	Sacramento River	Not Active	4

Preference (i.e., a y-intercept of 0.5)