

1 **Appendix 9B**2 **Aquatic Species Life History Accounts**

3 This appendix provides additional information on the life history characteristics of
 4 the target aquatic species assessed in the Remanded Biological Opinions on the
 5 Coordinated Long-Term Operation of the Central Valley Project (CVP) and State
 6 Water Project (SWP) Environmental Impact Statement (EIS). This information is
 7 intended to provide a more holistic understanding of how these species use the
 8 water bodies influenced by operation of the CVP and SWP and to help clarify
 9 relationships that provide the logical foundation for conclusions regarding the
 10 potential environmental consequences associated with changes in operation.

11 This appendix addresses the following species:

- 12 • River Lamprey
- 13 • Pacific Lamprey
- 14 • Green Sturgeon
- 15 • White Sturgeon
- 16 • Chinook Salmon
 - 17 – Winter-run Chinook Salmon
 - 18 – Central Valley Spring-run Chinook Salmon
 - 19 – Central Valley Fall-run and Late Fall-run Chinook Salmon
 - 20 – Upper Klamath and Trinity Rivers Spring-run Chinook Salmon
- 21 • Central Valley Steelhead
- 22 • Klamath Mountains Province Steelhead
- 23 • Sacramento Splittail
- 24 • Longfin Smelt
- 25 • American Shad
- 26 • Eulachon
- 27 • Striped Bass
- 28 • Southern Resident Killer Whale

29 **9B.1 River Lamprey (*Lampetra ayresii*)**30 **9B.1.1 Legal Status**

31 Federal: None

32 State: Species of Special Concern

33 River Lamprey was petitioned for listing by a number of conservation groups in
 34 2003, along with three other lamprey species (Klamath-Siskiyou Wildlands
 35 Center et al. 2003). The petition was declined by the U.S. Fish and Wildlife
 36 Service (USFWS) in 2004 because of insufficient evidence that listing was
 37 warranted.

9B.1.2 Distribution

1 River Lamprey are found in large coastal streams from just north of Juneau,
2 Alaska, to the San Francisco Bay (Vladykov and Follett 1958, Wydoski and
3 Whitney 1979). The Sacramento and San Joaquin basins are at the southern edge
4 of their range (Moyle et al. 2009). Little is known regarding their abundance and
5 distribution within California; they seem to be primarily associated with the lower
6 portions of certain large river systems, and most records for the state are from the
7 lower Sacramento-San Joaquin system, especially the Stanislaus and Tuolumne
8 rivers (Moyle et al. 1989, Moyle 2002). In the Sacramento River, they have been
9 documented upstream to at least Red Bluff Diversion Dam (RBDD) (Hanni et al.
10 2006, Moyle et al. 2009). River Lamprey have also been collected in the Feather
11 River, American River, Mill and Cache creeks (Vladykov and Follett 1958, Hanni
12 et al. 2006, Moyle et al. 2009). River Lamprey have not been documented during
13 rotary screw trapping efforts in Clear, Battle, and Deer creeks, or in the Yuba
14 River (Hanni et al. 2006). Other streams where they have been found in
15 California outside of the Central Valley include the Napa and Russian rivers, and
16 Alameda, Sonoma, and Salmon creeks (DWR et al. 2013).
17

9B.1.3 Life History and Habitat Requirements

18 River Lamprey are a small parasitic anadromous species. Most studies of their
19 biology have been conducted in British Columbia; relatively little is known
20 regarding their life history and habitat requirements in California (Moyle 2002).
21
22 Adult River Lamprey migrate from the ocean into spawning areas in the fall.
23 Adults of both sexes construct nests in gravel at the upstream end of riffles
24 (Wydoski and Whitney 1979, Beamish and Youson 1987, Moyle 2002). Eggs are
25 deposited and fertilized in these depressions, after which the adults typically die,
26 similar to other species of lampreys. In the Sacramento-San Joaquin basin of
27 California, most spawning is believed to occur in April and May (Vladykov and
28 Follett 1958; Scott and Crossman 1973) at temperatures of about 55 to 56 degrees
29 Fahrenheit (°F) (Wang 1986). Two females in Cache Creek were reported to have
30 11,400 and 37,300 eggs each (Vladykov and Follett 1958).
31
32 After hatching, young ammocoetes (the larval stage of lamprey) drift downstream
33 to settle in the silt-sand substrates of backwaters, eddies, and pools, where they
34 remain burrowed for approximately 3 to 5 years (Moyle 2002). At this stage, they
35 are filter feeders, with a diet consisting of algae (primarily diatoms) and other
36 organic detritus and microorganisms (Wydoski and Whitney 1979). Good water
37 quality and temperatures not exceeding 77°F are believed to be necessary for their
38 survival (Moyle 2002). Their metamorphosis into adults begins in July when they
39 reach about 12 centimeters (cm) (4.7 in) (Beamish 1980), and is not complete for
40 about 9 to 10 months until around April the following spring, when the esophagus
41 opens and adults are able to osmoregulate (Beamish and Youson 1987, Moyle
42 2002). This is a more extended period of metamorphosis than observed in other
43 lamprey species. During this time, they are believed to live in deep waters of the
44 river channel. Just prior to the completion of metamorphosis, the juvenile
45 lampreys (macrophthalmia) congregate immediately upstream of salt water and
enter the estuary or ocean from May to July (Beamish and Youson 1987).

1 Adults spend 3 to 4 months in salt water, remaining close to shore and growing to
2 lengths of about 25 to 31 cm. In the estuary or ocean, River Lamprey are obligate
3 parasites, typically killing their host in the process of feeding. They most
4 commonly parasitize fishes 10 to 30 cm long, feeding near the surface on smelt,
5 herring, and mid-size salmonids (Beamish 1980, Roos et al. 1973, Beamish and
6 Neville 1995). In Canada, they have been documented to be an important source
7 of mortality on salmon (Beamish and Neville 1995). In the fall, adults migrate
8 back upstream into spawning areas and cease to feed. Fidelity to the streams in
9 which they were spawned remains unknown.

10 The species is expected to use Delta habitats primarily as a migration corridor
11 (DWR et al. 2013), and have been collected in Suisun Bay, Montezuma Slough,
12 and Delta sloughs during California Department of Fish and Wildlife (DFW)
13 plankton sampling efforts. CVP and SWP salvage data indicate that they are
14 found in the salvage primarily from December through March (DWR et al. 2013).
15 Juveniles are weak swimmers, frequently becoming entrained in water diversions
16 or turbine intakes of hydroelectric projects or becoming impinged on screens
17 meant to bypass juvenile salmonids or other fish (USFWS 2007).

18 Very little is known regarding the distribution, habitat use, and life history of this
19 species in the action area. Numerous adults (less than 200 millimeters [mm]),
20 presumably of spawning age, have been captured in rotary screw traps at RBDD
21 from March through June (Hanni et al. 2006). Individuals smaller than most
22 adults (greater than 200 mm), likely outmigrating macrophthalmia, have been
23 captured at RBDD and Feather River rotary screw traps from late September
24 through early June (Hanni et al. 2006). Factors limiting River Lamprey
25 populations in the Sacramento River are likely similar to those limiting salmonids
26 (Moyle et al. 2009). Quantitative data on populations are extremely limited, but
27 loss and degradation of historical habitats suggest populations have likely
28 declined (Moyle et al. 2009).

29 **9B.1.4 References**

- 30 Beamish, R. J. 1980. Adult biology of the River Lamprey (*Lampetra ayresi*) and
31 the Pacific lamprey (*Lamptera tridentata*) from the Pacific Coast of
32 Canada. *Canadian Journal of Fisheries and Aquatic Science* 37:1906-
33 1923.
- 34 Beamish, R. J., and J. H. Youson. 1987. Life history and abundance of young
35 adult *Lampetra ayresi* in the Fraser River and their possible impact on
36 salmon and herring stocks in the Strait of Georgia. *Canadian Journal of*
37 *Fisheries and Aquatic Science* 44:525-537.
- 38 Beamish, R. J., and C. M. Neville. 1995. Pacific salmon and Pacific herring
39 mortalities in the Fraser River plume caused by River Lamprey (*Lampetra*
40 *ayresi*). *Canadian Journal of Fisheries and Aquatic Sciences* 52: 644-650.
- 41 DWR (California Department of Water Resources), Bureau of Reclamation, U.S.
42 Fish and Wildlife Service, and National Marine Fisheries Service. 2013.
43 *Environmental impact report/ environmental impact statement for the Bay*

- 1 *Delta Conservation Plan. Draft.* Prepared by ICF International,
2 Sacramento, California. March.
- 3 Hanni, J., B. Poytress, and H. N. Blalock-Herod. 2006. *Spatial and temporal*
4 *distribution patterns of Pacific and River Lamprey in the Sacramento and*
5 *San Joaquin rivers and delta.* U.S. Fish and Wildlife Service.
- 6 Klamath-Siskiyou Wildlands Center, Siskiyou Regional Education Project,
7 Umpqua Watersheds, Friends of the Eel, North Coast Environmental
8 Center, Environmental Protection Information Center, Native Fish
9 Society, Center for Biological Diversity, Oregon Natural Resources
10 Council, Washington Trout, and Umpqua Valley Audubon Society. 2003.
11 *A petition for rules to list: Pacific lamprey (Lampetra tridentate), River*
12 *Lamprey (Lampetra ayresi), western brook lamprey (Lampetra*
13 *richardsoni); and Kern brook lamprey (Lampetra hubbsi) as threatened or*
14 *endangered under the Endangered Species Act.*
- 15 Moyle, P. B. 2002. *Inland fishes of California. Second edition.* University of
16 California Press, Berkeley.
- 17 Moyle, P. B., L. R. Brown, S. D. Chase, and R. M. Quinones. 2009. Status and
18 conservation of lampreys in California. *American Fisheries Society*
19 *Symposium 72: 279-292.*
- 20 Moyle, P. B., R.N, Yoshiyama, J. E. Williams, and E. D. Wikramanayake. 1995.
21 *Fish species of special concern of California.* Department of Wildlife and
22 Fisheries Biology, University of California, Davis.
- 23 Roos, J. F., P. Gilhousen, S. R. Killick, and E. R. Zyblut. 1973. Parasitism on
24 juvenile Pacific salmon (*Oncorhynchus*) and Pacific herring (*Clupea*
25 *harengus pallasii*) in the Straight of Georgia by the River Lamprey
26 (*Lampetra ayresi*). *Journal of the Fisheries Research Board of Canada*
27 30:565-568.
- 28 Scott, W .B., and E. J. Crossman. 1973. *Freshwater fishes of Canada.* Fisheries
29 Research Board of Canada Bulletin No. 184.
- 30 USFWS (U.S. Fish and Wildlife Service). 2007. *Fact sheet. Pacific lamprey -*
31 *Lampetra tridentata.* Portland, Oregon.
- 32 Vladykov, V. D., and W. I. Follett. 1958. Redescription of *Lampetra ayersi*
33 (Gunther) of western North America, a species of lamprey
34 (Petromyzontidae) distinct from *Lampetra fluviatilis* (Linnaeus) of
35 Europe. *Journal of the Fisheries Research Board of Canada* 15: 47-77.
- 36 Wang, J. C. S. 1986. *Fishes of the Sacramento-San Joaquin estuary and*
37 *adjacent waters, California: a guide to the early life histories.* Technical
38 Report 9. Prepared for the Interagency Ecological Study Program for the
39 Sacramento-San Joaquin Estuary by California Department of Water
40 Resources, California Department of Fish and Game, U.S. Bureau of
41 Reclamation and U.S. Fish and Wildlife Service.

1 Wydoski, R., and R. Whitney. 1979. *Inland fishes of Washington*. University of
2 Seattle Press, Seattle.

3 **9B.2 Pacific Lamprey (*Entosphenus tridentatus*)**

4 **9B.2.1 Legal Status**

5 Federal: None

6 State: None

7 The Pacific Lamprey was petitioned for listing by 12 conservation groups in
8 2003, along with three other lamprey species (Klamath-Siskiyou Wildlands
9 Center et al. 2003). The petition was declined by USFWS in 2004 because of
10 insufficient evidence that listing was warranted (USFWS 2004).

11 **9B.2.2 Distribution**

12 The Pacific Lamprey is a widely distributed anadromous species found in river
13 systems along the northern margin of the Pacific Ocean from central Baja
14 California north along the west coast of North America to the Bering Sea in
15 Alaska (Ruiz-Campos and Gonzales-Guzman 1996, Lin et al. 2008). Historically,
16 Pacific Lamprey were generally distributed wherever salmon and steelhead
17 occurred and sometimes upstream of waterfalls that are impassable to anadromous
18 salmonids. In California, they were historically found along the entire coast and
19 far inland (Moyle et al. 2009). However, recent data and anecdotal accounts
20 indicate that distribution of the Pacific Lamprey has been reduced in many river
21 systems, including the Sacramento-San Joaquin (Moyle et al. 2009). Although
22 widely distributed in the Sacramento-San Joaquin basin, the species is absent
23 from as much as 80 percent of its historical spawning habitats, primarily due to
24 migratory barriers (Moyle et al. 2009).

25 **9B.2.3 Life History and Habitat Requirements**

26 **9B.2.3.1 Adult Migration**

27 Pacific Lamprey are anadromous, rearing in freshwater before outmigrating to the
28 ocean, where they grow to full size prior to returning to their natal streams to
29 spawn. Pacific Lamprey are thought to remain in the ocean for approximately
30 18 to 40 months before returning to freshwater as sexually immature adults,
31 typically from late winter until early summer (Kan 1975, Beamish 1980). After
32 entering freshwater from the ocean, adult Pacific Lamprey typically spend
33 approximately 1 year in freshwater prior to spawning (Robinson and Bayer 2005,
34 Clemens et al. 2009, Stillwater Sciences 2010, Lampman 2011). The adult
35 freshwater residence period can be divided into three distinct stages: (1) Initial
36 migration from the ocean to holding areas, (2) pre-spawning holding, and
37 (3) secondary migration to spawn (Robinson and Bayer 2005; Clemens et al.
38 2010, 2012).

1 The initial migration from the ocean to upstream holding areas occurs from
2 approximately January until early August (Stillwater Sciences 2010, McCovey
3 2011, Clemens et al. 2012). In the Eel River and the nearby Klamath River,
4 where ample information exists, entry into freshwater from the ocean generally
5 begins in January and ends by June (Petersen-Lewis 2009, McCovey 2010,
6 Stillwater Sciences 2010). Most individuals cease upstream migration by
7 mid-July, although some individuals continue moving into August (McCovey
8 2010). Data from mid-water trawls in Suisun Bay and the lower Sacramento and
9 San Joaquin rivers indicate that adults likely migrate into the Sacramento-
10 San Joaquin Basin from late winter through early summer (Hanni and
11 Blalock-Herod 2006).

12 The pre-spawning holding stage begins when individuals cease upstream
13 movement in the summer, and continues until fish began their secondary
14 migration to spawn, generally in late winter or early spring (Robinson and Bayer
15 2005, McCovey 2010). During this holding period, most fish remain stationary
16 throughout the summer and fall, but some individuals undergo additional
17 upstream movements in the winter following high flow events (Robinson and
18 Bayer 2005, McCovey 2010). In the Sacramento River, adults, likely either in the
19 holding or spawning stage, have been detected at Glenn-Colusa Irrigation District
20 (GCID) from December through July and nearly year-round at RBDD (Hanni and
21 Blalock-Herod 2006). It is expected that adult Pacific Lamprey with varying
22 levels of sexual maturity are present in the Sacramento-San Joaquin Basin
23 throughout the year.

24 After the pre-spawning holding period, individuals undergo a secondary migration
25 from holding areas to spawning areas. This migration generally begins in late
26 winter and continues through July, by which time most individuals have spawned
27 and died (Robinson and Bayer 2005, Stillwater Sciences 2010, Lampman 2011).
28 During this secondary migration, movement to spawning areas can be both
29 upstream and downstream (Robinson and Bayer 2005, Lampman 2011).

30 Unlike Pacific salmon and steelhead (and like the Great Lakes Sea Lamprey;
31 Bergstedt and Seelye 1995), Pacific Lamprey do not necessarily home to natal
32 spawning streams (Moyle et al. 2009). Instead, migratory lampreys may select
33 spawning locations based on the presence of a pheromone-like substance secreted
34 by ammocoetes (Bjerselius et al. 2000, Vrieze and Sorensen 2001, Yun et al.
35 2011). Results of recent genetics research supports lack of homing by the Pacific
36 Lamprey. A study of Pacific Lamprey population structure found few genetic
37 differences among individuals sampled at widely dispersed sites across their
38 range, indicating substantial genetic exchange among populations from different
39 streams (Goodman et al. 2006).

40 **9B.2.3.2 Spawning**

41 Spawning typically takes place from March through July depending on water
42 temperature and local conditions such as seasonal flow regimes (Kan 1975,
43 Brumo et al. 2009, Gunckel et al. 2009). Evidence from the Santa Clara River in
44 southern California suggests that individuals in the southern portion of the

1 species' range can spawn as early as January, with peak spawning from February
2 to April (Chase 2001), whereas inland and northern populations initiate spawning
3 considerably later in the spring (Kan 1975, Beamish 1980, Brumo et al. 2009).
4 Hannon and Deason (2007) have documented Pacific Lamprey spawning in the
5 American River between early January and late May, with peak spawning
6 typically occurring in early April. Spawning occurs in both the mainstem of
7 medium-sized rivers and smaller tributaries (Luzier et al. 2006, Brumo et al. 2009,
8 Gunckel et al. 2009), and generally takes place in pool and run tailouts and low
9 gradient riffles. Both males and females build redds that are approximately
10 40-by-40 cm in area and are constructed in gravel and cobble substrate (Brumo
11 2006, Gunckel et al. 2009). Spawning substrate size typically ranges from
12 approximately 25 to 90 mm (1.0 to 3.5 inches), with a median of 48 mm
13 (1.9 inches) (Gunckel et al. 2009). Water velocity above redds ranges from 0.2 to
14 1.0 meters per second (m/s) (median 0.6 m/s), and depth varies from
15 approximately 0.2 to 1.1 m (0.7 to 3.6 feet [ft]) (Gunckel et al. 2009). Depending
16 on their size, females lay between 30,000 and 240,000 eggs (Kan 1975), which
17 are approximately 1.4 mm (0.06 inch) in diameter (Meeuwig et al. 2004). In
18 comparison, Chinook Salmon generally lay approximately 4,000 to 12,000 eggs
19 (Jasper and Evensen 2006). During spawning, eggs are released in clutches of
20 about 500 every 2 to 5 minutes (Pletcher 1963). Upon fertilization, eggs adhere to
21 sandy substrate in the gravel redd (Pletcher 1963).

22 Depending on water temperature, hatching occurs in approximately 2 to 3 weeks,
23 and yolk-sac larvae known as prolarvae remain in redd gravels for approximately
24 2 to 3 more weeks before emerging at night as 8-to-9-mm larvae, and drift
25 downstream to rear in depositional areas (Meeuwig et al. 2005, Brumo 2006).
26 Pacific Lamprey typically die soon after spawning (Kan 1975; Brumo 2006),
27 although there is some anecdotal evidence that this is not always the case (Moyle
28 2002; Michael 1980; Michael 1984).

29 **9B.2.3.3 Juvenile Rearing and Outmigration**

30 After larvae emerge from redds drifting downstream, the eyeless, toothless larvae
31 known as ammocoetes settle out of the water column and burrow into fine silt and
32 sand substrate in low-velocity, depositional areas such as pools, alcoves, and side
33 channels (Moore and Mallatt 1980, Torgensen and Close 2004, Stone and Barndt
34 2005). Ammocoete presence has also been shown to be associated with presence
35 of woody debris (Roni 2003, Graham and Brun 2006). Rearing Pacific Lamprey
36 ammocoetes appear to prefer rearing temperatures below 68°F (20 degrees
37 Celsius [°C]) (BioAnalysts, Inc. 2000); and temperatures above 82.4°F (28°C)
38 result in mortality of ammocoetes (van de Wetering and Ewing 1999). Depending
39 on factors influencing their growth rates, they remain in this habitat from 4 to
40 10 years, filter-feeding on algae and detrital matter prior to metamorphosing into
41 an adult form (Pletcher 1963, Moore and Mallatt 1980, Beamish and Levings
42 1991, van de Wetering 1998). During the ammocoete stage, individuals may
43 periodically move and relocate in response to changing water levels, channel
44 adjustments, or substrate movements (ULEP 1998). These factors generally result
45 in a gradual downstream movement that may lead to higher densities in

1 downstream reaches (Richards 1980). During metamorphosis, individuals
2 develop eyes, a suctoral disc, sharp teeth, and more-defined fins (McGree et al.
3 2008). After metamorphosis, smolt-like individuals known as macrophthalmia
4 migrate to the ocean—typically in conjunction with high-flow events between fall
5 and spring (van de Wetering 1998). Data from rotary screw trapping at sites in
6 the Sacramento-San Joaquin Basin indicate that emigration of Pacific Lamprey
7 macrophthalmia peaks from early winter through early summer; however, some
8 outmigration has been observed year-round in the mainstem Sacramento River at
9 both RBDD and GCID (Hanni and Blalock-Herod 2006). When abundant,
10 outmigrating Pacific Lamprey may act to buffer predation on juvenile and smolt
11 salmon because they are easier to capture than salmonids (Close et al. 2002).

12 **9B.2.3.4 Ocean Residence**

13 In the ocean, adult Pacific Lamprey feed parasitically on a variety of marine and
14 anadromous fishes such as salmon, flatfish, rockfish, and pollock. Pacific
15 Lamprey are preyed upon by sharks, sea lions, and other marine animals
16 (Richards and Beamish 1981, Beamish and Levings 1991, Close et al. 2002), and
17 have been captured in depths from 300 to 2,600 ft and as far as 62 miles off the
18 coast (USFWS 2007).

19 **9B.2.4 Population Trends**

20 In recent years, state, federal, and tribal agencies have expressed concern at the
21 apparent decline of lamprey populations in the Northwestern United States (Close
22 et al. 2002; Moser and Close 2003; CRBLTW 2005). Widespread anecdotal
23 accounts of decreased Pacific Lamprey spawning and carcasses have been
24 supported by a substantial reduction in counts of migrating individuals at dams
25 since the late 1960s (Moser and Close 2003, Klamath-Siskiyou Wildlands Center
26 et al. 2003). Very few data on Pacific Lamprey populations are available to
27 assess status in the Sacramento-San Joaquin Basin; however, loss of access to
28 historical habitat throughout California indicates that populations are greatly
29 suppressed compared with historical levels (Moyle et al. 2009).

30 Factors limiting Pacific Lamprey populations are numerous and interrelated
31 (Moser and Close 2003, Moyle et al. 2009). Although very little data or
32 published studies are available for Pacific Lamprey in the region, parallels in their
33 life cycle with salmon and steelhead suggest that these species are adversely
34 affected by many of the same factors. Lack of access to historical spawning
35 habitats because of dams, entrainment by water diversions, agricultural practices,
36 urban development, harvesting, mining, transportation, estuary modification, prey
37 abundance, and nonnative invasive species have all been cited as important
38 anthropogenic factors limiting the viability of Pacific Lamprey populations in
39 California (Moyle et al. 2009). In the Delta, the impacts of agricultural practices,
40 development, estuary modification, and predation by nonnative species are
41 expected to be particularly pronounced.

1 **9B.2.5 References**

- 2 Beamish, R. J. 1980. Adult biology of the River Lamprey (*Lampetra ayresi*) and
3 the Pacific Lamprey (*Lampetra tridentata*) from the Pacific coast of
4 Canada. *Canadian Journal of Fisheries and Aquatic Science* 37: 1906–
5 1923.
- 6 Beamish, R. J., and C. D. Levings. 1991. Abundance and freshwater migrations
7 of the anadromous parasitic lamprey, *Lampetra tridentata*, in a tributary of
8 the Fraser River, British Columbia. *Canadian Journal of Fisheries and*
9 *Aquatic Sciences* 48: 1250–1263.
- 10 Bergstedt, R. A., and J. G. Seelye. 1995. Evidence for lack of homing by sea
11 lampreys. *Transactions of the American Fisheries Society* 124: 235–239.
- 12 BioAnalysts, Inc. 2000. *A status of Pacific lamprey in the mid-Columbia region.*
13 *Rocky Reach Hydroelectric Project, FERC Project No. 2145.* Prepared
14 for Public Utility District No. 1 of Chelan County, Wenatchee,
15 Washington.
- 16 Bjerselius, R., W. Li, J. H. Teeter, J. G. Seelye, P. B. Johnsen, P. J. Maniak, G. C.
17 Grant, C. N. Polkinghorne, and P. W. Sorensen. 2000. Direct behavioral
18 evidence that unique bile acids released by larval sea lamprey
19 (*Petromyzon marinus*) function as a migratory pheromone. *Canadian*
20 *Journal of Fisheries and Aquatic Sciences* 57: 557–569.
- 21 Brumo, A. F. 2006. Spawning, larval recruitment, and early life survival of
22 Pacific lampreys in the South Fork Coquille River, Oregon. Master's
23 thesis. Oregon State University, Corvallis.
- 24 Brumo, A. F., L. Grandmontagne, S. N. Namitz, and D. F. Markle. 2009.
25 *Evaluation of approaches used to monitor Pacific lamprey spawning*
26 *populations in a coastal Oregon stream.* Biology, management, and
27 conservation of lampreys in North America. Edited by L. R. Brown, S. D.
28 Chase, M. G. Mesa, R. J. Beamish, and P. B. Moyle. Pp. 204–222.
29 American Fisheries Society, Symposium 72, Bethesda, Maryland.
- 30 Chase, S. D. 2001. Contributions to the life history of adult Pacific lamprey
31 (*Lampetra tridentate*) in the Santa Clara river of southern California.
32 *Bulletin of the Southern California Academy of Sciences* 100: 74–85.
- 33 Clemens, B. J., S. J. van de Wetering, J. Kaufman, R. A. Holt, and C. B. Schreck.
34 2009. Do summer temperatures trigger spring maturation in adult Pacific
35 lamprey, *Entosphenus tridentatus*? *Ecology of Freshwater Fish* 18: 418-
36 426.
- 37 Clemens, B. J., T. R. Binder, M. F. Docker, M. L. Moser, and S. A. Sower. 2010.
38 Similarities, differences, and unknowns in biology and management of
39 three parasitic lampreys of North America. *Fisheries* 35: 580-594.
- 40 Clemens, B. J., M. G. Mesa, R. J. Magie, D. A. Young, and C. B. Schreck. 2012.
41 Pre-spawning migration of adult Pacific lamprey, *Entosphenus tridentatus*,

- 1 in the Willamette River, Oregon, U.S.A. *Environmental Biology of Fishes*
2 93: 245–254.
- 3 Close, D. A., M. S. Fitzpatrick, and H. W. Li. 2002. The ecological and cultural
4 importance of a species at risk of extinction, Pacific lamprey. *Fisheries*
5 27:19–25
- 6 CRBLTW (Columbia River Basin Lamprey Technical Workgroup). 2005.
7 April 19. *Critical uncertainties for lamprey in the Columbia River Basin:*
8 *results from a strategic planning retreat of the Columbia River Lamprey*
9 *Technical Workgroup.*
10 <http://www.fws.gov/columbiariver/lampreywg/docs/CritUncertFinal.pdf>
- 11 Goodman, D., S. Reid, and M. Docker. 2006. *A phylogeographic analysis of the*
12 *Pacific lamprey Entosphenus tridentatus.* Revised final project report.
13 Prepared for U.S. Fish and Wildlife Service, Portland, Oregon.
- 14 Graham, J. C., and C. V. Brun. 2006. *Determining lamprey species composition,*
15 *larval distribution, and adult abundance in the Deschutes River, Oregon,*
16 *subbasin.* 2005 Annual Report. Bonneville Power Administration,
17 Portland, Oregon.
- 18 Gunckel, S. L., K. K. Jones, and S. E. Jacobs. 2009. *Spawning distribution and*
19 *habitat use of adult Pacific and western brook lampreys in Smith River,*
20 *Oregon.* Edited by L. R. Brown, S. D. Chase, M. G. Mesa, R. J. Beamish,
21 and P. B. Moyle. Pp. 173–189. *Biology, management, and conservation*
22 *of lampreys in North America.* American Fisheries Society, Symposium
23 72, Bethesda, Maryland.
- 24 Hanni, J., and H. N. Blalock-Herod. 2006. *Spatial and temporal distribution*
25 *patterns of Pacific and River Lamprey in the Sacramento and San Joaquin*
26 *rivers and delta.* U.S. Fish and Wildlife Service, Stockton and
27 Sacramento, California.
- 28 Hannon, J., and B. Deason. 2008. *American River steelhead (Oncorhynchus*
29 *mykiss) spawning, 2001-2007.* U.S. Bureau of Reclamation, Sacramento,
30 California.
- 31 Jasper J. R., and D. F. Evensen. 2006. *Length-girth, length-weight, and fecundity*
32 *of Yukon River Chinook salmon, Oncorhynchus tshawytscha.* *Fishery Data*
33 *Series No. 06-70.* Alaska Department of Fish and Game, Division of
34 Commercial Fisheries, Anchorage.
- 35 Kan, T. T. 1975. *Systematics, variation, distribution, and biology of lampreys of*
36 *the genus Lampetra in Oregon.* Doctoral dissertation. Oregon State
37 University, Corvallis.

- 1 Klamath-Siskiyou Wildlands Center, Siskiyou Regional Education Project,
2 Umpqua Watersheds, Friends of the Eel, Northcoast Environmental
3 Center, Environmental Protection Information Center, Native Fish
4 Society, Center for Biological Diversity, Oregon Natural Resources
5 Council, Washington Trout, and Umpqua Valley Audubon Society. 2003.
6 *A petition for rules to list: Pacific lamprey (Lampetra tridentata); River*
7 *Lamprey (Lampetra ayresi); western brook lamprey (Lampetra*
8 *richardsoni); and Kern brook lamprey (Lampetra hubbsi) as threatened or*
9 *endangered under the Endangered Species Act.* Submitted to the U.S.
10 Fish and Wildlife Service.
- 11 Lampman, R. T. 2011. *Passage, migration, behavior, and autoecology of adult*
12 *Pacific lamprey at Winchester Dam and within the North Umpqua River*
13 *Basin, OR.* Master's thesis, Oregon State University, Department of
14 Fisheries and Wildlife, Corvallis.
- 15 Lin, B., Z. Zhang, Y. Wang, K. P. Currens, A. Spidle, Y. Yamazaki, and D. A.
16 Close. 2008. Amplified fragment length polymorphism assessment of
17 genetic diversity in Pacific lampreys. *North American Journal of*
18 *Fisheries Management* 28: 1182-1193.
- 19 Luzier, C. W., G. Silver, and T. A. Whitesel. 2006. *Evaluate habitat use and*
20 *population dynamics of lampreys in Cedar Creek.* 2005 Annual Report.
21 Bonneville Power Administration, Portland, Oregon.
- 22 McCovey, B. W., Jr. 2011. *A small scale radio bio-telemetry study to monitor*
23 *migrating Pacific lamprey (Lampetra tridentata) within the Klamath River*
24 *basin.* Final progress report. Yurok Tribal Fisheries Program, Klamath
25 River Division, Hoopa, California.
- 26 McGree M., T. A. Whitesel, and J. Stone. 2008. Larval metamorphosis of
27 individual Pacific lampreys reared in captivity. *Transactions of the*
28 *American Fisheries Society* 137: 1866–1878.
- 29 Meeuwig, M., J. M. Bayer, and R. Reiche. 2004. *Identification of larval Pacific*
30 *lampreys (Lampetra tridentata), River Lampreys (L. ayresi), and western*
31 *brook lampreys (L. richardsoni) and thermal requirements of early life*
32 *history stages of lampreys.* 2000 Annual Report. Bonneville Power
33 Administration, Portland, Oregon.
- 34 Meeuwig, M. H., J. M. Bayer, and J. G. Seelye. 2005. Effects of temperature on
35 survival and development of early life stage Pacific and western brook
36 lampreys. *Transactions of the American Fisheries Society* 134:19–27.
- 37 Michael, J. H. 1980. Repeat spawning of Pacific lamprey. *California Fish and*
38 *Game Notes* 66:186–187.
- 39 Michael, J. H. 1984. Additional notes on the repeat spawning by Pacific
40 lamprey. *California Fish and Game Notes* 70:186–188.
- 41 Moore, J. W., and J. M. Mallatt. 1980. Feeding of larval lamprey. *Canadian*
42 *Journal of Fisheries and Aquatic Sciences* 37: 1658–1664.

Appendix 9B: Aquatic Species Life History Accounts

- 1 Moser, M. L., and D. A. Close. 2003. Assessing Pacific lamprey status in the
2 Columbia River Basin. *Northwest Science* 77: 116–125.
- 3 Moyle, P. B. 2002. *Inland fishes of California*. Revised edition. University of
4 California Press, Berkeley.
- 5 Moyle, P. B., L. R. Brown, S. D. Chase, and R. M. Quinones. 2009. *Status and*
6 *conservation of lampreys in California*. Edited by L. R. Brown, S. D.
7 Chase, M. G. Mesa, R. J. Beamish, and P. B. Moyle. Pp. 279–292.
8 Biology, management, and conservation of lampreys in North America.
9 American Fisheries Society, Symposium 72, Bethesda, Maryland.
- 10 Petersen-Lewis, R. S. 2009. *Yurok and Karuk traditional ecological knowledge:*
11 *insights into Pacific lamprey populations of the lower Klamath Basin*.
12 Edited by L. R. Brown, S. D. Chase, M. G. Mesa, R. J. Beamish, and P. B.
13 Moyle. Pp. 1-40. Biology, management, and conservation of lampreys in
14 North America. American Fisheries Society, Symposium 72, Bethesda,
15 Maryland.
- 16 Pletcher, F. T. 1963. *The life history and distribution of lampreys in the Salmon*
17 *and certain other rivers in British Columbia, Canada*. Master's thesis.
18 University of British Columbia, Vancouver.
- 19 Richards, J. E. 1980. *Freshwater biology of the anadromous Pacific lamprey*
20 *Lampetra tridentata*. Master's thesis. University of Guelph, Guelph,
21 Ontario. As cited in Oregon Department of Fish and Wildlife *Oregon*
22 *Lampreys: Natural History Status and Analysis of Management Issues*,
23 February 25, 2002.
- 24 Richards, J. E., and F. W. H. Beamish. 1981. Initiation of feeding and salinity
25 tolerance in the Pacific lamprey *Lampetra tridentata*. *Marine Biology* 63:
26 73–77.
- 27 Robinson, T. C., and J. M. Bayer. 2005. Upstream migration of Pacific lampreys
28 in the John Day River, Oregon: behavior, timing, and habitat use.
29 *Northwest Science* 79: 106-119.
- 30 Roni, P. 2003. Responses of benthic fishes and giant salamanders to placement
31 of large woody debris in small Pacific Northwest streams. *North*
32 *American Journal of Fisheries Management* 23: 1087–1097.
- 33 Ruiz-Campos, G., and S. Gonzalez-Guzman. 1996. First freshwater record of
34 Pacific lamprey, *Lampetra tridentata*, from Baja California, Mexico.
35 *California Fish and Game* 82: 144–146.
- 36 Stillwater Sciences. 2010. *Pacific lamprey in the Eel River basin: a summary of*
37 *current information and identification of research needs*. Prepared by
38 Stillwater Sciences, Arcata, California for Wiyot Tribe, Loleta, California.
- 39 Stone, J., and S. Barndt. 2005. Spatial distribution and habitat use of Pacific
40 lamprey (*Lampetra tridentata*) ammocoetes in a western Washington
41 stream. *Journal of Freshwater Ecology* 20: 171-185.

- 1 Torgensen C. E., and D. A. Close. 2004. Influence of habitat heterogeneity on
2 the distribution of larval Pacific lamprey (*Lampetra tridentata*) at two
3 spatial scales. *Freshwater Biology* 49: 614–630.
- 4 ULEP (Umpqua Land Exchange Project). 1998. *Mapping rules for Pacific*
5 *lamprey (Lampetra tridentata)*. ULEP, Roseburg, Oregon. As cited by
6 Friant Water Users Authority and Natural Resources Defense Council
7 *Draft Restoration Strategies for the San Joaquin River*, February 2003.
- 8 USFWS (U.S. Fish and Wildlife Service). 2004. Endangered and threatened
9 wildlife and plants; 90-day finding on a petition to list three species of
10 lampreys as threatened or endangered. *Federal Register* 69: 77158–77167.
- 11 _____. 2007. *Fact sheet: Pacific lamprey - Lampetra tridentata*. Portland,
12 Oregon.
13 [http://www.fws.gov/oregonfwo/Species/Data/PacificLamprey/Documents/
14 012808PL-FactSheet.pdf](http://www.fws.gov/oregonfwo/Species/Data/PacificLamprey/Documents/012808PL-FactSheet.pdf)
- 15 van de Wetering, S. J. 1998. *Aspects of life history characteristics and*
16 *physiological processes in smolting pacific lamprey (Lampetra tridentata)*
17 *in a central Oregon coast stream*. Master's thesis. Oregon State
18 University, Corvallis.
- 19 van de Wetering, S. J., and R. E. Ewing. 1999. *Lethal temperatures for larval*
20 *Pacific lamprey, Lampetra tridentata*. Confederated Tribes of the Siletz
21 Indians, Siletz, Oregon. As cited by Confederated Tribes of Warm Springs
22 Reservation of Oregon *Pacific Lamprey Passage Evaluation and*
23 *Mitigation Plan: Phase I*, March 2012.
- 24 Vrieze, L. A., and P. W. Sorensen. 2001. Laboratory assessment of the role of a
25 larval pheromone and natural stream odor in spawning stream localization
26 by migratory sea lamprey (*Petromyzon marinus*). *Canadian Journal of*
27 *Fisheries and Aquatic Sciences* 58: 2374–2385.
- 28 Yun, S.-S., A. J. Wildbill, M. J. Siefkes, M. L. Moser, A. H. Dittman, S. C.
29 Corbett, W. Li, and D. A. Close. 2011. Identification of putative
30 migratory pheromones from Pacific lamprey (*Lampetra tridentata*).
31 *Canadian Journal of Fisheries and Aquatic Sciences* 68: 2194–2203.

32 **9B.3 Green Sturgeon (*Acipenser medirostris*)**

33 **9B.3.1 Legal Status**

34 Federal: Threatened, Designated Critical Habitat

35 State: Species of Special Concern

36 The National Marine Fisheries Service (NMFS) has divided North American
37 Green Sturgeon into two Distinct Population Segments (DPSs) using the Eel
38 River in California as the line of demarcation (Adams et al. 2002). The Southern
39 DPS of North American Green Sturgeon includes all coastal and Central Valley
40 populations south of the Eel River, including the Sacramento River basin

1 (NMFS 2006). Although the Southern DPS is considered a separate population
2 from the Northern DPS based on genetic data and spawning locations, their
3 ranges outside the spawning season overlap (DFG 2002, Israel et al. 2004, Moser
4 and Lindley 2007).

5 After a status review was completed in 2002 (Adams et al. 2002), NMFS
6 determined that the Southern DPS did not warrant listing as threatened or
7 endangered but should be identified as a Species of Concern. This determination
8 was challenged in April 2003, and NMFS was asked to consider new information
9 on the species. NMFS updated its status review in February 2005 and determined
10 that the Southern DPS should be listed as threatened under the Federal
11 Endangered Species Act (ESA) (NMFS 2005a). NMFS published a final rule
12 (NMFS 2006) in April 2006 that listed the Southern DPS as threatened; the rule
13 took effect on June 6, 2006.

14 NMFS made a final critical habitat designation for the Southern DPS in October
15 2009 (74 *Federal Register* [FR] 52300). Designated critical habitat in California
16 includes the Sacramento, lower Feather, and lower Yuba rivers; the Delta; and
17 Suisun, San Pablo, and San Francisco bays (NMFS 2014). NMFS published a
18 final 4(d) rule to apply ESA take prohibitions to the Southern DPS in July 2010
19 (75 FR 30714). In California, Green Sturgeon is a Class 1 Species of Special
20 Concern (qualifying as threatened under the California Endangered Species Act).

21 **9B.3.2 Distribution**

22 North American Green Sturgeon are the most wide-ranging sturgeon species, with
23 ocean migrations ranging between northern Mexico and southern Alaska (Adams
24 et al. 2002). Ocean abundance and densities of Green Sturgeon increase north of
25 the Golden Gate because both the Southern DPS and Northern DPS generally
26 migrate northward along the coast when at sea (NMFS 2005b), as confirmed by
27 radio telemetry studies conducted on Sacramento River Green Sturgeon (DFG
28 2002). Subadult and adult Green Sturgeon migrate thousands of miles along the
29 western coast of the United States, often venturing into coastal estuaries like
30 Willapa Bay and Grays Harbor in Washington, where they concentrate during
31 summer (Adams et al. 2002). Two adults tagged in Willapa Bay have been
32 detected by radio telemetry stations in the Sacramento River (Heublein et al.
33 2009), indicating that Green Sturgeon from the Sacramento River migrate as far
34 north as Washington before returning to the Sacramento River to spawn.
35 Concentrations of Green Sturgeon have also been detected near Vancouver Island
36 in Canada (NMFS 2005b).

37 Though Green Sturgeon migrate thousands of miles through rivers, estuaries, and
38 ocean, they do not readily establish new spawning populations; they are known
39 from only three river systems: the Sacramento, Rogue, and Klamath. However,
40 data suggest there may be spawning populations in both the Eel River and the
41 Umpqua River in Oregon (NMFS 2005b), which could indicate previously
42 undetected relict populations or the seeds of new subpopulations. The population
43 that spawns in the Sacramento River constitutes the only known spawning
44 population in the Southern DPS. Populations may have formerly spawned in the

1 San Joaquin and South Fork Trinity rivers, but have since been extirpated (Israel
2 and Klimley 2008).

3 Green Sturgeon juveniles, subadults, and adults are widely distributed in the
4 Sacramento-San Joaquin Delta and estuary areas including San Pablo Bay
5 (Beamesderfer et al. 2004). The Sacramento-San Joaquin Delta serves as a
6 migratory corridor, feeding area, and juvenile rearing area for North American
7 Green Sturgeon in the Southern DPS.

8 **9B.3.2.1 Current Distribution in Sacramento River**

9 Within the Sacramento River, data only support an approximation of spawning
10 locations. Larval Green Sturgeon have been captured routinely, but in small
11 numbers in the RBDD rotary screw traps (River Mile [RM] 243.5) and the GCID
12 fish facility (RM 206), suggesting that spawning generally occurs upstream of
13 Hamilton City (RM 199), though spawning may occur as far downstream as
14 Chico Landing (RM 194) (Heublein et al. 2009). Adult Green Sturgeon have
15 been observed congregating below RBDD during late spring and early summer
16 when the gates are down (Beamesderfer et al. 2004), suggesting that these may be
17 ripe adults trying to migrate upstream to spawn. Spawning may occur in reaches
18 upstream of RBDD (DFG 2002), but the upstream extent of spawning is
19 unknown. In 1999, USFWS placed egg mats in the Sacramento River from
20 Anderson Cottonwood Irrigation District (ACID) Dam (RM 298.4) to 10 miles
21 downstream of RBDD to identify Green Sturgeon spawning sites; however, only
22 two eggs were captured, both at mats downstream of RBDD, so the study did not
23 clarify the location of specific spawning sites or the upstream extent of spawning
24 (Beamesderfer et al. 2004). A radio telemetry study detected two adult Green
25 Sturgeon migrating past a remote monitoring station above RBDD, suggesting
26 possible spawning migration upstream (Heublein et al. 2009).

27 **9B.3.2.2 Historical Distribution in Sacramento River**

28 The location and character of spawning sites in the Rogue and Klamath rivers
29 suggest that Green Sturgeon spawned in the Sacramento River above Keswick
30 Dam (RM 302), including in the Pit, McCloud, and Little Sacramento rivers
31 (Nakamoto et al. 1995, NMFS 2005b). The timing of upstream migration
32 (February through July) corresponds with winter base and high flows and spring
33 snowmelt. Adult Green Sturgeon likely entered the Sacramento River during
34 winter, holding in pools in the middle and upper Sacramento River until high-
35 flow events triggered upstream migration; high flows would have allowed adults
36 to navigate through areas that might otherwise act as passage barriers at lower
37 flows, providing them with access to steeper reaches with higher-velocity flows
38 and coarser substrates for broadcast spawning. Such areas may have resulted in
39 higher egg survival—crevices between substrate particles would provide the
40 Green Sturgeon's relatively non-adhesive eggs to settle in areas less accessible to
41 egg predators.

42 The location and characteristics of preferred Green Sturgeon spawning habitats in
43 the Rogue and Klamath rivers suggest that most of the historical spawning habitat
44 in the Sacramento River likely occurred upstream of Keswick Dam (RM 302),

1 with dam construction in the 1940s creating a permanent barrier that eliminated
2 access to the majority of spawning habitat. Upstream passage may have been
3 impeded even earlier by the seasonal operation of the ACID Dam, which began in
4 1916. Later-arriving adults would have even less access to spawning habitat
5 because of the operation of RBDD, which blocked upstream passage when the
6 gates were lowered in mid-May. Beginning in the late 1800s, those adults that
7 successfully spawned upstream might have had their larvae entrained by water
8 diversions such as the GCID diversion near Hamilton City.

9 **9B.3.3 Life History and Habitat Requirements**

10 Sturgeon live 40 to 50 years, delay maturation to large sizes (125 cm total length),
11 and spawn multiple times over their lifespan. This life history strategy has been
12 successful through normal environmental variation in the large river habitats
13 where spawning occurs. Their long lifespan, repeat spawning in multiple years,
14 and high fecundity allow them to persist through periodic droughts and
15 environmental catastrophes. The high fecundity associated with large size allows
16 them to produce large numbers of offspring when suitable spawning conditions
17 occur and compensate for years of poor reproductive and juvenile rearing
18 conditions. Adult Green Sturgeon do not spawn every year, and only a fraction of
19 the population enters fresh water where they might be at risk of a catastrophic
20 event (Beamesderfer et al. 2007). Though there are general descriptions of
21 preferred habitat conditions for Green Sturgeon, much of this information is
22 derived from Rogue River and Klamath River data, and little is known about
23 specific spawning, rearing, or holding locations in the Sacramento River.

24 **9B.3.3.1 Adult Migration**

25 Though Green Sturgeon spend most of their life in marine and estuarine
26 environments, they periodically migrate into freshwater streams to spawn,
27 spending up to 6 months in fresh water during their spawning migration.
28 Upstream migration generally begins in February and may last until late July
29 (Adams et al. 2002). In the Rogue River, telemetry studies have shown that adult
30 Green Sturgeon hold in low-velocity, deep-water habitats prior to migrating
31 upstream to spawn (Erickson et al. 2002). The adults move around in the pools
32 and may stray short distances, but the scope of their movement is limited. In the
33 Sacramento River, adult Green Sturgeon begin their upstream spawning
34 migrations into the San Francisco Bay in March and reach Knights Landing on
35 the Sacramento River during April (Heublein et al. 2006).

36 **9B.3.3.2 Spawning**

37 Spawning occurs between March and July, peaking between mid-April and mid-
38 June (Emmett et al. 1991). Based on the distribution of sturgeon eggs, larvae, and
39 juveniles in the Sacramento River, DFG (2002) indicated that Green Sturgeon
40 spawn in late spring and early summer above Hamilton City, possibly up to
41 Keswick Dam (Brown 2007). Israel and Klimley (2008) state that Green
42 Sturgeon spawn in the mainstem from the confluence of Battle Creek (river
43 kilometer 438) to the area upstream of Molinos, but may also spawn below
44 RBDD closer to GCID in some years. Adults spawn within about a week,

1 and females appear to spawn regardless of habitat conditions (Beamesderfer
2 et al. 2007).

3 Green Sturgeon prefer areas of fast, deep, turbulent water in mainstem channels
4 for spawning (Moyle 2002). They spawn in a variety of substrates, from clean
5 sand to bedrock, but prefer bed surfaces composed of coarse cobble (Moyle
6 2002). In the Rogue River, suspected spawning sites (inferred from the
7 movement of radio-tagged Green Sturgeon) have beds composed of cobbles and
8 boulders, with water depths greater than 10 to 15 feet (3 to 4.6 meters) and
9 turbulent water over slope breaks in the channel (Wildlife Conservation Society
10 2005). The interstitial spaces between large particles may provide eggs with
11 cover from predation (Moyle 2002). Eggs and larvae require cool water
12 temperatures and high dissolved oxygen concentrations while digesting their yolk
13 sac (Van Eenennaam et al. 2005).

14 Female Green Sturgeon produce 59,000 to 242,000 eggs, about 4.34 mm in
15 diameter (Van Eenennaam et al. 2001, 2006). Green Sturgeon eggs have the
16 largest mean diameter of any sturgeon species (Cech et al. 2000), but they lay
17 fewer eggs. The larger eggs may allow embryos to grow larger before hatching
18 and emerging from cover, increasing their survival relative to other sturgeon
19 species. Fecundity peaks at around age 24 years (Beamesderfer et al. 2007).

20 **9B.3.3.3 Juvenile Rearing**

21 Hatchling Green Sturgeon embryos seek nearby cover and remain under rocks
22 (Deng et al. 2002). After about 6 to 9 days, the hatchlings develop into larvae and
23 initiate exogenous foraging on the benthos (Deng et al. 2002, Kynard et al. 2005).
24 After a day or so, larvae disperse downstream for 1 to 2 weeks. Movements and
25 foraging activity during this period are nocturnal (Cech et al. 2000, Kynard et al.
26 2005). Larval Green Sturgeon are regularly captured during this dispersal stage at
27 about 2 weeks old (24- to 34-mm fork length) in rotary screw traps at RBDD
28 (DFG 2002, USFWS 2002) and 3 weeks old when captured farther downstream at
29 the GCID fish facility (Van Eenennaam et al. 2001). Following emergence in
30 early summer, larval Green Sturgeon migrating downstream with snowmelt flows
31 between May and July, growing quickly and becoming more tolerant of
32 increasing water temperatures and salinities. The upper thermal limit for optimal
33 development and hatching is between 17 to 18°C; temperatures higher than this
34 may affect development and hatching success, and complete mortality occurs at
35 temperatures above 23°C (Van Eenennaam et al. 2005).

36 Young Green Sturgeon appear to rear for the first 1 to 2 months in the Sacramento
37 River between Keswick Dam and Hamilton City (DFG 2002). Larvae and post-
38 larvae are present in the lower Sacramento River and North Delta between May
39 and October, primarily in June and July (DFG 2002). Little is known of
40 distribution and movements of young-of-the-year and riverine juveniles, but
41 observations suggest they may be distributed primarily in the mainstem
42 Sacramento River downstream of Anderson and in the brackish portions of the
43 north and interior Delta (Israel and Klimley 2008). Juvenile Green Sturgeon have
44 been captured in the Delta during all months of the year (Borthwick et al. 1999,

1 DFG 2002). Catches of 1- and 2-year-old Southern DPS Green Sturgeon on the
2 shoals in the lower San Joaquin River, at the CVP/SWP fish salvage facilities, and
3 in Suisun and San Pablo bays indicate that some fish rear in the estuary for at least
4 2 years (DFG 2002). Larger juvenile and subadult Green Sturgeon occur
5 throughout the estuary, possibly temporarily, after spending time in the ocean
6 (DFG 2002, Kelly et al. 2007).

7 The rearing habitat preferences of Green Sturgeon larvae and juveniles in the
8 Sacramento River are not well understood. Laboratory research has identified
9 water temperature thresholds for larval Green Sturgeon. Water temperatures
10 above 68°F (20°C) were found to be lethal to Green Sturgeon embryos by Cech
11 et al. (2000), and temperatures above 63 to 64°F (17 to 18°C) were found to be
12 stressful by Van Eenennaam et al. (2005). Cech et al. (2000) found that optimal
13 growth of larvae occurred at 59°F (15°C), with growth slowing at temperatures
14 below 52°F (11°C) and above 62°F (19°C).

15 Several studies suggest that juvenile Green Sturgeon rear in fresh water for 1 to
16 4 years, acclimating gradually to brackish environments before migrating to the
17 ocean (Beamesderfer and Webb 2002, Nakamoto et al. 1995). Larval Green
18 Sturgeon are captured at RBDD and the GCID fish facility between May and
19 August, with peak capture at RBDD in June and July and at the GCID fish facility
20 in July (Adams et al. 2002). Green Sturgeon larvae trapped at RBDD average
21 1.1 inches (2.9 cm) in length, while larvae trapped at the GCID fish facility
22 average 1.4 inches (3.6 cm) (Adams et al. 2002), suggesting that larvae move
23 downstream soon after hatching; however, it is not clear how long larval and
24 juvenile Green Sturgeon remain in the middle Sacramento River. Larval Green
25 Sturgeon grow quickly, reaching 2.9 inches (74 mm) by the time they become
26 juveniles at around 45 days posthatching (Deng 2000). Klamath River studies
27 indicate that juvenile Green Sturgeon can grow to 12 inches (30 cm) in their first
28 year and 24 inches (60 cm) within 2 to 3 years (Nakamoto et al. 1995). The small
29 size of salvaged juvenile Green Sturgeon at the CVP and SWP fish facilities
30 indicates that they move downstream to rear in the Bay-Delta estuary (Adams
31 et al. 2002), though it is unclear how long they remain before migrating to
32 the ocean.

33 While in the riverine environment, juveniles occupy low-light habitat and are
34 active at night (Kynard et al. 2005). Older juveniles may be adapted to move
35 through habitats with variable gradients of salinity, temperature, and dissolved
36 oxygen (Kelly et al. 2007, Moser and Lindley 2007). Their diet during their
37 Sacramento River residence is unknown, but likely consists of drifting and
38 benthic aquatic macroinvertebrates (Israel and Klimley 2008).

39 Stomach contents from adult and juvenile Green Sturgeon captured in the
40 Sacramento-San Joaquin Delta included shrimp, mollusks, amphipods, and small
41 fish (Radtke 1966, Houston 1988, Moyle et al. 1992). Stomachs of Green
42 Sturgeon caught in Suisun Bay contained *Corophium* sp. (amphipod), *Cragon*
43 *franciscorum* (bay shrimp), *Neomysis awatchensis* (Opossum shrimp:
44 synonymous with *Neomysis mercedis*), and annelid worms (Ganssle 1966).
45 Stomachs of Green Sturgeon caught in San Pablo Bay contained *C. franciscorum*,

1 *Macoma* sp. (clam), *Photis californica* (amphipod), *Corophium* sp., *Synidotea*
2 *laticauda* (isopod), and unidentified crab and fish (Ganssle 1966). Stomachs of
3 Green Sturgeon caught in the Delta contained *Corophium* sp. and *N. awatchensis*
4 (Radtke 1966). As a result of recent changes in the species composition of
5 macroinvertebrates inhabiting the Bay-Delta estuary due to nonnative species
6 introductions, the current diet of Green Sturgeon is likely to differ from that
7 reported in the 1960s.

8 In the Rogue River, adults hold in deep pools after spawning until late fall or early
9 winter, when they emigrate to downstream estuaries or the ocean, perhaps cued by
10 winter freshets that cause water temperatures to drop (Erickson et al. 2002).
11 Erickson et al. (2002) noted that adult downstream migration appeared correlated
12 with water temperatures below 50°F (10°C).

13 **9B.3.3.4 Ocean Residence**

14 Green Sturgeon from the Southern DPS pass through the San Francisco Bay to the
15 ocean where they commingle with other sturgeon populations (DFG 2002).
16 Subadult and adult sturgeon tagged in San Pablo Bay overwinter in bays and
17 estuaries along the coast of California, Oregon, and Washington, between
18 Monterey Bay and Willapa Bay, before moving farther north in the fall to
19 overwinter north of Vancouver Island. Individual Southern DPS Green Sturgeon
20 tagged by DFW in the San Francisco estuary have been recaptured off Santa Cruz,
21 California; in Winchester Bay on the southern Oregon coast; at the mouth of the
22 Columbia River; and in Grays Harbor, Washington (USFWS 1993, Moyle 2002).
23 Most Southern DPS Green Sturgeon tagged in the San Francisco estuary have
24 been returned from outside that estuary (Moyle 2002).

25 Subadult and adult Green Sturgeon generally migrate north along the coast once
26 they reach the ocean, concentrating in coastal estuaries like Willapa Bay, Grays
27 Harbor, and the Columbia River estuary during summer (Adams et al. 2002). The
28 strategy underlying summer visits to coastal estuaries is unclear because sampling
29 indicates they have relatively empty stomachs, suggesting they may not be
30 entering the estuaries to feed (Beamesderfer 2000). Females reach sexual
31 maturity after about 17 years and males after about 15 years (Adams et al. 2002).
32 Spawning was believed to occur every 3 to 5 years (Tracy 1990), but may occur
33 as frequently as every 2 years (NMFS 2005a).

34 **9B.3.4 Population Trends**

35 Empirical estimates of Green Sturgeon abundance are not available for any west
36 coast population including the Sacramento River population. Interpretations of
37 available time series of abundance index data for Green Sturgeon are confounded
38 by small sample sizes, intermittent reporting, fishery-dependent data, lack of
39 directed sampling, subsamples representing only a portion of the population, and
40 potential confusion with White Sturgeon (Adams et al. 2002). Musick et al.
41 (2000) noted that the North American Green Sturgeon population has declined by
42 88 percent throughout much of its range. The current population status of
43 Southern DPS Green Sturgeon is unknown (Beamesderfer et al. 2007, Adams
44 et al. 2007). Based on captures of Green Sturgeon during surveys for White

1 Sturgeon in San Francisco Bay (USFWS 1995), the population is believed to
2 range from several hundred to a few thousand adults.

3 Population estimates of Green Sturgeon in the Sacramento River have been
4 derived from data collected by monitoring programs that generally focus on other
5 species because few monitoring programs specifically address Green Sturgeon in
6 the Sacramento River. Green Sturgeon larvae are captured annually in the RBDD
7 rotary screw traps, the GCID fish screen, and the CVP/SWP fish salvage facilities
8 in the South Delta. DFW conducts annual trammel net surveys in San Pablo Bay
9 to track the White Sturgeon population, and Green Sturgeon often form part of the
10 incidental catch. Eggs, larvae, and post-larval Green Sturgeon are now commonly
11 reported in sampling directed at Green Sturgeon and other species (Beamesderfer
12 et al. 2004, Brown 2007). Young-of-the-year Green Sturgeon have been observed
13 annually since the late 1980s in fish sampling efforts at RBDD and the Glenn-
14 Colusa Canal (Beamesderfer et al. 2004). Green Sturgeon in the Sacramento
15 River are believed to have declined over the last 2 decades, with fewer than
16 50 spawning adults observed annually in the best spawning habitat along the
17 middle section of the Sacramento River (Israel and Klimley 2008).

18 Similar to other anadromous fish, Green Sturgeon in the Sacramento River likely
19 exhibit seasonal behavioral patterns in response to changes in flows, water
20 temperature, or other environmental cues affected by flows, but it is not clear if
21 anthropogenically induced changes in the flow regime have contributed to the
22 apparent decline in Green Sturgeon spawners. Researchers have hypothesized
23 that high spring flows, or the turbidity associated with them, may act as an
24 upstream migration cue. The annual catch of larval sturgeon at the RBDD and
25 GCID fish screens suggests that spawning occurs in the Sacramento River in most
26 years, regardless of water year type; however, it is unclear how many adults
27 return to spawn each year and whether there is a relationship between flows and
28 the number of adult spawners in any given year. The relationship between flow
29 and water temperature in the Sacramento River may influence Green Sturgeon
30 through controlling the amount of suitable rearing habitat available for larvae and
31 juveniles (Adams et al. 2002).

32 The most consistent sample data for Sacramento Green Sturgeon are for subadults
33 captured in San Pablo Bay during periodic White Sturgeon assessments since
34 1948. The California Department of Fish and Game (now DFW) measured and
35 identified 15,901 sturgeon of both species between 1954 and 1991 (USFWS
36 1996). Catches of subadult and adult North American Green Sturgeon by the
37 Interagency Ecological Program between 1996 and 2004 ranged from 1 to
38 212 Green Sturgeon per year, with the highest catch in 2001. Various attempts
39 have been made to infer Green Sturgeon abundance based on White Sturgeon
40 mark-recapture estimates and relative numbers of White and Green Sturgeon in
41 the catch (USFWS 1996, Moyle 2002). However, low catches of Green Sturgeon
42 preclude estimates or indices of Green Sturgeon abundance from these data
43 (Schaffter and Kohlhorst 1999, Gingras 2005). It is unclear if the high annual
44 variability in length distributions in these samples reflects variable recruitment
45 and abundance or is an artifact of small sample sizes, pooling of sample years, or

1 variable distribution patterns between freshwater and ocean portions of the
2 population.

3 Anecdotal information is also available on young-of-the-year Green Sturgeon
4 from juvenile fish monitoring efforts at RBDD and the GCID pumping facility on
5 the upper Sacramento River. Fish traps at these facilities captured between 0 and
6 2,068 juvenile Green Sturgeon per year (Adams et al. 2002), which suggests that
7 at least some Green Sturgeon reproduction occurred during the 1990s.

8 Approximately 3,000 juvenile Green Sturgeon have been observed in rotary screw
9 traps operated for juvenile salmon at RBDD from 1994 to 2000. Annual catches
10 have declined from 1995 through 2000 although the relationship of these catches
11 to actual abundance is unknown. Recent data indicate that little production
12 occurred in 2007 and 2008 (13 and 3 larvae, respectively, were captured in the
13 rotary screw traps at RBDD) (Poytress et al. 2009). Larger production occurred
14 in 2009, 2010, and 2011 (45, 122, and 643 larvae, respectively, were captured
15 using a benthic D-net), and no larvae were captured in 2012 (Poytress et al. 2010,
16 2011, 2012, 2013).

17 More than 2,000 juvenile Green Sturgeon have been collected in fyke and rotary
18 screw traps operated at the GCID diversion from 1986 to 2003. Operation of the
19 screw trap at the GCID site began in 1991 and has continued year-round with the
20 exception of 1998. Juvenile Green Sturgeon at the GCID site were consistently
21 larger in average size, but the number captured varied widely with no apparent
22 patterns in abundance between the two sites. Abundance of juveniles peaked
23 during June and July with a slightly earlier peak at RBDD (Adams et al. 2002).

24 Variable numbers of juvenile Green Sturgeon are observed each year from two
25 south Delta water diversion facilities (DFG 2002). When water is exported
26 through the CVP/SWP export facilities, fish become entrained into the diversion.
27 Since 1957, Reclamation has salvaged fish at the CVP Tracy Fish Collection
28 Facility. DFW's Fish Facilities Unit, in cooperation with DWR, began salvaging
29 fish at the SWP Skinner Delta Fish Protective Facility in 1968. The salvaged fish
30 are trucked daily and released at several sites in the western Delta. Salvage of
31 fish at both facilities is conducted 24 hours a day, 7 days a week, at regular
32 intervals. Salvaged fish are subsampled for species composition and numbers.
33 Numbers of Green Sturgeon observed at these fish facilities have declined since
34 the 1980s, which contributed to NMFS' decision to list the Southern DPS as a
35 threatened species. From the SWP Skinner Fish Facility, Green Sturgeon counts
36 averaged 87 individuals per year between 1981 and 2000 and 20 individuals per
37 year from 2001 through 2007. From the CVP Tracy Fish Collection Facility,
38 Green Sturgeon counts averaged 246 individuals per year between 1981 and 2000
39 and 53 individuals per year from 2001 through 2007 (Reclamation 2008).
40 Patterns were similar between total numbers per year and numbers adjusted for
41 water export volumes, which increased during the 1970s and 1980s. Annual
42 counts of Green Sturgeon from the SWP and CVP fish facilities are not
43 significantly correlated (Beamesderfer 2005).

1 USFWS (1996) reported substantial uncertainty in the interpretation of salvage
2 data for Green Sturgeon because of poor quality control on both counts and
3 species identification, expansions from small sample sizes, variability in sturgeon
4 dispersal patterns and collection vulnerability in response to complex changes in
5 Delta flow dynamics, and changes in configuration and operations over time.
6 Estimated sturgeon salvage numbers are expanded from subsamples, and actual
7 numbers of Green Sturgeon observed are substantially smaller. Historical
8 expansions were based on variable expansion rates (subsample duration) ranging
9 from 15 seconds per 2 hours when fish numbers were high to 100 percent
10 counting during periods when fish numbers were low. Under current conditions,
11 NMFS (2004) requires sampling of fish salvage at both the SWP and CVP
12 facilities at intervals of no less than 10 minutes every 2 hours. Green Sturgeon
13 salvage estimates reported for years before 1993 may be in error because of
14 uncertainty whether smaller sturgeon were correctly identified (USFWS 1996,
15 DFG 2002). Reclamation and DWR recommended that only more recent (from
16 1993 and later) CVP and SWP salvage data be used to analyze the effects of water
17 project operations on Green Sturgeon and other anadromous fishes.

18 **9B.3.5 References**

- 19 Adams, P. B., C. B. Grimes, J. E. Hightower, S. T. Lindley, and M. L. Moser.
20 2002. *Status review for North American green sturgeon, Acipenser*
21 *medirostris*. National Marine Fisheries Service, Santa Cruz, California.
- 22 Adams, P. B., C. B. Grimes, J. E. Hightower, S. T. Lindley, M. L. Moser, and M.
23 J. Parsley. 2007. Population Status of North American Green Sturgeon,
24 *Acipenser medirostris*. *Environmental Biology of Fishes* 79:339–356.
- 25 Beamesderfer, R. C. 2000. *Agenda and notes for green sturgeon workshop,*
26 *22-23 March 2000, Weitchpec, California*. Oregon Department of Fish
27 and Wildlife, Portland.
- 28 _____. 2005. *Technical Review of Recent Status Review and Proposed Listing*
29 *of Green Sturgeon*. Prepared for State Water Contractors. Available at:
30 http://www.fishsciences.net/reports/2005/tech_review_recent_status.pdf.
- 31 Beamesderfer, R. C. P., and M. A. H. Webb. 2002. *Green sturgeon status review*
32 *information*. S.P. Cramer and Associates, Gresham, Oregon.
- 33 Beamesderfer, R., M. Simpson, G. Kopp, J. Inman, A. Fuller, and D. Demko.
34 2004. *Historical and current information on green sturgeon occurrence*
35 *in the Sacramento and San Joaquin rivers and tributaries*. S.P. Cramer &
36 Associates, Oakdale, California. Prepared for State Water Contractors,
37 Sacramento, California.
- 38 Beamesderfer, R. C. P., M. L. Simpson, and G. J. Kopp. 2007. Use of life history
39 information in a population model for Sacramento green sturgeon.
40 *Environmental Biology of Fishes* 79: 315-337.

- 1 Borthwick, S. M., R. R. Corwin, and C. R. Liston. 1999. *Investigations of fish*
2 *entrainment by archimededs and internal helical pumps at the Red Bluff*
3 *Research Pumping Plant, Sacramento California: February 1997-June*
4 *1998*. Bureau of Reclamation, Red Bluff, California.
- 5 Brown, K. 2007. *Evidence of spawning by green sturgeon, Acipenser*
6 *medirostris, in the upper Sacramento River, California*.
- 7 Cech, J. J. Jr., S. I. Doroshov, G. P. Moberg, B. P. May, R. G. Schaffter, and D.
8 M. Kohlhorst. 2000. *Biological assessment of green sturgeon in the*
9 *Sacramento-San Joaquin watershed (Phase 1)*. Project No. 98-C-15,
10 Contract No. B-81738. Final report to CALFED Bay-Delta Program. As
11 cited by Adams et al. 2002.
- 12 Deng, X. 2000. *Artificial reproduction and early life stages of the green surgeon*
13 *(Acipenser medirostris)*. Doctoral dissertation. University of California,
14 Davis. As cited by Adams et al. 2002.
- 15 Deng X, J. P. Van Eenennaam, and S. I. Doroshov. 2002. *Comparison of early*
16 *life stages and growth of green and white sturgeon*. Biology,
17 management, and protection of North American sturgeon. Edited by W.
18 Van Winkle, P. J. Anders, D. H. Secor, and D. A. Dixon, 237-248.
19 Symposium 28. American Fisheries Society, Bethesda, Maryland.
- 20 DFG (California Department of Fish and Game). 2002. *California Department*
21 *of Fish and Game comments to NMFS regarding green sturgeon listing*.
22 Sacramento.
- 23 Emmett, R. L., S. L. Stone, S. A. Hinton, and M. E. Monaco. 1991. *Distribution*
24 *and abundance of fishes and invertebrates in west coast estuaries*.
25 Volume 2: Species life history summaries. ELMR Report No. 8.
26 NOS/NOAA Strategic Environmental Assessment Division, Rockville,
27 Maryland.
- 28 Erickson, D. L., J. A. North, J. E. Hightower, J. Weber, and L. Lauck. 2002.
29 Movement and habitat use of green sturgeon *Acipenser medirostris* in the
30 Rogue River, Oregon, USA. *Journal of Applied Ichthyology* 18: 565-569.
- 31 Ganssle, D. 1966. *Fishes and decapods of San Pablo and Suisun bays*.
32 Ecological studies of the Sacramento-San Joaquin Estuary, Part 1.
33 Compiled by D. W. Kelley, 1-40. California Department of Fish and
34 Game Bulletin 133.
- 35 Gingras, M. 2005. (San Pablo Bay white sturgeon abundance) X (green
36 sturgeon:white sturgeon catch ratio): is the product an index of green
37 sturgeon abundance? Symposium on green sturgeon and their
38 environment at Cal-Neva American Fisheries Society Annual Meeting.
39 Sacramento, California. As cited by Beamesderfer, R.C.P, G. Kopp, D.
40 Demko *Review of the Distribution, Life History and Population Dynamics*
41 *of Green Sturgeon with Reference to California's Central Valley*, 2005.

- 1 Heublein, J. C., J. T. Kelly, and A. P. Klimley. 2006. *Spawning migration and*
2 *habitat of green sturgeon, Acipenser medirostris, in the Sacramento River.*
3 Presentation at the CALFED Science Conference, Sacramento California.
4 As cited in DWR et al. 2013
- 5 Heublein, J. C., J. T. Kelly, C. E. Crocker, A. P. Klimley, and S. T. Lindley.
6 2009. Migration of green sturgeon *Acipenser medirostris* in the
7 Sacramento River. *Environmental Biology of Fishes* 84: 245-258.
- 8 Israel, J. A., J. F. Cordes, M. A. Blumberg, and B. May. 2004. Geographic
9 patterns of genetic differentiation among western U.S. collections of North
10 American green sturgeon (*Acipenser medirostris*). *North American*
11 *Journal of Fisheries Management* 24:922-931.
- 12 Israel, J. A., and A. P. Klimley. 2008. *Life history conceptual model for North*
13 *American green sturgeon (Acipenser medirostris).* Prepared for the Delta
14 Regional Ecosystem Restoration and Implementation Plan (DRERIP) by
15 University of California, Davis.
- 16 Kelly, J. T., A. P. Klimley, and C. E. Crocker. 2007. Movements of green
17 sturgeon, *Acipenser medrostris*, in the San Francisco Bay Estuary,
18 California. *Environmental Biology of Fishes* 79: 281-295.
- 19 Kynard, B., E. Parker, and T. Parker. 2005. Behavior of early life intervals of
20 Klamath River green sturgeon, *Acipenser medirostris*, with a note on body
21 color. *Environmental Biology of Fishes* 72:85-97.
- 22 Moser, M. L., and S. T. Lindley. 2007. Use of Washington estuaries by subadult
23 and adult green sturgeon. *Environmental Biology of Fishes* 79: 243-253.
- 24 Moyle, P. B. 2002. *Inland fishes of California.* Revised edition. University of
25 California Press, Berkeley.
- 26 Moyle, P. B., P. J. Foley, and R. M. Yoshiyama. 1992. *Status of green sturgeon,*
27 *Acipenser medirostris, in California.* Report by University of California
28 at Davis to the National Marine Fisheries Service, Terminal Island,
29 California.
- 30 Musick, J. A., M. M. Harbin, S. A. Berkeley, G. H. Burgess, A. M. Eklund, L.
31 Findley, R. G. Gilmore, J. T. Golden, D. S. Ha, G. R. Huntsman, J. C.
32 McGovern, S. J. Parker, S. G. Poss, E. Sala, T. W. Schmidt, G. R.
33 Sedberry, H. Weeks, and S. G. Wright. 2000. Marine, Estuarine, and
34 Diadromous fish stocks at Risk of Extinction in North America (exclusive
35 of Pacific Salmonids). *Fisheries* 25(11):6-30.
- 36 Nakamoto, R. J., T. T. Kisanuki, and G. H. Goldsmith. 1995. *Age and growth of*
37 *Klamath River green sturgeon (Acipenser medirostris).* Project 93-FP-13.
38 U.S. Fish and Wildlife Service, Coastal California Fish and Wildlife
39 Office, Arcata, California.
- 40 NMFS (National Marine Fisheries Service). 2004. *Endangered Species Act –*
41 *Section 7 consultation biological opinion on the long-term Central Valley*

- 1 *Project and state water project operations, criteria, and plan (OCAP BO).*
2 Southwest Region. Long Beach, California.
- 3 _____. 2005a. *Green Sturgeon (Acipenser medirostris) Status Review Update.*
4 NOAA Fisheries, Southwest Fisheries Science Center.
- 5 _____. 2005b. Endangered and threatened wildlife and plants: proposed
6 threatened status for Southern Distinct Population Segment of North
7 American green sturgeon. *Federal Register* 70: 17386-17401.
- 8 _____. 2006. Endangered and threatened wildlife and plants: threatened status
9 for Southern Distinct Population Segment of North American green
10 sturgeon: final rule. *Federal Register* 71: 17757-17766.
- 11 _____. 2014. Green Sturgeon. NOAA Fisheries Office of Protected Resources.
12 Available at:
13 <http://www.nmfs.noaa.gov/pr/species/fish/greensturgeon.htm>. Updated
14 June 2, 2014.
- 15 Piper, R. G., I. B. McElwain, L. E. Orme, J. P. McCraren, L. G. Fowler, and J. R.
16 Leonard. 1982. *Fish hatchery management*. U.S. Fish and Wildlife
17 Service.
- 18 Poytress, W. R., J. J. Gruber, D. A. Trachtenbarg, and J. P. Van Eenennaam.
19 2009. *2008 Upper Sacramento River Green Sturgeon Spawning Habitat*
20 *and Larval Migration Surveys*. March. Annual Report of U.S. Fish and
21 Wildlife Service to Bureau of Reclamation, Red Bluff Fish Passage
22 Program, Red Bluff, CA.
- 23 Poytress, W. R., J. J. Gruber, and J. Van Eenennaam. 2010. *2009 Upper*
24 *Sacramento River Green Sturgeon Spawning Habitat and Larval*
25 *Migration Surveys*. Final Annual Report. July. Annual Report of U.S.
26 Fish and Wildlife Service to Bureau of Reclamation, Red Bluff Fish
27 Passage Program, Red Bluff, CA.
- 28 _____. 2011. *2010 Upper Sacramento River Green Sturgeon Spawning Habitat*
29 *and Larval Migration Surveys*. Final Annual Report. February. Annual
30 Report of U.S. Fish and Wildlife Service to Bureau of Reclamation, Red
31 Bluff Fish Passage Program, Red Bluff, CA.
- 32 _____. 2012. *2011 Upper Sacramento River Green Sturgeon Spawning Habitat*
33 *and Larval Migration Surveys*. Final Annual Report. March. Annual
34 Report of U.S. Fish and Wildlife Service to Bureau of Reclamation, Red
35 Bluff Fish Passage Program, Red Bluff, CA.
- 36 Poytress, W. R., J. J. Gruber, C. E., Praetorius, and J. P. Van Eenennaam. 2013.
37 *2012 Upper Sacramento River Green Sturgeon Spawning Habitat and*
38 *Young of the Year Migration Surveys*. Annual Report of U.S. Fish and
39 Wildlife Service to Bureau of Reclamation, Red Bluff, CA.
- 40 Radtke, L. D. 1966. Distribution of smelt, juvenile sturgeon, and starry flounder
41 in the Sacramento-San Joaquin Delta with observations on food of
42 sturgeon. Ecological studies of the Sacramento-San Joaquin Estuary. Part

- 1 II. Edited by Turner, J. L. and D. W. Kelly. California Department of
2 Fish and Game. *Fish Bulletin* 136: 115-119.
- 3 Reclamation (Bureau of Reclamation). 2008. *Long-term Central Valley Project
4 and State Water Project Operations, Criteria, and Plan (OCAP).*
5 Biological assessment.
- 6 Schaffter, R. G, and D. W. Kohlhorst. 1999. Status of white sturgeon in the
7 Sacramento-San Joaquin Estuary. *California Fish and Game* 85: 37-41.
- 8 Tracy, C. 1990. *Green sturgeon meeting and comments.* Memorandum.
9 Washington Department of Fisheries. As cited by Adams et al. 2002.
- 10 USFWS (U.S. Fish and Wildlife Service). 1993. *Endangered and threatened
11 wildlife and plants: determination of threatened status for the delta smelt.*
12 *Federal Register* 58:2854–12863.
- 13 _____. 1995. *Working Paper: Habitat Restoration Actions to Double Natural
14 Production of Anadromous Fish in the Central Valley of California.*
15 Volume 2. May 9. Prepared under the direction of the Anadromous Fish
16 Restoration Program Core Group, Stockton, CA.
- 17 _____. 1996. *Recovery plan for the Sacramento-San Joaquin Delta native fishes.*
18 U.S. Fish and Wildlife Service, Portland Oregon.
- 19 _____. 2002. *Spawning areas of green sturgeon Acipenser medirostris in the
20 upper Sacramento River, California.* U.S. Fish and Wildlife Service, Red
21 Bluff, California.
- 22 Van Eenennaam, J. P., M. A. H. Webb, X. Deng, S. I. Doroshov, R. B. Mayfield,
23 J. J. Cech Jr., D. C. Hillemeier, and T. E. Willson. 2001. Artificial
24 spawning and larval rearing of Klamath River green sturgeon.
25 *Transactions of the American Fisheries Society* 130: 159-165.
- 26 Van Eenennaam, J. P., J. Linares-Casenave, X. Deng, and S. I. Doroshov. 2005.
27 Effect of incubation temperature on green sturgeon embryos, *Acipenser
28 medirostris.* *Environmental Biology of Fishes* 72: 145-154.
- 29 Van Eenennaam, J. P., J. Linares-Casenave, S. I. Doroshov, D. C. Hillemeier, T.
30 E. Wilson, and A. A. Nova. 2006. Reproductive conditions of Klamath
31 River green sturgeon. *Transactions of the American Fisheries Society*
32 135:151-163.
- 33 Wildlife Conservation Society. 2005. Research on green sturgeon spawning in
34 the Rogue River received by Michael Fainter, Stillwater Sciences,
35 Berkeley, California, on July 14, 2005, via phone conversation with Dan
36 Erickson, Wildlife Conservation Society.

1 **9B.4 White Sturgeon (*Acipenser transmontanus*)**

2 **9B.4.1 Legal Status**

3 Federal: None

4 State: None

5 **9B.4.2 Distribution**

6 White Sturgeon have a marine distribution spanning from the Gulf of Alaska
7 south to Mexico, but a spawning distribution ranging only from the Sacramento
8 River northward. Currently, self-sustaining spawning populations are only known
9 to occur in the Sacramento, Fraser, and Columbia rivers.

10 In California, the largest numbers are in the San Francisco Bay estuary, with
11 spawning occurring mainly in the Sacramento and Feather rivers. White Sturgeon
12 historically ranged into upper portions of the Sacramento system including the Pit
13 River, and a substantial number were trapped in and above Lake Shasta when
14 Shasta Dam was closed in 1944 and successfully reproduced until the early 1960s
15 (State Water Contractors 2004). They may have occurred historically in the
16 San Joaquin River based on habitat similarities with these other watersheds.

17 Adult sturgeon were caught in the sport fishery industry in the San Joaquin River
18 between Mossdale and the confluence with the Merced River in late winter and
19 early spring, suggesting this was a spawning run (Kohlhorst 1976). Kohlhorst
20 et al. (1991) estimated that approximately 10 percent of the Sacramento River
21 system spawning population migrated up the San Joaquin River. Spawning may
22 occur in the San Joaquin River when flows and water quality permit; however, no
23 evidence of spawning is present (Kohlhorst 1976, Kohlhorst et al. 1991).

24 Landlocked populations are located above major dams in the Columbia River
25 basin, and residual non-reproducing fish above the Shasta Dam and Friant Dam
26 have been occasionally found.

27 Adult White Sturgeon are occasionally noted in the San Joaquin River during
28 DFW fall midwater trawls, DFW summer townet surveys, and University of
29 California Davis Suisun Marsh fisheries monitoring. White Sturgeon spawning
30 has recently been confirmed in the lower San Joaquin River (Jackson and Van
31 Eenennaam 2013), and the U.S. Geological Survey (USGS) is currently mapping
32 and characterizing White Sturgeon spawning habitat in the lower portion of the river
33 (USGS 2015).

34 **9B.4.3 Life History and Habitat Requirements**

35 White Sturgeon are long-lived, late maturing, and have a high fecundity (Israel et
36 al. 2015) Because White Sturgeon require a long time to mature, large year
37 classes are typically associated with years of high outflow (Kohlhorst et al. 1991,
38 Schaffter and Kohlhorst 1999), and population size can fluctuate to extremes
39 (Schaffter and Kohlhorst 1999).

1 Reports of maximum size and age of White Sturgeon are as great as 6 meters fork
2 length (FL) (820 kilograms) and greater than 100 years, although they generally
3 do not exceed 2 meters FL or 27 years of age. Males mature in 10 to 12 years
4 (75 to 105 centimeters FL) and females in 12 to 16 years (95 to 135 centimeters
5 FL). Maturation depends largely on temperature and photoperiod.

6 **9B.4.3.1 Adult Migrations and Spawning**

7 White Sturgeon migrate upstream in late winter. Upstream migration is usually
8 initiated by a large pulse flow (Schaffter 1997), and not all adults will spawn each
9 year. Because of this, successful year classes tend to occur at irregular intervals,
10 and therefore numbers of adult fish within a population can fluctuate significantly.
11 Although males may spawn each year, females usually spawn once every 2 to
12 4 years. White Sturgeon have high fecundities, and typical females may have as
13 many as 200,000 eggs. Spawning occurs over deep gravel riffles or in deep pools
14 with swift currents and rock bottoms between late February and early June when
15 temperatures are between 8°C and 19°C. Eggs become adhesive subsequent to
16 fertilization, and adhere to the substrate until they hatch 4 to 12 days later,
17 depending on temperature. Once the eggs have been deposited, the adults move
18 back downstream to the estuary. Larvae hatch in 1 to 2 weeks, depending on
19 temperature. Once the yolk sac is absorbed (approximately 1 week after
20 hatching), the larvae can begin to actively forage along the benthos.

21 In the Sacramento River, most White Sturgeon spawn downstream of the Glenn-
22 Colusa Irrigation Dam.

23 **9B.4.3.2 Juvenile Rearing**

24 White Sturgeon are benthic feeders, and adults may move into food-rich areas to
25 forage. Juveniles consume mainly crustaceans, especially amphipods and
26 opossum shrimp. Adult diets include invertebrates (mainly clams, crabs, and
27 shrimp), as well as fish, especially herring, anchovy, Striped Bass, and smelt.
28 White Sturgeon are opportunistic predators and may feed on many introduced
29 species.

30 Juvenile sturgeon are often found in upper reaches of estuaries in comparison to
31 adults, which suggests that there is a correlation between size and salinity
32 tolerance.

33 **9B.4.3.3 Estuary and Ocean Residence**

34 White Sturgeon primarily live in brackish portions of estuaries where they tend to
35 concentrate in deep sections having soft substrate. They move according to
36 salinity changes, and may swim into intertidal zones to feed at high tide.

37 Recent stomach content analysis of White Sturgeon from the San Francisco Bay
38 estuary indicates that the invasive overbite clam, *Corbula amurensis*, may now be
39 a major component of the White Sturgeon diet (Zeug et al. 2014), and unopened
40 clams were often observed throughout the alimentary canal (Kogut 2008).
41 Kogut's study found that at least 91 percent of clams that passed through sturgeon
42 digestive tracts were alive. This suggests sturgeon are potential vehicles for

1 transport of adult overbite clams and also raise concern about the effect of this
2 invasive clam on sturgeon nutrition and contaminant exposure.

3 In the ocean, White Sturgeon have been known to migrate long distances, but
4 spend most of their life in brackish portions of large river estuaries.

5 **9B.4.4 Population Trends**

6 There is a relatively strong relationship between Delta outflow and year class
7 strength during the period when white sturgeon are spawning and young white
8 sturgeon are migrating downstream (March-July). There is a threshold at about
9 50,000 cfs such that year classes are generally strong when flows are above the
10 threshold (Gingras et al. 2014). NMFS (2005) also noted a relationships between
11 flow and apparent White Sturgeon spawning success. A sturgeon population
12 study conducted by the California Department of Fish and Wildlife has been
13 ongoing intermittently since 1967. In 2014, catch per 100 net-fathom hour of
14 white sturgeon within the current slot limit (102-152 cm FL) was 0.46 ± 0.05
15 (SE); in 2013, catch per 100 net-fathom hour of white sturgeon within the current
16 slot limit was 0.4 ± 0.1 (SE). Both of these values are well below the historical
17 average of 2.8 (DuBois et al. 2014). Large numbers of young white sturgeon
18 have only been produced twice in the last 15 years, in 1998 and 2006 (Gingras et
19 al. 2014). The 2010-2014 White Sturgeon length frequency distributions show:
20 (1) strong cohorts (from mid-to-late 1990s) within the legally-harvestable size
21 range have substantially diminished; and (2) the progression of a strong cohort
22 (from 2006) toward harvestable size (DuBois et al. 2014). Given the trends in
23 catch-per-unit-effort (CPUE) and harvest, the amount of harvest, and harvest
24 rates, it's quite clear that harvest is the main reason CPUE and abundance have
25 declined so steeply (Gingras et al. 2014).

26 Periodic high flows in the 1990s produced small increases in White Sturgeon
27 salvage catches, but salvage numbers were much lower than prior to 1985.
28 USFWS (1996) in the *Sacramento/San Joaquin Delta Native Fishes Recovery*
29 *Plan* also reported that juvenile sturgeon are probably more vulnerable to
30 entrainment at the SWP and CVP at low to intermediate flows during those years
31 when river and Delta inflow are normal or below normal.

32 **9B.4.5 References**

- 33 Brown, L. R., and P. B. Moyle. 1993. Distribution, ecology, and status of fishes of
34 the San Joaquin River drainage, California. *California Fish and Game*
35 *Bulletin* 79:96-113.
- 36 DuBois, J., M. Harris, and L. Warkentin. 2014. 2014 Field Season Summary for
37 the Sturgeon Population Study. California Department of Fish and
38 Wildlife, Bay Delta Region (Stockton). 18 November 2014.
- 39 Gingras, M., J. DuBois, and M. Fish. 2014. Impact of Water Operations and
40 Overfishing on White Sturgeon. Presentation at the IEP Annual
41 Workshop, Folsom, CA, 27 February 2014.

- 1 Israel, J., A. Drauch, and M. Gingras. 2015. Life History Conceptual Model for
2 White Sturgeon (*Acipenser transmontanus*). DRERIP Delta Conceptual
3 Model. Sacramento (CA): Delta Regional Ecosystem Restoration
4 Implementation Plan.
5 http://www.dfg.ca.gov/ERP/drerip_conceptual_models.asp (Accessed
6 October 17, 2015).
- 7 Jackson, Z. J., and J. P. Van Eenennaam. 2013. 2012 San Joaquin River Sturgeon
8 Spawning Survey. Stockton Fish and Wildlife Office, Anadromous Fish
9 Restoration Program, U.S. Fish and Wildlife Service, Lodi, California.
- 10 Kogut, N. 2008. Overbite clams, *Corbula amerensis*, defecated alive by White
11 Sturgeon, *Acipenser transmontanus*. *California Fish and Game* 94:143-
12 149.
- 13 Kohlhorst, D. W. 1976. Sturgeon spawning in the Sacramento River in 1973, as
14 determined by distribution of larvae. *California Fish and Game* 62:32-40.
- 15 Kohlhorst, D. W., L. W. Botsford, J. S. Brennan, and G. M. Cailliet. 1991.
16 Aspects of the structure and dynamics of an exploited central California
17 population of White Sturgeon (*Acipenser transmontanus*). In *Acipenser*,
18 pp. 277-293. Edited by P. Williot. CEMAGREF, Bordeaux, France.
- 19 Moyle, P. B. 2002. *Inland Fishes of California*. Revised edition. University of
20 California Press, Berkeley.
- 21 NMFS (National Marine Fisheries Service). 2005. Endangered and threatened
22 wildlife and plants: proposed threatened status for Southern Distinct
23 Population Segment of North American Green Sturgeon. *Federal Register*
24 70: 17386-17401.
- 25 Schaffter, R. G. 1997. White Sturgeon spawning migrations and location of
26 spawning habitat in the Sacramento River, California. *California Fish and*
27 *Game* 83: 1-20.
- 28 Schaffter, R. G., and D. W. Kohlhorst. 1999. Status of White Sturgeon in the
29 Sacramento-San Joaquin Estuary. *California Fish and Game* 85: 37-41.
- 30 State Water Contractors. 2004. *Historical and Current Information on Green*
31 *Sturgeon Occurrence in the Sacramento and San Joaquin Rivers and*
32 *Tributaries*. Prepared by R. Beamesderfer, M. Simpson, G. Kopp, J.
33 Inman, A. Fuller, and D. Demko, S.P. Cramer and Associates, Oakdale,
34 California, for State Water Contractors, Sacramento, California.
- 35 USFWS (U.S. Fish and Wildlife Service). 1996. *Sacramento-San Joaquin Delta*
36 *Native Fishes Recovery Plan*. Portland, Oregon.
- 37 USGS (U.S. Geological Survey). 2015. Mapping Sturgeon Spawning Habitat in
38 the Lower San Joaquin River. [http://ca.water.usgs.gov/projects/2011-](http://ca.water.usgs.gov/projects/2011-20.html)
39 [20.html](http://ca.water.usgs.gov/projects/2011-20.html). Website accessed on June 2, 2015.
- 40 Zeug, S.C., A. Brodsky, N. Kogut, A.R. Stewart, and J.E. Merz. 2014. Ancient
41 fish and recent invaders: white sturgeon *Acipenser transmontanus* diet

1 response to invasivespecies-mediated changes in a benthic prey
2 assemblage. Mar. Ecol. Prog. Ser. Vol. 514: 163-174, 2014. doi:
3 10.3354/meps11002

4 **9B.5 Chinook Salmon (*Oncorhynchus tshawytscha*)**

5 **9B.5.1 Introduction**

6 The Sacramento-San Joaquin Delta functions as a migration corridor and potential
7 rearing area for adult and juvenile Chinook Salmon in the Sacramento and
8 San Joaquin River basins. The Sacramento River basin supports four runs of
9 Chinook Salmon: winter-run, spring-run, fall-run, and late fall-run. The
10 San Joaquin River basin currently supports fall-run (and possibly late fall-run)
11 Chinook Salmon in its lower tributaries: the Merced, Tuolumne, and Stanislaus
12 rivers. The winter-run consists of a single population spawning in the Sacramento
13 River mainstem below Keswick Dam. The other runs consist of populations that
14 spawn in multiple tributaries. Three ESUs of Chinook Salmon are represented in
15 the combined basins: Sacramento River winter-run (federally listed as
16 endangered), Sacramento River spring-run (federally listed as threatened), and
17 Central Valley fall-run and late fall-run (species of concern). Each of these runs
18 exhibits a variety of different life-history strategies.

19 **9B.5.2 Chinook Salmon Habitat Requirements**

20 The Sacramento River basin is the largest watershed in California (about
21 27,000 mi²) and empties into the largest estuary on the west coast of the United
22 States. This diverse basin is unique in that it supports four runs of Chinook
23 Salmon, including the winter-run, which only occurs in the Sacramento River
24 basin. Because the four runs exhibit a variety of different life-history strategies,
25 anthropogenic activities in the basin have affected each of the runs differently.
26 The habitat requirements and the life-history strategies of the four runs are
27 discussed below.

28 **9B.5.2.1 Upstream Migration and Holding**

29 Adult Chinook Salmon require water deeper than 0.8 ft (24 cm) and water
30 velocities less than 8 ft/s (2.4 m/s) for successful upstream migration (Thompson
31 1972). Adult Chinook Salmon appear to be less capable of negotiating fish
32 ladders, culverts, and waterfalls during upstream migration than Coho Salmon or
33 steelhead (Nicholas and Hankin 1989), due in part to slower swimming speeds
34 and inferior jumping ability compared to steelhead (Reiser and Peacock 1985,
35 Bell 1986). The maximum jumping height for Chinook Salmon has been
36 calculated to be approximately 7.9 ft (2.4 m) (Bjornn and Reiser 1991).

37 Both winter-run and spring-run Chinook Salmon return to the Sacramento River
38 when reproductively immature, typically holding for a few months in deep pools
39 near spawning areas until spawning. Adult winter-run and spring-run Chinook
40 Salmon require large, deep pools with flowing water for summer holding, tending
41 to hold in pools with depths greater than 4.9 ft (greater than 1.5 m) that contain

1 cover from undercut banks, overhanging vegetation, boulders, or woody debris
2 (Lindsay et al. 1986), and have water velocities ranging from 0.5 to 1.2 ft/s (15 to
3 37 cm/s) (Marcotte 1984). Water temperatures for adult Chinook holding are
4 reportedly best when less than 60.8°F (less than 16°C), and lethal when greater
5 than 80.6°F (greater than 27°C) (Moyle et al. 1995). Spring-run Chinook Salmon
6 in the Sacramento River system typically hold in pools below 69.8 to 77°F (21 to
7 25°C).

8 In general, adult Chinook Salmon appear capable of migrating upstream under a
9 wide range of temperatures. Bell (1986) reported that salmon and steelhead
10 migrate upstream in water temperatures that range from 3 to 20°C (37 to 68°F).
11 Bell (1986) reports that temperatures ranging from 3 to 13°C (37 to 55°F) are
12 suitable for upstream migration of spring-run Chinook Salmon, and 10 to 19°C
13 (50 to 66°F) is suitable for upstream migration of fall-run Chinook Salmon. In a
14 review of available literature, Marine (1992) reported a water temperature range
15 of 6 to 14°C (43 to 57°F) as optimal for pre-spawning broodstock survival,
16 maturation, and spawning for adult Chinook Salmon.

17 **9B.5.2.2 Spawning**

18 Most Chinook Salmon spawn in larger rivers or tributaries, although spawning
19 has been observed in streams as small as 7 to 10 ft (2 to 3 m) wide (Vronskiy
20 1972). Chinook Salmon typically spawn in low- to moderate-gradient reaches of
21 streams, but can navigate shorter reaches with steeper gradients to access suitable
22 spawning areas. Armantrout (ULEP 1998) concluded that Chinook Salmon
23 seldom inhabit streams with gradients greater than 3 percent after examining
24 extensive inventory data from Oregon. The upper extent of Chinook Salmon
25 distribution in the Umpqua River basin in Oregon appears to occur where
26 gradients are less than 3 percent (ULEP 1998).

27 Upon arrival at the spawning grounds, adult females dig shallow depressions or
28 pits (redds) in suitably sized gravels (discussed in further detail below), deposit
29 eggs in the bottom during the act of spawning, and cover them with additional
30 gravel. Over a period of one to several days, the female gradually enlarges the
31 redd by digging additional pits in an upstream direction (Burner 1951). Redd
32 areas vary considerably depending on female size, substrate size, and water
33 velocities, and can range from 5.4 (Neilson and Banford 1983) to 482 ft² (0.5 to
34 44.8 m²) (Chapman et al. 1986).

35 Chinook Salmon tend to seek spawning sites with high rates of intergravel flow.
36 Upwelling, which is associated with a concave bed profile, may be an important
37 feature selected by spawning Chinook Salmon (Vaux 1968).

38 Chinook Salmon are capable of spawning within a wide range of water depths and
39 velocities, provided that intergravel flow is adequate for delivering sufficient
40 oxygen to eggs and alevins (Healey 1991). Depths most often recorded for
41 Chinook Salmon redds range from 4 to 80 inches (10 to 200 cm) (Burner 1951,
42 Chambers et al. 1955, Vronskiy 1972), and velocities range from 0.5 to 3.3 ft/s
43 (15 to 100 cm/s) (Burner 1951, Chambers et al. 1955, Thompson 1972, Vronskiy
44 1972, Smith 1973), although values may vary between races and stream basins.

1 Fall-run Chinook Salmon, for instance, are able to spawn in deeper water with
 2 higher velocities such as the mainstem Sacramento River because of their larger
 3 size (Hallock et al. 1957).

4 Substrate particle size composition has been shown to have a significant influence
 5 on intragravel flow dynamics (Platts et al. 1979). Chinook Salmon may therefore
 6 have evolved to select redd sites with specific particle size criteria that will ensure
 7 adequate delivery of dissolved oxygen to their incubating eggs and developing
 8 alevins. In addition, salmon are limited by the size of substrate that they can
 9 physically move during the redd building process. Substrates selected likely
 10 reflect a balance between water depth and velocity, substrate composition and
 11 angularity, and fish size. As depth, velocity, and fish size increase, Chinook
 12 Salmon are able to displace larger substrate particles. D50 values (the median
 13 diameter of substrate particles found within a redd) for spring-run Chinook have
 14 been found to range from 10.8 to 78.0 mm (0.43 to 3.12 inches) (Platts et al.
 15 1979; Chambers et al. 1954, 1955).

16 In 1997, USFWS researchers collected data on substrate particle size, velocity,
 17 and depth at hundreds of Chinook Salmon redds in the Sacramento River between
 18 Keswick Dam and Battle Creek to develop habitat suitability criteria for use in
 19 models that can aid in determining instream flows beneficial for anadromous
 20 salmonids. Redds in both shallow and deep areas were sampled. Table 9B.1
 21 summarizes habitat suitability criteria data collected in this study for three of the
 22 four runs (too few spring-run redds were found from which to collect data).
 23 Much more detail on the methods used and results can be found in USFWS
 24 (2003).

25 **Table 9B.1 Range of Suitable Habitat Values for Chinook Salmon Spawning in the**
 26 **Sacramento River (USFWS 2003)**

Run	Range of Suitable Values Velocity ft/s	Range of Suitable Values Velocity m/s	Range of Suitable Values Depth ft	Range of Suitable Values Depth m	Range of Suitable Values Substrate in	Range of Suitable Values Substrate cm
Fall	0.93 to 2.66	0.28 to 0.81	1–14	0.3–4	1–3 to 3–5	3–8 to 8–13
Late fall	0.90 to 2.82	0.27 to 0.86	1–14	0.3–4	1–3 to 4–5	3–8 to 10–13
Winter	1.54 to 4.10	0.47 to 1.25	3–16	0.9–5	1–3 to 3–5	3–8 to 8–13

27 **9B.5.2.3 Egg Incubation and Alevin Development**

28 Once redd construction is completed, a key determinant of survival from egg
 29 incubation through fry emergence is the amount of fine sediment in the gravel
 30 (McCuddin 1977; Reiser and White 1988). High concentrations of fine sediment
 31 in (or on) a streambed can reduce permeability and intergravel flow within the
 32 redd. This can result in reduced delivery rate of oxygen and increasingly elevated
 33 metabolic waste levels around incubating eggs, larvae, and sac-fry as they
 34 develop within egg pockets (Kondolf 2000), which can in turn lead to high
 35 mortality. Several studies have correlated reduced dissolved oxygen levels with

1 mortality, impaired or abnormal development, delayed hatching and emergence,
2 and reduced fry size at emergence in anadromous salmonids (Wickett 1954,
3 Alderdice et al. 1958, Coble 1961, Silver et al. 1963, McNeil 1964a, Cooper
4 1965, Shumway et al. 1964, Koski 1981). Silver et al. (1963) found that low
5 dissolved oxygen concentrations are related to mortality and reduced size in
6 Chinook Salmon and steelhead embryos. Fine sediments in the gravel interstices
7 can also physically impede fry emergence, trapping (or entombing) them within
8 the redd (Phillips et al. 1975, Hausle and Coble 1976).

9 The effects of high fine sediment concentrations may be counteracted to a certain
10 extent by the redd construction process itself. As adult salmon build redds, they
11 displace fine material downstream and coarsen the substrate locally (Kondolf
12 et al. 1993, Peterson and Foote 2000, Moore et al. 2004). However, the effects of
13 sediment reduction during redd construction may be rapidly reversed by
14 infiltration of fine sediment into the redds during the incubation period (Kondolf
15 et al. 1993).

16 Suitable water temperatures are required for proper embryo development and
17 emergence. Incubating Chinook Salmon eggs can withstand constant
18 temperatures between 35.1 (Combs and Burrows 1957) and 62.1°F (1.7 and
19 16.7°C) (USFWS 1999); however, substantial mortality may occur at the
20 extremes. Myrick and Cech (2004) conclude that temperatures between 43 and
21 54°F (6 and 12°C) are best for ensuring egg and alevin survival. Sublethal stress
22 and/or mortality of incubating eggs resulting from elevated temperatures would be
23 expected to begin at temperatures of about 58°F (14.4°C) for constant exposures
24 (Combs and Burrows 1957, Combs 1965, Healey 1979).

25 Some have suggested that the eggs and fry of winter-run Chinook Salmon may be
26 slightly more tolerant of warm water temperatures than those of fall-run Chinook
27 Salmon. One study by USFWS (1999) showed fall-run Chinook Salmon egg
28 mortality increasing at lower temperatures (53.6°F [12°C]) than winter-run
29 (56.0°F [13.3°C]). Greater tolerance to temperature was also observed in the
30 post-hatching period, as was also found by Healey (1979). According to Myrick
31 and Cech (2001), however, temperature tolerances of winter-run eggs and fry
32 generally agree with those found for populations in more northern regions, and
33 there does not appear to be much variation, if any, with regard to egg thermal
34 tolerances between runs of Chinook Salmon (Healey 1979, Myrick and Cech
35 2001).

36 **9B.5.2.4 Fry Rearing**

37 Following emergence, fry occupy low-velocity, shallow areas near stream
38 margins, including backwater eddies and areas associated with bank cover such as
39 large woody debris (Lister and Genoe 1970, Everest and Chapman 1972, McCain
40 1992). As the fry grow, they tend to move into deeper and faster water further
41 from banks (Hillman et al. 1987, Everest and Chapman 1972, Lister and Genoe
42 1970). Everest and Chapman (1972) suggests that habitat with water velocities
43 less than 0.5 ft/s (15 cm/s) and depths less than 24 inches (60 cm) are suitable for
44 newly emerged fry.

1 Although fry typically drift downstream following emergence (Healey 1991),
2 movement upstream or into cooler tributaries following emergence has also been
3 observed in some systems (Lindsay et al. 1986, Taylor and Larkin 1986). On the
4 Sacramento River, juvenile Chinook Salmon are more commonly found in
5 association with natural banks and shaded riparian cover than banks stabilized
6 with riprap (DFG 1983; Michny and Hampton 1984; Michny and Deibel 1986;
7 Michny 1987, 1988, 1989; Fris and DeHaven 1993). DeHaven (1989) found this
8 association to be weaker at lower water temperatures than at temperatures over
9 70°F (21°C).

10 **9B.5.2.5 Juvenile Rearing**

11 Little is known regarding habitat selection of juvenile Chinook Salmon in the
12 Sacramento River system specifically. Habitat preferences of Chinook Salmon
13 may vary depending on channel confinement, substrate and bank characteristics,
14 abundance of small and large wood, presence of other salmonids (particularly
15 Coho Salmon), and whether the Chinook display an ocean- or stream-type life
16 history. Juvenile habitat use may also change seasonally, diurnally, or as a
17 function of growth, with larger juveniles tending to occupy habitats with higher
18 water velocities.

19 Several researchers have shown relationships between velocity and juvenile
20 Chinook Salmon habitat use, with juveniles generally occupying areas with water
21 velocities less than 15 to 30 cm/s (Thompson 1972, Hillman et al. 1987, Steward
22 and Bjornn 1987, Murphy et al. 1989, Beechie et al. 2005), as well as a preference
23 for areas with cover provided by brush, large wood, or undercut banks (Hillman
24 et al. 1987, Johnson et al. 1992, Beechie et al. 2005). Lister and Genoe (1970)
25 found that juvenile Chinook Salmon preferred “slow water adjacent to faster
26 water (40 cm/s),” and Shirvell (1994) suggested that preferred habitat locations
27 vary by activity. For feeding, they are likely to select positions with optimal
28 velocity conditions, whereas for predator avoidance, optimal light conditions are
29 more likely to be important (Shirvell 1994). At night, juvenile Chinook Salmon
30 appear to move to quiet water or pools and settle to the bottom, returning the next
31 day to the riffle and glide habitats they had occupied the previous day
32 (Edmundson et al. 1968, Chelan County Public Utility District 1989).

33 Although some researchers have found juvenile Chinook Salmon to reside
34 primarily in pools, they may also use glides and runs as well as riffles. Chinook
35 Salmon may prefer deeper pools with low water velocities during spring and
36 summer as well as during winter (Lister and Genoe 1970, Everest and Chapman
37 1972, Swales et al. 1986, Hillman et al. 1987). In the Elk River in Oregon,
38 Burnett and Reeves (2001) found most juvenile ocean-type Chinook Salmon (in
39 sympatry with Coho Salmon and steelhead) in valley segments with deeper pools,
40 larger volume pools, and pools with greater densities of large wood. In Elk River
41 tributaries, the juveniles were observed almost exclusively in pools. Roper et al.
42 (1994) also found age-0+ Chinook to be strongly associated with pools in the
43 South Umpqua River basin in Oregon. In the Sacramento and American rivers,
44 CDFG (1997) found juvenile Chinook Salmon densities to be highest in runs,
45 closely followed by pools, with fish also occupying riffles and glides.

9B.5.2.6 Summer Rearing

Juvenile growth rates are an important influence on survival because juvenile salmon are gape-limited predators that are themselves subject to gape-limited predation by larger fish. Thus, faster growth both increases the range of food items available to them and decreases their vulnerability to predation (Myrick and Cech 2004). Temperatures have a significant effect on juvenile Chinook Salmon growth rates. On maximum daily rations, growth rate increases with temperature to a certain point and then declines with further increases. Reduced rations can also result in reduced growth rates; therefore, declines in juvenile salmonid growth rates are a function of both temperature and food availability. Laboratory studies indicate that juvenile Chinook Salmon growth rates are highest at rearing temperatures from 65 to 70°F (18.3 to 21.1°C) in the presence of unlimited food (Clarke and Shelbourn 1985, Banks et al. 1971, Brett et al. 1982, Rich 1987), but decrease at higher temperatures. Myrick and Cech (2004) note that two studies have been published on the relationship between temperature and growth of Central Valley Chinook Salmon—one by Marine and Cech (2004) on Sacramento River fall-run Chinook Salmon, and one by Myrick and Cech (2002) on American River fall-run Chinook Salmon. Provided that food is not limited, these studies showed that optimum temperatures for growth were between 63 and 68°F (17 and 20°C). Under natural conditions, it is unlikely that Chinook Salmon will feed at 100 percent rations, and disease, competition, and predation are also factors that may affect survival. To determine temperatures that might be optimal for growth of juvenile Chinook under natural conditions, Brett et al. (1982) used a value of 60 percent rations, based on field studies that suggested fish in the wild fed at roughly 60 percent of their physiological maximum. When used in a model developed for sockeye salmon, Brett determined that juvenile Chinook Salmon would reach their optimal growth at a temperature of about 59°F (15°C) (Brett et al. 1982). Nicholas and Hankin (1989) suggest that the duration of freshwater rearing is tied to water temperatures, with juveniles remaining longer in rivers with cool water temperatures.

Temperatures of greater than 74°F (23.3°C) are considered potentially lethal to juvenile Chinook Salmon (State Water Contractors 1990). Myrick and Cech (2004) summarized available information on juvenile Chinook Salmon temperature tolerances. Incipient upper lethal temperature (IULT) studies, which may be the most biologically relevant for studying juvenile temperature tolerances, are lacking for Central Valley Chinook Salmon. Sacramento River fall-run Chinook Salmon were reared at temperatures between 70 and 75°F (21 and 24°C) by Marine and Cech (2004) without significant mortality; however, Rich (1987) observed significant mortality after only 8 days of rearing at 75°F (24°C) (Myrick and Cech 2004). Myrick and Cech (2004) suggests that, until IULT studies are conducted on Central Valley Chinook Salmon, managers use Brett's (1952) and Brett et al.'s (1982) data on more northern Chinook Salmon, which determined that the IULT is in the range of 24 to 25°C (75 to 77°F). More detail on temperature tolerances of various Chinook life stages can be found in Myrick and Cech (2001, 2004).

1 Chronic exposure to high temperatures may result in greater vulnerability to
2 predation. Marine (1997) found that Sacramento River fall-run Chinook Salmon
3 reared at the highest temperatures (21 to 24°C [70 to 75°F]) were preyed upon by
4 Striped Bass more often than those reared at low or moderate temperatures.
5 Consumption rates of piscivorous fish such as Sacramento pikeminnow, Striped
6 Bass, and largemouth bass increase with temperature, which may compound the
7 effects of high temperature on juvenile and smolt predation mortality.

8 **9B.5.2.7 Winter Rearing**

9 Juvenile Chinook Salmon rearing in tributaries may disperse downstream into
10 mainstem reaches in the fall and take up residence in deep pools with LWD, in
11 interstitial habitat provided by boulder and rubble substrates, or along river
12 margins (Swales et al. 1986, Healey 1991, Levings and Lauzier 1991). During
13 high flow events, juveniles have been observed to move to deeper areas in pools,
14 and they may also move laterally in search of slow water (Shirvell 1994, Steward
15 and Bjornn 1987). Hillman et al. (1987) found that individuals remaining in
16 tributaries to overwinter chose areas with cover and low water velocities, such as
17 areas along well-vegetated, undercut banks. There is very little information
18 available on Chinook Salmon use of floodplains and off-channel habitats such as
19 sloughs and oxbows compared to Coho Salmon. However, studies in the
20 Sacramento and Cosumnes rivers have shown that shallow, seasonally inundated
21 floodplains can provide suitable rearing habitat for Chinook Salmon.

22 In winter, juvenile Chinook Salmon may make use of the interstitial spaces
23 between coarse substrates as cover (Bjornn 1971, Hillman et al. 1987). Hillman
24 et al. (1987) found that the addition of cobble substrate to heavily sedimented
25 glides in the fall substantially increased winter rearing densities, with juvenile
26 Chinook Salmon using the interstitial spaces between the cobbles as cover. Fine
27 sediment can act to reduce the value of gravel and cobble substrate as winter
28 cover by filling interstitial spaces between substrate particles. This may cause
29 juveniles to avoid these embedded areas and move elsewhere in search of suitable
30 winter cover (Stuehrenberg 1975, Hillman et al. 1987).

31 Over much of the Chinook Salmon's range, winter temperatures are too cold to
32 allow for much growth in the winter. The low-temperature threshold for positive
33 growth in juvenile Chinook Salmon is believed to be about 40.1°F (4.5°C), with
34 39.4°F (4.1°C) being the lower limit for zero net growth in a juvenile Chinook
35 Salmon population (Armour 1990). In the Sacramento River, water temperatures
36 rarely fall below 43°F (6°C), however, allowing for growth throughout the winter.

37 Within the action area, where juvenile Chinook Salmon are rearing in mainstem
38 channels downstream of reservoirs, water temperatures rarely fall below 43°F
39 (6°C), allowing for growth throughout the winter months. Under these
40 conditions, habitat shifts are less related to seasonal temperature changes and
41 more strongly affected by growth (i.e., as individuals grow, they can take
42 advantage of habitats with stronger flow and are better able to escape predation).

1 In the Sacramento/San Joaquin system, some juvenile Chinook Salmon rear on
2 seasonally inundated floodplains in the winter. Sommer et al. (2001) found
3 higher growth and survival rates of juveniles that reared on the Yolo Bypass
4 floodplain than in the mainstem Sacramento River, and Moyle (2000) observed
5 similar results on the Cosumnes River floodplain. On the Yolo Bypass,
6 bioenergetic modeling suggested that increased prey availability on the floodplain
7 was sufficient to offset increased metabolic demands from higher water
8 temperatures (9°F [5°C] higher than mainstem). The Yolo Bypass has a relatively
9 smooth topography with few pits and depressions, which possibly enhances its
10 value as floodplain rearing habitat by reducing stranding mortality as floodwaters
11 recede and juvenile salmon return to the main stem (Sommer et al. 2001).

12 **9B.5.2.8 Smoltification and Outmigration**

13 Juveniles of all four runs of Chinook Salmon in the Central Valley must pass
14 through the Sacramento-San Joaquin Delta and San Francisco Bay Estuary on
15 their way to the ocean, and many rear there for varying periods prior to ocean
16 entry. Williams (2012) found evidence that many naturally produced fall-run
17 Chinook Salmon that survived to return as adults had left freshwater at lengths
18 greater than 55 mm, while juvenile Chinook Salmon from other Central Valley
19 runs were older and larger upon entering the estuary and likely passed through it
20 more quickly (Williams 2012).

21 In many systems within the species' distribution, juvenile Chinook Salmon spend
22 up to several months in estuaries feeding and growing before entering the ocean
23 (Healey 1991); in productive estuaries, this strategy can result in ocean entry at a
24 larger size with a higher chance of survival, presumably by reducing predation at
25 this critical juncture. Although wetlands and floodplains may have been
26 extensive enough in the Delta under historical conditions (Atwater et al. 1979) to
27 support high juvenile production in an environment where there were fewer
28 predators, Delta marsh habitats and native fish communities have undergone such
29 extreme changes from historical conditions (Kimmerer et al. 2008) that few
30 locations in the eastern and central Delta currently provide suitable habitat for
31 rearing Chinook Salmon. For example, substantial numbers of fry may be found
32 in the Delta from January through March, but relatively few were found in the
33 remaining months of the year during sampling from 1977 to 1997 (Brandes and
34 McLain 2001). The annual abundance of fry (defined as less than 2.8 inches
35 [70 mm] fork length) in the Delta during this period appears related to flow, with
36 the highest numbers observed in wet years (Brandes and McLain 2001).

37 Although growth rates of juvenile Chinook Salmon may be high at temperatures
38 approaching 66°F (19°C), cooler temperatures may be required for Chinook
39 Salmon to successfully complete the physiological transformation from parr to
40 smolt. Smoltification in juvenile Sacramento River fall-run Chinook Salmon was
41 studied by Marine (1997), who found that juveniles reared under a high
42 temperature regime of 70 to 75°F (21 to 24°C) exhibited altered and impaired
43 smoltification patterns relative to those reared at low 55 to 61°F (13 to 16°C) and
44 moderate 63 to 68°F (17 to 20°C) temperatures. Some alteration and impairment
45 of smoltification was also seen in the juveniles reared at moderate temperatures.

1 **9B.5.3 Winter-Run Chinook Salmon**

2 **9B.5.3.1 Legal Status**

3 Federal: Endangered, Designated Critical Habitat

4 State: Endangered

5 Although Chinook Salmon range from California's Central Valley to Alaska and
6 the Kamchatka Peninsula in Asia, winter-run Chinook Salmon are only found in
7 the Sacramento River. Chinook Salmon of this race are unique because they
8 spawn during the summer months when air temperatures usually approach their
9 yearly maximum. As a consequence, winter-run Chinook Salmon require stream
10 reaches with cold water sources that will protect embryos and juveniles from the
11 warm ambient conditions in the summer. Historically, high-elevation reaches of
12 tributaries to the upper Sacramento River (e.g., McCloud River) provided the cold
13 water reaches that supported summer spawning by winter-run Chinook Salmon.
14 Currently, hypolimnetic releases from Shasta Lake provide the cold water
15 temperatures that allow winter-run Chinook Salmon to persist downstream of the
16 dam, despite the complete loss of historical spawning habitat, access to which was
17 cut off upon completion of Shasta Dam (1963).

18 The California-Nevada chapter of the American Fisheries Society petitioned
19 NMFS to list the run as a threatened species in 1985 (AFS 1985) and, following a
20 dangerously low year-class in 1989, NMFS issued an emergency listing for
21 Sacramento River winter-run Chinook Salmon as a threatened species (NMFS
22 1989); the California Fish and Game Commission listed the winter run as
23 endangered in the same year. After several years of low escapements in the early
24 1990s, the status of winter-run was changed from threatened to endangered by
25 NMFS in 1994, which was reaffirmed in 2005 and 2011 (NMFS 1994, 2005,
26 2011).

27 The ESU includes fish that are propagated as part of a conservation hatchery
28 program managed by the USFWS at Livingston Stone National Fish Hatchery
29 (LSNFH). Since 2000, the proportion of the ESU spawning in the Sacramento
30 River that are of hatchery origin has generally ranged from 5 to 10 percent of the
31 total population, but reached a high of 20 percent in 2005 (NMFS 2011).
32 USFWS's goal is to manage the LSNFH program such that hatchery origin fish
33 are less than 20 percent of total in-river escapement. Hatchery fish were
34 estimated to be 12 percent of the total in-river spawners in 2010, based on carcass
35 surveys (DFG 2010). Over the last 10 years, hatchery returns have averaged
36 8 percent of total escapement (NMFS 2011).

37 Critical habitat was designated as the Sacramento River from Keswick Dam at
38 river mile (RM) 302 to Chipps Island (RM 0) at the westward margin of the
39 Delta; all waters from Chipps Island westward to the Carquinez Bridge, including
40 Honker Bay, Grizzly Bay, Suisun Bay, and the Carquinez Strait; all waters of
41 San Pablo Bay westward of the Carquinez Bridge; and all waters of San Francisco
42 Bay (north of the San Francisco-Oakland Bay Bridge) to the Golden Gate Bridge
43 (NMFS 1993).

1 9B.5.3.1.1 Distribution

2 Winter-run Chinook Salmon are found only in the Sacramento River basin. The
3 distribution of winter-run Chinook Salmon spawning has shifted over time in
4 response to changes in upstream passage caused by water supply development
5 and operations. Prior to construction of Shasta Dam in the 1940s, winter-run
6 Chinook Salmon spawned in the upper Sacramento River system (in the Little
7 Sacramento, McCloud, and possibly Pit and Fall rivers) and in nearby Battle
8 Creek (Yoshiyama et al. 1998). Since the construction of Shasta Dam, winter-run
9 Chinook Salmon have been limited to the mainstem Sacramento River below
10 Keswick Dam (RM 302), although a few adults occasionally stray into tributaries
11 (e.g., Battle and Mill creeks) to spawn (Harvey-Arrison 2001). The distribution
12 of spawning likely shifted again in 1966, when the construction and operation of
13 RBDD (RM 243.5) impeded access to upstream reaches, forcing more winter-run
14 adults to spawn downstream of the diversion dam. A radio-tag survey of winter-
15 run adults between 1979 and 1981 indicated that adults were delayed at RBDD
16 between 1 and 40 days, with an average delay of 18 days (Hallock and Fisher
17 1985). The dam also forced winter-run adults to spawn downstream of Red Bluff,
18 where summer water temperatures were frequently too high to support successful
19 egg incubation and emergence. Beginning in 1986, the Bureau of Reclamation
20 (Reclamation) began raising RBDD gates during the winter to facilitate upstream
21 passage of winter-run Chinook (Reclamation 2004), which precipitated an
22 upstream shift in the distribution of winter-run spawning. In 2012, the RBDD
23 gates were opened to allow year-round passage.

24 Until 2001, most winter-run spawning occurred downstream of ACID Dam
25 (RM 298.4); however, an improvement of this dam's fish passage facilities in
26 2001 allowed another upstream shift in the distribution of spawning (DFG 2002a,
27 2004).

28 9B.5.3.1.2 Life History and Habitat Requirements

29 General habitat requirements for Chinook Salmon are described above; the
30 following describes life history strategies and habitat requirements unique to the
31 winter-run or of primary importance to its life history. The winter-run Chinook
32 Salmon's life history is unique to the Sacramento River because it provides the
33 thermal conditions that allow for the success of this strategy. Because winter-run
34 Chinook Salmon spawn in late spring and early summer, they require access to
35 stream reaches with summer water temperatures cool enough to allow egg
36 incubation. The spawning reaches and reaches downstream have sufficiently
37 warm water temperatures to support growth throughout the winter, allowing
38 juveniles to grow large enough to smolt and outmigrate before water temperatures
39 become too high the following spring and summer. This life-history strategy
40 reduces competition for spawning habitat with other runs. However, it also makes
41 the run reliant on year-round coldwater sources, which limits the potential for
42 expanding the range of the run in the Sacramento River basin.


43 Table 9B.2 illustrates life history timing for winter-run Chinook Salmon in the
44 Sacramento River basin. Winter-run Chinook Salmon display a life history that is

1 intermediate between ocean-type and stream-type. They spend between 5 and
 2 10 months rearing in fresh water before migrating to sea, which is longer than for
 3 typical ocean-type Chinook Salmon, but shorter than for other stream-type
 4 Chinook Salmon (Healey 1991).

5 **Table 9B.2 Life History Timing of Winter-run Chinook Salmon in the Sacramento**
 6 **River Basin**

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Adult entry into San Francisco Bay ^a												
Migration past RBDD ^b												
Spawning ^c												
Incubation ^c												
Fry emergence ^c												
Rearing (age 0+)												
Presence at CVP/SWP salvage facilities ^c												
Outmigration toward and through the Delta ^c												

7 Notes:
 8 a. Van Woert 1958; Hallock et al. 1957
 9 b. Hallock and Fisher 1985
 10 c. NMFS 2012 (unpubl. data)



11 **9B.5.3.1.3 Adult Upstream Migration and Spawning**

12 Adult winter-run Chinook Salmon enter San Francisco Bay from November
 13 through June (Van Woert 1958, Hallock et al. 1957). Migration past RBDD
 14 begins in mid-December and can continue into early August, but the majority of
 15 winter-run adults migrate past RBDD between January and May, with a peak in
 16 mid-March (Hallock and Fisher 1985). In recent years, upstream passage of
 17 winter-run adults at RBDD was addressed by raising the gates between
 18 September 15 and May 15, which encompasses the vast majority of the upstream
 19 migration period for winter-run Chinook Salmon. As of 2012, the gates at RBDD
 20 are open year-round to allow for upstream passage.

1 Like spring-run Chinook Salmon, winter-run Chinook Salmon enter spawning
2 streams while still reproductively immature. Adults hold for a few months in
3 deep pools near spawning areas, which provides time for gonadal development.
4 Spawning occurs from mid-April to mid-August, peaking in May and June, in the
5 Sacramento River reach between Keswick Dam and RBDD (Reclamation 1991).
6 With the changes in RBDD gate operations, volitional spawning below RBDD is
7 negligible in most years. Since fish passage improvements were completed at the
8 ACID Dam in 2001, winter-run Chinook Salmon spawning has shifted upstream.
9 The majority of winter-run Chinook Salmon in recent years (i.e., more than
10 50 percent since 2007) spawn in the area from Keswick Dam to the ACID Dam
11 (approximately 5 miles) (NMFS 2009).

12 **9B.5.3.1.4 Juvenile Rearing and Outmigration**

13 Winter-run fry emerge from the spawning gravels from mid-June through mid-
14 October (NMFS 1997). Because spawning is concentrated upstream in the
15 reaches below Keswick Dam, the entire Sacramento River can serve as a nursery
16 area for juveniles as they migrate downstream. Emigrating juvenile Sacramento
17 River winter-run Chinook Salmon pass the RBDD beginning as early as mid-July,
18 typically peaking in September, and can continue through March in dry years
19 (Reclamation 1991, NMFS 1997). Many juveniles apparently rear in the
20 Sacramento River below RBDD for several months before they reach the Delta
21 (Williams 2006). From 1995 to 1999, all Sacramento River winter-run Chinook
22 Salmon outmigrating as fry passed the RBDD by October, and all outmigrating
23 presmolts and smolts passed the RBDD by March (Martin et al. 2001).

24 Juvenile Sacramento River winter-run Chinook Salmon occur in the Delta
25 primarily from November through early May based on data collected from trawls
26 in the Sacramento River at West Sacramento, although the overall timing may
27 extend from September to early May (NMFS 2012). The timing of migration
28 varies somewhat because of changes in river flows, dam operations, seasonal
29 water temperatures, and hydrologic conditions (water year type). Winter-run
30 Chinook Salmon juveniles remain in the Delta until they are between 5 and
31 10 months of age, after reaching a fork length of approximately 118 mm. Distinct
32 emigration pulses from the Delta appear to coincide with periods of high
33 precipitation and increased turbidity (Del Rosario et al. 2013).

34 The entire population of the Sacramento River winter-run Chinook Salmon passes
35 through the Delta as migrating adults and emigrating juveniles. Because winter-
36 run Chinook Salmon use only the Sacramento River system for spawning, adults
37 are likely to migrate upstream primarily along the western edge of the Delta
38 through the Sacramento River corridor. Juveniles likely use a wider area within
39 the Delta for migration and rearing than adults; juvenile winter-run salmon have
40 been collected at various locations in the Delta, including the SWP and CVP
41 south Delta export facilities. Studies using acoustically tagged juvenile and adult
42 Chinook Salmon are ongoing to further investigate the migration routes,
43 migration rates, reach-specific mortality rates, and the effects of hydrologic
44 conditions (including the effects of SWP/CVP export operations) on salmon
45 migration through the Delta. Tagging studies have indicated that juvenile salmon

1 entering the interior Delta via the Delta Cross Channel and Georgiana Slough
2 survive at a lower rate than fish migrating within the Sacramento River (Newman
3 and Brandes 2010; Perry et al. 2010, 2012). Juvenile winter-run Chinook Salmon
4 likely inhabit Suisun Marsh for rearing and may inhabit the Yolo Bypass when
5 flooded, although use of these two areas is not well understood.

6 **9B.5.3.1.5 Population Trends**

7 There is little historical data available to characterize winter-run Chinook Salmon
8 escapements prior to the construction of Shasta Dam; indeed, the agencies did not
9 recognize winter-run Chinook Salmon as a distinct run until the 1940s (Needham
10 et al. 1943). In the late 1930s, the pending construction of Shasta Dam prompted
11 the agencies to commission a study of potential salmon salvage options. As part
12 of this investigation, researchers placed a counting weir at ACID Dam between
13 1937 and 1939 to estimate the size of the salmon run in the Sacramento River
14 (Hatton 1940). The counting weir enabled scientists to estimate the run size of
15 the fall-run Chinook Salmon populations; however, the removal of flashboards
16 from the ACID Dam during winter prevented observations of winter-run Chinook
17 Salmon during their period of upstream migration (December–May).

18 There were no direct observations of winter-run Chinook Salmon spawning in the
19 mainstem Sacramento River between 1943 and 1946—the first years when the
20 construction of Shasta Dam blocked upstream passage. Nevertheless, incidental
21 observations of winter-run salmon during trap-and-haul operations for spring-run
22 salmon, coupled with poor environmental conditions in the Sacramento River and
23 Deer Creek, led Slater to conclude that “the winter-run populations were small” in
24 the years when Shasta Dam was being constructed (1963).

25 Slater (1963) hypothesized that the winter-run salmon population began to
26 rebound in 1947, and that “this initial recovery seems to have been both
27 substantial and rapid” from the “low point of 1943–1946.” He cites an angling
28 survey conducted by Smith (1950), which evaluated the 1947–1948 and 1949–
29 1950 sport fishery in the upper Sacramento River. “Increased catches of winter-
30 run Chinook Salmon in January and February 1949” (Slater 1963) led Smith
31 (1950) to conclude that a “sizable” winter-run population existed. Similarly,
32 Slater cited an increase in the number of winter-run salmon that were harvested
33 by Coleman National Fish Hatchery between 1949 and 1956 (as part of the fall-
34 run salmon propagation program) (Azevedo and Parkhurst 1958) as evidence that
35 winter-run salmon escapements increased in the late 1940s and early 1950s.
36 Although these qualitative assessments do not permit a detailed tracking of
37 winter-run salmon abundance, they do suggest a positive trend in the population
38 in the years after Shasta Dam was completed.

39 This positive trend seems to have continued through the 1950s, because Hallock
40 estimated that 11,000 winter-run adults were harvested from the Sacramento
41 River by anglers in the winter of the 1961–1962 fishing season (Slater 1963).
42 Hallock’s estimate of the percentage of winter-run Chinook Salmon caught in the
43 in-river recreational harvest suggests that total winter-run escapements in the
44 winter of 1961–1962 numbered in the tens of thousands. In June 1963, Slater

1 personally observed winter-run Chinook Salmon spawning in the vicinity of
2 Redding in numbers that approached the fall-run population that spawned in the
3 same sites (Slater 1963). For context, the four years before Slater's observation
4 of winter-run spawning in 1963 (1959–1962) had fall-run salmon escapement
5 estimates ranging from 115,500 to 250,000 salmon. Although Slater observed
6 spawning in only a small portion of the habitat available to both winter-run and
7 fall-run salmon in the Sacramento River, his observation suggests that the winter-
8 run salmon population had increased substantially from the few hundred fish
9 captured during the trap-and-haul salvage operation in 1943 and 1945. His
10 observation also suggests that the winter-run salmon population had recovered
11 from a probable year-class failure in 1943 and a partial year-class failure in 1944.

12 Beginning in 1967, agency biologists began estimating annual winter-run
13 escapements by monitoring adults migrating through the fish passage facilities of
14 RBDD. Although the dam facilitated a more accurate account of the winter-run
15 population, gate operations interfered with upstream passage. Gate operations
16 were modified beginning in winter 1986 to facilitate the upstream passage of
17 winter-run Chinook Salmon. However, raising the dam gates rendered winter-run
18 escapement estimates less reliable, because migrating salmon could bypass the
19 dam's fish counting facilities.

20 The RBDD counts permitted agency biologists to track the decline in winter-run
21 Chinook abundance beginning in the 1970s. The drought of 1976–1977 caused a
22 precipitous decline in abundance between 1978 and 1979, when escapements fell
23 below 2,500 fish. Population abundance remained very low through the mid-
24 1990s, with adult abundance in some years less than 500 fish (DFW 2014).

25 Beginning in the mid-1990s and continuing through 2006, adult escapement
26 showed a trend of increasing abundance, approaching 20,000 fish in 2005 and
27 2006. However, recent population estimates of winter-run Chinook Salmon
28 spawning upstream of the RBDD have declined since the 2006 peak. The
29 escapement estimate for 2007 through 2014 has ranged from a low of 738 adults
30 in 2011 to a high of 5,959 adults in 2013. The escapement estimate of 738 adults
31 in 2011 was the lowest total escapement estimate since the all-time low
32 escapement estimate of 144 adults in 1994. Poor ocean productivity (Lindley
33 et al. 2009), drought conditions from 2007 to 2009, and low in-river survival
34 (National Marine Fisheries Service 2011) are suspected to have contributed to the
35 recent decline in escapement of adult winter-run Chinook Salmon. Table 9B.3
36 shows winter-run Chinook Salmon natural and hatchery escapement subsequent
37 to 2004.

1 **Table 9B.3 Recent Winter-run Chinook Salmon Natural and Hatchery Escapement**

Year	Sacramento River above RBDD	Sacramento River below RBDD	Subtotal	CNFH Transfers	LSNFH Transfers	Battle Creek	Total
Dec 1990-Aug 1991	177	0	177	33	–	–	211
Dec 1991-Aug 1992	1,159	44	1,203	34	–	–	–
Dec 1992-Aug 1993	369	9	378	–	–	–	–
Dec 1993-Aug 1994	144	0	144	42	–	–	–
Dec 1994-Aug 1995	1,159	7	1,166	43	–	88	–
Dec 1995-Aug 1996	1,012	0	1,012	–	–	325	–
Dec 1996-Aug 1997	836	0	836	–	–	44	–
Dec 1997-Aug 1998	2,831	62	2,893	–	99	–	–
Dec 1998-Aug 1999	3,264	0	3,264	–	24	–	–
Dec 1999-Aug 2000	1,261	0	1,261	–	89	2	–
Dec 2000-Aug 2001	8,085	35	8,120	–	104	–	–
Dec 2001-Aug 2002	7,325	12	7,337	–	104	–	–
Dec 2002-Aug 2003	8,105	28	8,133	–	85	–	–
Dec 2003-Aug 2004	7,784	0	7,784	–	85	–	–
Dec 2004-Aug 2005	15,730	0	15,730	36	109	0	15,875
Dec 2005-Aug 2006	17,157	48	17,205	5	93	6	17,304
Dec 2006-Aug 2007	2,487	0	2,487	1	54	0	2,542
Dec 2007-Aug 2008	2,725	0	2,725	0	105	0	2,830

Appendix 9B: Aquatic Species Life History Accounts

Year	Sacramento River above RBDD	Sacramento River below RBDD	Subtotal	CNFH Transfers	LSNFH Transfers	Battle Creek	Total
Dec 2008-Aug 2009	4,537	0	4,537	0	121	0	4,658
Dec 2009-Aug 2010	1,533	0	1,533	0	63	0	1,596
Dec 2010-Aug 2011	738	0	738	2	86	1	827
Dec 2011-Aug 2012	2,578	0	2,578	0	93	–	2,671
Dec 2012-Aug 2013	5,920	0	5,920	0	164	–	6,084
Dec 2013-Aug 2014	2,627	0	2,627	0	388	–	3,015

1 Source: DFW 2014

2 Note:

3 CNFH = Coleman National Fish Hatchery

1 Winter-run Chinook Salmon escapement to the Sacramento River in 2011 was
2 827 fish, which is the smallest number since 1994 and only 10 percent of the
3 40-year-average of approximately 8,000 fish (Azat 2012). Unusual ocean
4 conditions appear to have been affecting the ESU in the past 5 years, along with
5 other Central Valley Chinook Salmon stocks (NMFS 2011). Climate change and
6 future variations in ocean conditions, along with the many factors affecting
7 survival during freshwater life stages, may pose a serious risk to the ESU (NMFS
8 2011).

9 **9B.5.4 Central Valley Spring-Run Chinook Salmon**

10 **9B.5.4.1 Legal Status**

11 Federal: Threatened, Designated Critical Habitat

12 State: Threatened

13 Spring-run Chinook Salmon were probably the most abundant salmonid in the
14 Central Valley under historical conditions (Mills and Fisher 1994); however, large
15 dams eliminated access to vast amounts of historical habitat, and the spring run
16 has exhibited the severest declines of any of the four Chinook Salmon runs in the
17 Sacramento River basin (Fisher 1994).

18 The Central Valley spring-run Chinook Salmon ESU was federally listed as
19 threatened in 1999, and the listing was reaffirmed in 2005 when critical habitat
20 was also designated (NMFS 1999a, 2005). Spring-run Chinook Salmon was
21 listed as a threatened species under the California Endangered Species Act
22 (CESA) in February 1999. The ESU includes all naturally spawned populations
23 of spring-run Chinook Salmon in the Sacramento River and its tributaries in
24 California, including the Feather River. Feather River Hatchery spring-run
25 Chinook Salmon are also included in the ESU. This ESU largely consists of three
26 self-sustaining wild populations (i.e., Mill, Deer, and Butte creeks). Fish in these
27 streams spawn outside of the action area but pass through it on their upstream and
28 downstream migrations. Spring-run Chinook Salmon in the Feather River and
29 Clear Creek spawn within the action area.

30 Designated critical habitat for Central Valley spring-run Chinook Salmon
31 includes stream reaches of the American, Feather, Yuba, and Bear rivers;
32 tributaries of the Sacramento River, including Big Chico, Butte, Deer, Mill,
33 Battle, Antelope, and Clear creeks; and the main stem of the Sacramento River
34 from Keswick Dam through the Delta. Designated critical habitat in the Delta
35 includes portions of the Delta Cross Channel, Yolo Bypass, and portions of the
36 network of channels in the northern Delta. Critical habitat for spring-run Chinook
37 Salmon was not designated for the Stanislaus or San Joaquin rivers.

38 **9B.5.4.2 Distribution**

39 Prior to the construction of dams in the Sacramento and San Joaquin basins,
40 spring-run Chinook Salmon migrated during the spring snowmelt flows to access
41 coldwater holding and spawning habitat higher up in the basins. These steeper,
42 higher-elevation reaches are often characterized by falls and cascades that may be
43 obstacles to upstream movement of salmonids at lower flows. By migrating

1 during the high spring snowmelt flows, spring-run Chinook Salmon can also
 2 access areas above reaches that become too warm for salmon in the summer and
 3 fall, isolating them from the fall run. Thus, under historical conditions, the
 4 spring- and fall-run Chinook Salmon were geographically isolated in terms of
 5 where they spawned in the basin, which maintained their genetic integrity.

6 Spring-run Chinook Salmon once occupied all major river systems in California
 7 where there was access to cool reaches that would support oversummering adults.
 8 Historically, they were widely distributed in streams of the Sacramento-
 9 San Joaquin basin, spawning and rearing over extensive areas in the upper and
 10 middle reaches (elevations ranging from 1,400 to 5,200 ft [450 to 1,600 m]) of the
 11 San Joaquin, American, Yuba, Feather, Sacramento, McCloud, and Pit rivers
 12 (Myers et al. 1998). Spring Chinook Salmon runs in the San Joaquin River were
 13 extirpated in the mid- to late 1940s following the closure of Friant Dam and
 14 diversion of water for agricultural purposes to the San Joaquin Valley.

15 In the Sacramento River, the closure of Shasta Dam in 1945 cut off access to the
 16 spring run's major historical spawning grounds in the McCloud, Pit, and upper
 17 Sacramento rivers. This represented a loss of 70 percent of spring-run spawning
 18 habitat in the Sacramento River basin (Yoshiyama et al. 2001). Populations of
 19 spawning spring-run Chinook Salmon in the Sacramento River basin are more
 20 common in east-side tributaries to the Sacramento River upstream of the mouth of
 21 the American River. The most important spawning populations are in Deer, Mill,
 22 and Butte creeks because of their relative lack of past hatchery influence, as well
 23 as relatively stable numbers. Some spawning also takes place in Big Chico,
 24 Antelope, Cottonwood, Beegum, Clear, and Battle Creeks, and in the mainstem
 25 Sacramento River downstream of Keswick Dam and upstream of RBDD
 26 (Association of California Water Agencies and California Urban Water Agencies
 27 1997; DFG 1998, 2002b, 2012 [GrandTab data]). A spring run in the Feather
 28 River basin is maintained by hatchery production; however, the stock is believed
 29 to have been hybridized with the fall run to a great extent (Lindley et al. 2004).

30 **9B.5.4.2.1 Changes in Distribution and Hybridization with Fall** 31 **Chinook Salmon**

32 Dams have reduced or eliminated spatial segregation between spawning spring-
 33 and fall-run Chinook Salmon in some areas, particularly in the mainstem
 34 Sacramento River, leading to increased potential for hybridization on the
 35 spawning grounds. The completion of Keswick and Shasta dams in the mid-
 36 1940s blocked spring-run Chinook Salmon access to habitat in the McCloud, Pit,
 37 and Little Sacramento rivers. After construction of the dams, spring-run Chinook
 38 Salmon were forced to spawn in the mainstem Sacramento River below Keswick
 39 Dam. Historically, water temperatures would have been too high in the mainstem
 40 Sacramento River for spring-run Chinook Salmon to hold in this area during the
 41 summer. But because of hypolimnetic releases from Shasta Lake, this reach
 42 provides temperatures during the summer that are now suitable for spring-run
 43 Chinook Salmon holding and spawning, where before they were only suitable for
 44 fall-run spawning once temperatures cooled in the fall. However, coldwater
 45 releases from Shasta Dam can warm relatively rapidly during the very hot days

1 typical of the Sacramento Valley in summer and early fall. As a result, both the
2 fall and spring runs must spawn in close enough proximity to Keswick Dam to
3 benefit from these releases. The elimination of the spatial segregation that had
4 existed between the fall and spring runs results in competition between the runs
5 for the limited spawning habitat. Since fall-run Chinook Salmon spawn slightly
6 later than spring-run, spring-run redds may also be superimposed by spawning
7 fall-run fish. This may have contributed to the loss of the spring-run population,
8 along with hybridization between the two runs, as described below.

9 The majority of spring-run Chinook Salmon used to spawn upstream in tributaries
10 rather than in the mainstem Sacramento River; however, the completion and
11 operation of Shasta Dam reduced water temperatures in the main stem
12 downstream of Keswick Dam, which permitted spring-run Chinook Salmon to
13 spawn there, resulting in hybridization with fall-run stocks. Although spring-run
14 Chinook Salmon spawn earlier than fall-run, the timing of spawning of the two
15 runs overlaps enough that hybridization can occur where they share the same
16 spawning areas. Where the spring run is now forced to share spawning grounds
17 in the mainstem Sacramento River with the fall run, fall-run Chinook Salmon may
18 dominate because of their longer growth period in the ocean, slightly larger size,
19 and less time spent holding in the stream prior to spawning. Hybridization
20 between the two runs has tended to be to the detriment of the spring run life
21 history.

22 Because of this hybridization with fall-run Chinook Salmon in the mainstem
23 channel, there are considered to be only three “pure” self-sustaining populations
24 of wild spring-run Chinook Salmon remaining in Deer, Mill, and Butte creeks.

25 Similar patterns have been observed in the Feather River, where the spring run
26 historically spawned upstream of the location of Oroville Dam, and where they
27 are now forced to spawn in the same area as the fall run, as well as in the Yuba
28 and American rivers, where forced sympatry on the spawning grounds and
29 subsequent hybridization following dam construction led to DFW concluding that
30 the spring run was “extinct” in those rivers.

31 **9B.5.4.3 Life History and Habitat Requirements**

32 General habitat requirements for Chinook Salmon are described above; the
33 following describes life history strategies and habitat requirements unique to the
34 spring run or of primary importance to its life history. Spring-run Chinook
35 Salmon display a stream-type life history strategy—adults migrate upstream while
36 sexually immature, hold in deep cold pools over the summer, and spawn in late
37 summer and early fall. Juvenile outmigration is highly variable, with some
38 juveniles outmigrating in winter and spring, and others oversummering and then
39 emigrating as yearlings. Table 9B.4 illustrates life-history timing for spring-run
40 Chinook Salmon in the Sacramento River basin. The table illustrates some of the
41 changes in timing that have been observed for the run over the years, particularly
42 with regard to upstream migration and spawning.


Appendix 9B: Aquatic Species Life History Accounts

1 **Table 9B.4 Life History Timing of Spring-run Chinook Salmon in the Sacramento River Basin**

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Adult entry into Sacramento-San Joaquin Delta Estuary												
“Historical” adult migration past Red Bluff Diversion Dam ^a												
“Recent” adult migration past Red Bluff Diversion Dam ^b												
Entry into spawning tributaries (current) ^c												
Adult holding												
Historical spawning in Sacramento River basin ^d												
Spawning (Deer, Mill, Butte creeks ^e)												
Spawning (mainstem Sacramento River ^f)												
Incubation												
Fry emergence												
Fry/juvenile outmigration from tributaries ^g												
Subyearling/Yearling outmigration from tributaries ^{g, h}												
Presence at CVP/SWP salvage facilities ⁱ												
Outmigration toward and through the Delta ⁱ												
Ocean entry (yearlings)												

2 Sources: Fisher 1994; Myers et al. 1998; Hill and Weber 1999; Ward and McReynolds 2001; USFWS 2005

- 1 Notes:
- 2 a. As observed in the 1970s (Association of California Water Agencies and California Urban Water Agencies 1997)
- 3 b. As observed in the 1980s (Association of California Water Agencies and California Urban Water Agencies 1997)
- 4 c. Association of California Water Agencies and California Urban Water Agencies (1997), Hill and Webber (1999)
- 5 d. Rutter (1908), Parker and Hanson (1944)
- 6 e. Harvey (1995), Moyle et al. (1995)
- 7 f. Association of California Water Agencies and California Urban Water Agencies (1997)
- 8 g. Some spring run disperse downstream soon after emergence as fry in March and April, with others smolting after several months of rearing, and
- 9 still others remaining to oversummer and emigrate as yearlings (USFWS 1995).
- 10 h. Based on outmigrant trapping in Butte Creek in 1999 and 2000, up to 69% of age 0+ juveniles outmigrate through the lower Sacramento River
- 11 and Sacramento-San Joaquin Delta between mid-November and mid-February, with a peak in December and January (DFG 1998, Hill and Weber
- 12 1999, Ward and McReynolds 2001). A smaller number remain in Butte Creek and outmigrate in late spring or early summer; and in both Butte
- 13 and Mill creeks, some of these oversummer and outmigrate as yearlings from October to March, with a peak in November (Association of
- 14 California Water Agencies and California Urban Water Agencies 1997, Hill and Webber 1999)
- 15 i. NMFS 2012 (unpublished data)

 Period of activity
Period of peak activity

16

1 9B.5.4.3.1 Adult Upstream Migration and Spawning

2 Adult spring-run Chinook Salmon may return between the ages of 2 to 5 years.
3 Historically, adults of this run are believed to have returned predominantly at ages
4 4 and 5 years at a large size. Most spring-run Chinook Salmon now return at
5 age 3, although some portion returns at age 4 (Fisher 1994, McReynolds et al.
6 2005) probably because of intense ocean harvest (which removes the largest fish
7 from the population and selects for fish that spend fewer years at sea). In 2003,
8 an estimated 69 percent of the spring run in Butte Creek returned at age 4 (Ward
9 et al. 2004); however, in most years, the proportion of age 4 adults is much
10 smaller.

11 Adult Central Valley spring-run Chinook Salmon begin their upstream migration
12 in late January and early February (DFG 1998) and enter the Sacramento River
13 between February and September, primarily in May and June (DFG 1998, Myers
14 et al. 1998). Lindley et al. (2006) reported that adult Central Valley spring-run
15 Chinook Salmon enter native tributaries from the Sacramento River primarily
16 between mid-April and mid-June. Adults enter Deer and Mill creeks beginning in
17 March, peaking in May, and concluding in June (Vogel 1987a, 1987b;
18 Association of California Water Agencies and California Urban Water Agencies
19 1997). Their upstream migration is timed to take advantage of spring snowmelt
20 flows, which allow them access to upstream holding areas where temperatures are
21 cool enough to hold over the summer prior to the spawning season (NMFS
22 1999a). In the Sacramento River, upstream migration of spring-run Chinook
23 Salmon overlaps to a certain extent with that of winter-run Chinook Salmon; and
24 adults from particular runs are not generally distinguishable from one another by
25 physical appearance alone, making it difficult to pinpoint migration timing with
26 precision (Healey 1991).

27 Adults require large, deep pools with moderate flows for holding over the summer
28 prior to spawning in the fall. Marcotte (1984) reported that suitability of pools
29 declines at depths less than 7.9 ft (2.4 m) and that optimal water velocities range
30 from 0.5 to 1.2 ft/s (15 to 37 cm/s). In the John Day River in Oregon, spring-run
31 adults usually hold in pools deeper than 4.9 ft (1.5 m) that contain cover from
32 undercut banks, overhanging vegetation, boulders, or woody debris (Lindsay et al.
33 1986).

34 In Sacramento River tributaries, adults will pack densely in the limited holding
35 pool habitat that is available. Some fish remain to spawn at the tails of the
36 holding pools, while most move upstream to the upper watersheds to spawn, and
37 still others move back downstream to spawn. Although there are several deep
38 pools in the upper Sacramento River that may provide holding habitat for adult
39 spring-run Chinook Salmon, it is not clear which pools are heavily used. As a
40 result of cold water releases from Shasta Reservoir and natural channel
41 characteristics, numerous deep pools with suitable holding habitat are located
42 between Keswick Dam and Red Bluff (Northern California Water Association
43 and Sacramento Valley Water Users 2011).

1 Water temperatures for adult spring-run Chinook Salmon holding and spawning
2 are reportedly best when less than 60.8°F (16°C), and are lethal when greater than
3 80.6°F (27°C) (Hinze 1959, Boles et al. 1988, DFG 1998). Spring Chinook
4 Salmon in the Sacramento River typically hold in pools below 69.8 to 77°F (21 to
5 25°C). Adults may be particularly sensitive to temperatures during July and
6 August, when energy reserves are low and adults are preparing to spawn. There is
7 evidence that spring-run Chinook Salmon in the San Joaquin River were exposed
8 to high temperatures during migration and holding under historical conditions
9 (Clark 1943, Yoshiyama et al. 2001). It is possible that Central Valley spring-run
10 Chinook Salmon are adapted to tolerate warmer temperatures than other Chinook
11 Salmon stocks; however, there is no experimental evidence to confirm this
12 hypothesis, and short-term exposure to temperatures as high as 25 to 27°C (77 to
13 80.6°F) is known to be tolerated by adult Chinook Salmon (Boles et al. 1988).

14 Habitat suitability studies conducted by USFWS (2004) indicate that suitable
15 spawning velocities for spring-run Chinook Salmon in Butte Creek range from
16 0.80 to 3.22 ft/s (24.4 to 98 cm/s), and suitable substrate size ranges from 1 to
17 5 inches (2.5 to 12.7 cm) in diameter. Adult Chinook have been observed
18 spawning in water greater than 0.8 foot deep and in water velocities of 1.2 to
19 3.5 ft/s (DFG 1998).

20 The timing of spring run spawning in the mainstem Sacramento River has shifted
21 later in the year, which is believed to be a result of genetic introgression with the
22 fall run (Association of California Water Agencies and California Urban Water
23 Agencies 1997). Populations in Deer and Mill creeks, which do not appear to
24 have significantly hybridized with the fall run, generally spawn earlier than those
25 in the main stem (Lindley et al. 2004). Rutter (1908) noted that most spawning in
26 the late 1800s/early 1900s in the Sacramento River basin occurred in August.
27 Parker and Hanson (1944) observed intensive spawning of spring-run Chinook
28 Salmon from the first week of September through the end of October in 1941.
29 Redd counts have indicated that spring-run Chinook Salmon spawning typically
30 begins in late August, peaks in September, and concludes in October in both Deer
31 and Mill creeks (Harvey 1995, Moyle et al. 1995, NMFS 2004a).

32 In the Feather River, the time of river entry for spring-run Chinook Salmon has
33 apparently shifted to later in the season, and is now intermediate between timing
34 of entry of spring run into other tributaries and timing of entry of the fall run.
35 Whereas wild-type spring-run Chinook Salmon enter Deer and Mill creeks
36 primarily in mid-April to mid-June, coded-wire tag data and anecdotal
37 information from anglers indicate that Feather River fish do not enter fresh water
38 until June or July (Association of California Water Agencies and California
39 Urban Water Agencies 1997).

40 **9B.5.4.3.2 Egg Incubation and Alevin Development**

41 In the Sacramento River and its tributaries, egg incubation for spring-run Chinook
42 Salmon extends from August to March (Fisher 1994, Ward and McReynolds
43 2001). Egg incubation generally lasts between 40 and 90 days at water
44 temperatures of 42.8 to 53.6°F (6 to 12°C) (Vernier 1969, Bams 1970, Heming

1 1982). At temperatures of 37°F (2.7°C), time to 50 percent hatching can take up
2 to 159 days (Alderdice and Velsen 1978). Alevins remain in the gravel for 2 to
3 3 weeks after hatching while absorbing their yolk sacs. Emergence from the
4 gravels occurs from November to March in the Sacramento River basin (Fisher
5 1994, Ward and McReynolds 2001). Once fry emerge from the gravel, they
6 initially seek areas of shallow water and low velocities while they finish
7 absorbing the yolk sac (Moyle 2002). As juvenile Chinook Salmon grow, they
8 move into deeper water with higher current velocities, but still seek shelter and
9 velocity refugia to minimize energy expenditures (Healey 1991). USFWS catches
10 of juvenile salmon in the Sacramento River near West Sacramento showed that
11 larger juvenile salmon were captured in the main channel and smaller fry were
12 typically captured along the channel margins (USFWS 1997).

13 **9B.5.4.3.3 Juvenile Rearing and Outmigration**

14 Fry and juvenile rearing takes place in the natal streams, the mainstem of the
15 Sacramento River, inundated floodplains (including the Sutter and Yolo
16 bypasses), and the Delta. During the winter, some spring-run juveniles have been
17 found rearing in the lower portions of non-natal tributaries and intermittent
18 streams (Maslin et al. 1997, Snider et al. 2001).

19 The rearing and outmigration patterns exhibited by spring-run Chinook Salmon
20 are highly variable, with fish rearing anywhere from 3 to 15 months before
21 outmigrating to the ocean (Fisher 1994). Variation in length of juvenile residence
22 may be observed both within and among streams (e.g., Butte versus Mill creeks,
23 [USFWS 1996]). Some may disperse downstream soon after emergence as fry in
24 March and April, with others smolting after several months of rearing, and still
25 others remaining to oversummer and emigrate as yearlings (USFWS 1996). Scale
26 analysis indicates that most returning adults have emigrated as subyearlings
27 (Myers et al. 1998). Calkins et al. (1940) conducted an analysis of scales of
28 returning adults, and estimated that more than 90 percent had emigrated as
29 subyearlings, at about 3.5 inches (88 mm).

30 The term “yearling” is generally applied to any juveniles that remain to
31 oversummer in their natal stream. Yearling outmigrants are common in Deer and
32 Mill creeks, but rare in Butte Creek (Association of California Water Agencies
33 and California Urban Water Agencies 1997). Extensive outmigrant trapping in
34 Butte Creek has shown that spring-run Chinook Salmon outmigrate primarily as
35 juvenile (age 0+) fish from November through June, with a small proportion
36 remaining to emigrate as yearlings beginning in mid-September and extending
37 through March, with a peak in November (Association of California Water
38 Agencies and California Urban Water Agencies 1997, Hill and Webber 1999,
39 Ward et al. 2004). Peak movement of juvenile spring-run Chinook Salmon in the
40 Sacramento River at Knights Landing generally occurs in December, and again in
41 March. However, juveniles also have been observed migrating between
42 November and the end of May (Snider and Titus 1998, 2000b, c, d; Vincik et al.
43 2006; Roberts 2007).

1 Coded-wire-tag studies conducted on Butte Creek spring-run Chinook Salmon
2 have shown that juveniles use the Sutter Bypass as a rearing area until it begins to
3 drain in the late winter or spring (Hill and Webber 1999). Few juvenile Chinook
4 Salmon are observed in the bypass after mid-May. Five recaptures indicate that
5 juveniles leaving the Sutter Bypass migrate downstream rapidly and do not use
6 the mainstem Sacramento River as rearing habitat (Hill and Webber 1999).

7 Within the Delta, juvenile Chinook Salmon forage in shallow areas with
8 protective cover, such as tidally influenced sandy beaches and shallow water areas
9 with emergent aquatic vegetation (Meyer 1979, Healey 1980). Very little
10 information is available on the estuarine rearing of spring-run Chinook Salmon
11 (NMFS 2004a). NMFS (2004a) postulates that, because spring-run Chinook
12 Salmon yearling outmigrants are larger than fall-run Chinook Salmon smolts, and
13 are ready to smolt upon entering the Delta, they may spend little time rearing in
14 the estuary. Most have presumably left the estuary by mid-May (DFG 1995).
15 Once in the ocean, spring-run Chinook Salmon perform extensive offshore
16 migrations before returning to their natal streams to spawn.

17 **9B.5.4.4 Population Trends**

18 At one time, spring-run Chinook Salmon may have been the most abundant race
19 in the Central Valley, with escapement in the hundreds of thousands (Mills and
20 Fisher 1994). Spring-run Chinook Salmon have since declined to remnant
21 populations totaling a few thousand fish, sometimes approaching 30,000 to
22 40,000 in good years (Mills and Fisher 1994, NMFS 1999a). Loss of access to
23 upstream spawning and rearing areas due to the construction of dams in the
24 Sacramento and San Joaquin rivers is believed to have been a major cause of the
25 decline of the spring run.

26 Under historical conditions, it is doubtful that spring-run Chinook Salmon
27 spawned in the mainstem Sacramento in significant numbers (Lindley et al.
28 2004). After the closure of Shasta and Keswick dams, spring-run Chinook
29 Salmon began to spawn in the mainstem Sacramento River when changes in
30 temperatures made this a viable life-history strategy. Throughout the 1970s and
31 1980s, thousands of spring-run Chinook Salmon passed RBDD en route to
32 spawning grounds farther upstream. By the 1990s, escapements had declined;
33 however, changes in the RBDD gate operations beginning in 1986 complicated
34 the process of estimating spring-run Chinook Salmon abundance. Identification
35 of the spring run at RBDD is also complicated by their low escapements and the
36 difficulty of distinguishing fish of this run from those of the fall run. The two
37 runs cannot be distinguished reliably by physical characteristics or run timing
38 (Healey 1991) because of the naturally protracted run timing of the abundant fall
39 run, and the apparent shift to later upstream migration timing by the spring run,
40 which results in the runs being more temporally overlapped than they were
41 historically.

42 Populations of spring-run Chinook Salmon in Butte Creek increased after the
43 1990s, and Butte Creek currently has the largest naturally spawning spring-run
44 population (DFW 2014, GrandTab data). A few naturally spawning fish are also

1 present in Battle, Clear, Cottonwood, Antelope, Mill, Deer, and Big Chico creeks
2 (DFW 2014, GrandTab data). In general, spring-run Chinook Salmon that are
3 most genetically similar to the runs that occurred historically in the Sacramento
4 basin are currently confined to spawning primarily in Deer, Mill, and Butte
5 creeks, with perhaps a few spawning in the mainstem Sacramento River.

6 Restrictions on ocean harvest to protect winter-run Chinook Salmon, as well as
7 improved ocean conditions, have likely had a positive impact on spring-run
8 Chinook Salmon adult returns to the Central Valley. In 2008, abundance in key
9 indicator streams (e.g., Mill, Deer, and Butte Creeks) was at historical levels;
10 however, between 2008 and 2011, spring-run populations in these same streams
11 dropped closer to historical lows (as based on preliminary DFW 2014, GrandTab
12 data). Spring-run Chinook Salmon populations generally increased from 1990
13 through 2006, but then returned to very low levels by 2008 and remained low
14 through 2011. The preliminary total spring-run Chinook Salmon escapement
15 count for 2013 was 23,697 adults, which was the highest count since 2003
16 (30,697 adults) and over three times that of 2011 (7,408 adults) (DFW 2014)
17 (Table 9B.5).

1 **Table 9B.5 Recent Spring-run Chinook Salmon Natural and Hatchery Escapement**

YEAR	Sacramento River Mainstem	Battle Ck^a	Clear Ck	Cottonwood Ck	Antelope Ck	Mill Ck	Deer Ck	Big Chico Ck	Butte Ck Snorkel	Butte Ck Carcass	Feather River Hatchery^b	TOTAL SPRING RUN
1990	4,198	2	–	–	–	844	496	–	250	–	1,893	7,683
1991	825	–	–	–	–	319	479	–	–	–	4,303	5,926
1992	371	–	–	–	0	237	209	–	730	–	1,497	3,044
1993	391	–	1	1	3	61	259	38	650	–	4,672	6,076
1994	862	–	0	–	0	723	485	2	474	–	3,641	6,187
1995	426	66	2	8	7	320	1,295	200	7,500	–	5,414	15,238
1996	378	35	–	6	1	253	614	2	1,413	–	6,381	9,083
1997	128	107	–	0	0	202	466	2	635	–	3,653	5,193
1998	1,115	178	47	477	154	424	1,879	369	20,259	–	6,746	31,649
1999	262	73	35	102	40	560	1,591	27	3,679	–	3,731	10,100
2000	43	78	9	122	9	544	637	27	4,118	–	3,657	9,244
2001	621	111	0	245	8	1,104	1,622	39	9,605	18,670	4,135	26,663
2002	195	222	66	125	46	1,594	2,195	0	8,785	16,409	4,189	25,043
2003	0	221	25	73	46	1,426	2,759	81	4,398	17,404	8,662	30,697
2004	370	90	98	17	3	998	804	0	7,390	10,558	4,212	17,150
2005	30	73	69	47	82	1,150	2,239	37	10,625	17,592	1,774	23,093
2006	0	221	77	55	102	1,002	2,432	299	4,579	6,537	2,181	12,906
2007	248	291	194	34	26	920	644	0	4,943	6,871	2,635	11,144

Appendix 9B: Aquatic Species Life History Accounts

YEAR	Sacramento River Mainstem	Battle Ck^a	Clear Ck	Cottonwood Ck	Antelope Ck	Mill Ck	Deer Ck	Big Chico Ck	Butte Ck Snorkel	Butte Ck Carcass	Feather River Hatchery^b	TOTAL SPRING RUN
2008	52	105	200	0	3	381	140	0	3,935	11,046	1,460	13,387
[2009]	0	194	120	0	0	220	213	6	2,059	2,763	989	4,505
[2010]	0	172	21	15	17	482	262	2	1,160	1,991	1,661	4,623
[2011]	0	157	8	2	6	366	271	124	2,130	4,505	1,969	7,408
[2012]	0	799	68	1	1	768	734	0	8,615	16,140	3,738	22,249
[2013]	0	608	659	1	0	644	708	0	11,470	16,783	4,294	23,697
[2014]	0	429	95	2	7	679	830	0	3,616	5,083	2,776	9,901

1 Source: DFW 2014, GrandTab data.

2 Notes:

3 Data for years in brackets are preliminary.

4 a. In 2009, USFWS conducted a comprehensive analysis of Battle Creek coded wire tag data from 2000-2008 to estimate numbers of fall- and late
5 fall-run Chinook Salmon returning to Battle Creek. Previously, a cutoff date of December 1 was used to assign run. This changed some Battle
6 Creek estimates.

7 b. Feather River Hatchery implemented a methodology change in 2005 for distinguishing spring- from fall-run. Fish arriving prior to the spring-run
8 spawning period were tagged and returned to the river. The spring-run escapement was the number of these tagged fish that subsequently
9 returned to the hatchery during the spring-run spawning period.

1 **9B.5.5 Central Valley Fall-run and Late Fall-run Chinook Salmon**

2 **9B.5.5.1 Legal Status**

3 Federal: Species of Concern

4 State: Central Valley fall-run – None; Central Valley late fall-run – Species of
5 Special Concern

6 Fall-run populations occur throughout the range of Chinook Salmon and are
7 currently the most abundant and widespread of the salmon runs in California and
8 the Central Valley, largely because the construction of dams was not as damaging
9 in terms of loss of historical habitat compared to the runs that spawned at higher
10 elevations. Fall-run abundance is also a function of hatchery supplementation,
11 because fall-run Chinook Salmon have been the primary focus of hatchery
12 production at Central Valley hatcheries for several decades. As the most
13 abundant salmonid species in the Central Valley, fall-run Chinook Salmon
14 constitute an important component of the commercial and recreational salmon
15 fishery in California. NMFS designated the Central Valley Fall (and Late fall)
16 Chinook Salmon ESU as a Species of Concern in 2004 (NMFS 2004b).

17 NMFS classifies late fall-run Chinook Salmon as part of the Central Valley fall-
18 run and late fall-run Chinook Salmon ESU, reasoning that the late fall-run
19 population represents a life-history variation of the fall-run salmon population
20 rather than a distinct run (NMFS 2004b). However, agencies generally treat late
21 fall-run salmon in the Sacramento River basin as a distinct run, conducting
22 separate carcass and redd surveys for them, and publishing separate reports to
23 address the fall-run and late fall-run populations. Agencies also manage the
24 hatchery propagation of late fall-run separately from fall-run Chinook Salmon.
25 Except for hatchery propagation, there are relatively few restoration and
26 management activities that focus specifically on late fall-run Chinook Salmon in
27 the Sacramento River, as compared to the other runs of Chinook Salmon in the
28 basin (USFWS 1996).

29 **9B.5.5.2 Distribution**

30 **9B.5.5.2.1 Fall-run Chinook Salmon**

31 Within the range of the Central Valley ESU, large populations of fall-run Chinook
32 Salmon are found in the Sacramento River and its major tributaries. Fall-run
33 Chinook Salmon are the most widely distributed salmonid in the Sacramento
34 River basin, with significant spawning populations documented as far north as the
35 upstream limit of anadromy in the upper Sacramento River (Keswick Dam at
36 RM 302) and as far south as the American River near Sacramento. Sizeable
37 spawning populations occur in other tributaries to the Sacramento River—Clear
38 Creek, Battle Creek, Butte Creek, and Feather River—with more modest
39 spawning populations in numerous smaller tributaries (e.g., Deer, Mill, Cow, and
40 Antelope creeks). The San Joaquin River system once supported large runs of
41 both spring-run and fall-run Chinook Salmon. Fall-run Chinook Salmon
42 historically spawned in the mainstem San Joaquin River upstream of the Merced

1 River confluence and in the mainstem channels of the major tributaries—the
2 Merced, Tuolumne, and Stanislaus rivers. Dam construction and water diversion
3 dewatered much of the mainstem San Joaquin River, limiting fall-run Chinook to
4 the three major tributaries where they currently spawn and rear downstream of
5 mainstem dams.

6 **9B.5.5.2.2 Late Fall-run Chinook Salmon**

7 Little is known about the historical distribution of late fall-run salmon in the
8 Sacramento River valley. Late fall-run Chinook Salmon currently spawn
9 primarily in the mainstem Sacramento River between Red Bluff (RM 243.5) and
10 Keswick Dam (RM 302). DFW conducts aerial redd surveys that target the late
11 fall-run spawning period, and an analysis of the surveys suggests that adults
12 generally spawn upstream of RBDD (RM 243.5). Yoshiyama et al. (1996)
13 gleaned incidental references to late fall-run fish from historical documents to
14 suggest that late fall-run Chinook Salmon historically spawned in the mainstem
15 reaches of the upper Sacramento River and tributaries such as the Little
16 Sacramento, Pit, and McCloud rivers. Because a significant fraction of juvenile
17 late fall-run Chinook Salmon overwinter in natal streams before emigrating,
18 mainstem reaches close to coldwater sources were likely the most important
19 historical spawning areas for late fall-run Chinook Salmon. Unfortunately, there
20 is little historical data on water temperatures in the upper Sacramento River basin
21 to analyze the stream reaches that may have been important spawning and rearing
22 areas for the late fall-run. Yoshiyama et al. (1996) also suggested the presence of
23 historical spawning populations of late fall-run Chinook Salmon in the American
24 and San Joaquin rivers prior to the era of large dam construction.

25 **9B.5.5.3 Life History and Habitat Requirements**

26 General habitat requirements for Chinook Salmon were described previously.
27 Only habitat requirements specific to fall-run and late fall-run Chinook Salmon
28 are described here.

29 Historically, the summer water temperature regime in the Sacramento River was a
30 key variable that influenced the life history timing and strategy of the different
31 salmonids that occur in the basin. Fall-run Chinook Salmon avoid stressful
32 summer conditions by migrating upstream in the fall (September–November)
33 when both air and water temperatures begin to cool. Because they arrive at
34 spawning grounds with fully developed gonads, adult fall-run can spawn
35 immediately (October–November), which allows their progeny to emerge in time
36 to emigrate from the Sacramento River as fry in the subsequent spring (February–
37 May) before water temperatures become too high.

38 Because fall-run Chinook Salmon adults migrate upstream during periods of low
39 fall baseflows, spawning is generally limited to the alluvial reaches of mainstem
40 rivers below flow-related obstacles. There is relatively little overwintering
41 habitat in these lower mainstem reaches to support a yearling life history strategy,
42 so the majority of fall-run juveniles emigrate as fry before spring water
43 temperatures become lethal. Historically, warming spring water temperatures

1 may have imposed a lethal penalty on the progeny of any late-arriving fall-run
2 adults.

3 Yoshiyama et al. (1996) suggested that spawning populations of late fall-run
4 salmon occurred in the Sacramento River prior to the construction of Shasta Dam,
5 citing what are mostly incidental references to late fall-run salmon in several
6 historical documents. Although these historical accounts indicate the occurrence
7 of salmon migrating upstream and spawning in December or later on several
8 different Central Valley tributaries, it is not clear whether such migration and
9 spawning activity occurred consistently or in substantial numbers. These
10 historical references to late fall-run fish may document fall-run stragglers whose
11 progeny perished the subsequent spring and contributed little to the population, or
12 they may indicate passage barriers that delayed the upstream migration and
13 spawning of fall-run fish en masse.

14 Late fall-run salmon in the Sacramento River have been a collateral beneficiary of
15 the operation of the Shasta and Trinity divisions of the CVP, which maintain
16 suitable water conditions for endangered winter-run Chinook Salmon. Since
17 1994, coldwater releases designed to protect winter-run eggs incubating through
18 the summer months have likely expanded suitable oversummering habitat for late
19 fall-run juveniles downstream. Fall-run juveniles could continue to emigrate as
20 fry or spend a summer growing in the river before emigrating as subyearlings.

21 The late fall-run Chinook Salmon strategy is successful because a substantial
22 fraction of juveniles oversummer in the Sacramento River before emigrating,
23 which allows them to avoid predation through both their larger size and greater
24 swimming ability (larger juvenile salmon can evade a certain amount of predation
25 through size alone). One implication of this life history strategy is that rearing
26 habitat is most likely the limiting factor for late fall-run Chinook Salmon,
27 especially if availability of cool water determines the downstream extent of
28 spawning habitat for late fall-run salmon.

29 Tables 9B.6 and 9B.7 display the life-history timing of fall-run and late fall-run
30 Chinook Salmon in the action area.

Appendix 9B: Aquatic Species Life History Accounts

1 **Table 9B.6 Life History Timing of Central Valley Fall-run Chinook Salmon**


Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Adult migration past Red Bluff Diversion Dam												
Spawning												
Incubation												
Fry emergence ^a												
Rearing in mainstem Sacramento River ^b												
Outmigration past Red Bluff Diversion Dam												
Presence at CVP/SWP salvage facilities												
Emigration toward and through the Delta ^c												

2 Notes:

3 a. Northern California Water Association and Sacramento Valley Water Users (2011) shows emergence ending in February; Williams (2006)
 4 shows emergence ending in April.

5 b. A few fall-run Chinook Salmon remain upstream of RBDD location to rear to a yearling life stage.


6 c. NMFS (2012, unpublished data)



1 **Table 9B.7 Life History Timing of Central Valley Late Fall-run Chinook Salmon**

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Adult entry into mainstem Sacramento River ^{a, b}	Peak											
Migration past Red Bluff Diversion Dam ^{a, b, c}												
Adult holding ^d												
Spawning ^{a, b, c, e, f, g}												
Incubation												
Fry emergence ^{a, c}				Peak	Peak							
Stream residency ^{a, c}												
Fry outmigration past Red Bluff Diversion Dam ^b				Peak	Peak							
Smolt outmigration past Red Bluff Diversion Dam ^b										Peak		
Presence at CVP/SWP salvage facilities	Peak	Peak										
Emigration toward and through the Delta ^c	Peak	Peak										
Smolt outmigration ^a												
Ocean entry ^c												

- 2 Sources:
- 3 a. Yoshiyama et al. 1998
- 4 b. Association of California Water Agencies and California Urban Water Agencies
- 5 c. Fisher 1994
- 6 d. Moyle 2002
- 7 e. Snider et al. 1998, 1999, 2000
- 8 f. Northern California Water Association and Sacramento Valley Water Users 2011
- 9 g. Williams 2006



10

1 9B.5.5.3.1 Adult Upstream Migration and Spawning

2 Adult fall-run Chinook Salmon migrate into the Sacramento River and its
3 tributaries from June through December in mature condition, with upstream
4 migration peaking in September and October. Fall-run Chinook Salmon in the
5 San Joaquin system typically enter spawning streams from September through
6 November. Adults spawn soon after arriving at their spawning grounds between
7 late September and December, with peak spawning activity in late October and
8 early November.

9 Adult late fall-run Chinook Salmon migrate up the Sacramento River between
10 mid-October and mid-April, with peak migration occurring in December
11 (Reclamation 1991) (Table 9B.7). Adults spawn soon after reaching spawning
12 areas between January and April. Fisher reports that peak spawning in the
13 Sacramento River occurs in early February (1994), but carcass surveys conducted
14 in the late 1990s suggest that peak spawning may occur in January (Snider et al
15 1998, 1999, 2000).

16 Fall-run and late fall-run Chinook Salmon are generally able to spawn in deeper
17 water with higher velocities than Chinook Salmon in other runs because of their
18 larger size (Healey 1991). Late fall-run salmon tend to be the largest individuals
19 of the Chinook Salmon species that occur in the Sacramento River basin (USFWS
20 1996).

21 Fry emergence occurs from December through March, and fry rear in freshwater
22 for only a few months before migrating downstream to the ocean as smolts
23 between March and July (Yoshiyama et al. 1998). Late fall-run fry emerge from
24 redds between April and June (Vogel and Marine 1991).

25 9B.5.5.3.2 Juvenile Rearing and Outmigration

26 Fall-run Chinook Salmon in the Sacramento River generally exhibit two rearing
27 strategies: migrating to the lower reaches of the river or Delta as fry, or remaining
28 to rear in the gravel-bedded reach for about 3 months and then smolting and
29 outmigrating. The highest abundances of fry in the Delta are observed in wet
30 years (Brandes and McLain 2001). Fall-run Chinook Salmon fry rear during a
31 time and in a location where floodplain inundation is most likely to occur, thereby
32 expanding the amount of rearing habitat available. Relative survival of fry appears
33 to be higher in the upper Sacramento River than in the Delta or bay, especially in
34 wet years (Brandes and McClain 2001).

35 One potential disadvantage of early emergence and emigration and rearing in
36 mainstem channels and the estuary is the possibility of higher predation mortality
37 because of the relatively small size of emigrants. However, fall-run Chinook
38 Salmon fry exhibit several characteristics to combat predation mortality.
39 Predators often occupy deep pools in mainstem channels, so fry generally use
40 shallow water habitat found along channel margins or in runs and riffles to avoid
41 predators. Because rearing habitat is not limiting for fall-run Chinook Salmon
42 fry, they do not exhibit territorial behavior, which allows them to rear, smolt, and
43 outmigrate in higher densities. By emigrating synchronously in schools rather

1 than as individuals, fall-run Chinook Salmon fry and smolts can swamp potential
2 predators to avoid significant losses to predation; and by emigrating in late spring,
3 they have the advantage of higher discharge fueled by early snowmelt, which can
4 reduce their exposure to predation.

5 Fall-run Chinook Salmon juvenile smolt during early spring, prior to increases in
6 water temperatures. Juvenile Chinook Salmon feed and grow as they move
7 downstream in spring and summer; larger individuals are more likely to move
8 downstream earlier than smaller juveniles (Nicholas and Hankin 1989, Beckman
9 et al. 1998), and it appears that in some systems juveniles that do not reach a
10 critical size threshold will not outmigrate, but will remain to oversummer
11 (Bradford et al. 2001). Bell (1958) suggests that the timing of yearling smolt
12 outmigration corresponds to increasing spring discharges and temperatures.
13 Kjelson et al. (1981) observed that peak seine catches of Chinook Salmon fry in
14 the Sacramento-San Joaquin Delta correlated with increases in flow associated
15 with storm runoff. Flow accounted for approximately 30 percent of the variability
16 in the fry catch.

17 As fall-run Chinook Salmon fry and parr migrate downstream, they also use the
18 lower reaches of non-natal tributaries as rearing habitat (Maslin et al. 1997).
19 During periods of high winter and spring runoff, fall-run Chinook Salmon
20 juveniles are also diverted into the bypasses that border the Sacramento River,
21 where growing conditions are generally better than mainstem rearing habitats,
22 which can facilitate higher rates of juvenile survival (Sommer et al. 2001).
23 Natural floodplain or riparian areas that become inundated during high flows may
24 also provide good habitat for juvenile Chinook Salmon and prevent them from
25 being displaced downstream (The Nature Conservancy 2003).

26 Research conducted in the Central Valley suggests that seasonally inundated,
27 shallow water habitats may provide superior rearing habitat for juvenile salmonids
28 than mainstem channels (Sommer et al. 2001). Juvenile fall-run salmon migrate
29 downstream between January and June when floodplains and bypasses are
30 periodically flooded during wet water years. By promoting faster growth,
31 prolonged floodplain inundation likely helps the fall-run population by increasing
32 juvenile salmon survival.

33 As described above, the timing of late fall-run spawning in January through
34 March means that fry emerge between April and June. Water temperatures in the
35 lower Sacramento River are often too high in May and June to support fry
36 survival, so later-emerging fry that migrate downstream likely suffer high rates of
37 mortality and contribute little to the population. This suggests that a significant
38 fraction of late fall-run juveniles rear in the upper Sacramento River throughout
39 the summer before emigrating in the following fall and early winter as large
40 subyearlings (Fisher 1994). Summer rearing is made possible by the cold water
41 releases from the Shasta-Trinity divisions of the CVP. Late fall-run juveniles
42 generally leave the Sacramento River by December (Vogel and Marine 1991),
43 with peak emigration of smolts in October.

1 Although growth rates of juvenile Chinook Salmon may be high at temperatures
2 approaching 19°C (66°F), cooler temperatures may be required to successfully
3 complete the physiological transformation from parr to smolt. Smoltification in
4 juvenile Sacramento River fall-run Chinook Salmon was studied by Marine
5 (1997), who found that juveniles reared under a high temperature regime of 21 to
6 24°C (70 to 75°F) exhibited altered and impaired smoltification patterns relative
7 to those reared at low 55 to 61°F (13 to 16°C) and moderate 17 to 20°C (63 to
8 68°F) temperatures. Some alteration and impairment of smoltification was also
9 seen in the juveniles reared at the moderate temperatures.

10 Chronic exposure to high temperatures may also result in greater vulnerability to
11 predation. In this same study by Marine (1997), Sacramento River fall-run
12 Chinook Salmon reared at the highest temperatures (21 to 24°C [70 to 75°F]) were
13 preyed upon by Striped Bass more often than those reared at low or moderate
14 temperatures. Consumption rates of piscivorous fish such as Sacramento
15 pikeminnow, Striped Bass, and largemouth bass increase with temperature, which
16 may compound the effects of high temperature on juvenile and smolt predation
17 mortality. Juvenile growth rates are an important influence on survival; faster
18 growth thus both increases the range of food items available to them and decreases
19 their vulnerability to predation (Myrick and Cech 2004).

20 **9B.5.5.3.3 Ocean Residence**

21 When fall-run Chinook Salmon produced from the Sacramento-San Joaquin
22 system enter the ocean, they appear to head north to inhabit the northern
23 California-southern Oregon coast (Oregon Department of Fish and Wildlife
24 1987). They typically have a greater tendency to remain along the continental
25 shelf than do stream-type Chinook Salmon (Healey 1983). The age of returning
26 Chinook Salmon adults in California ranges from 2 to 5 years.

27 **9B.5.5.4 Population Trends**

28 Although NMFS considers fall-run and late fall-run Chinook Salmon as part of
29 the same ESU in the Central Valley, most resource agencies have tracked the two
30 runs separately. For example, DFW has conducted aerial redd surveys
31 specifically targeting late fall-run salmon, and the Anadromous Fish Restoration
32 Program (AFRP) has tracked late fall-run salmon escapements as a separate
33 population. However, reports on fall-run escapement estimates vary because
34 some include late fall-run in the estimates, while others do not. Because the older
35 reports often fail to clarify which runs are being enumerated in the escapement
36 estimate, care must be exercised when using fall-run escapement estimates,
37 especially from different sources.

38 **9B.5.5.4.1 Fall-run Chinook Salmon**

39 Fall-run Chinook Salmon estimates are available from 1940; however, systematic
40 counts of Chinook Salmon in the San Joaquin Basin began in 1953, long after
41 construction of large dams on the major San Joaquin basin rivers. Comparable
42 estimates of population size before 1940 are not available. Since population
43 estimates began, the number of fall-run Chinook returning to the San Joaquin

1 Basin annually has fluctuated widely. Escapement in the Tuolumne River
2 dropped from a high of 40,300 in 1985 to a low of about 100 resulting from the
3 1987 to 1992 dry period (TID/MID 1997). With increased precipitation and
4 improved flow conditions, escapement increased to 3,300 in 1996 (TID/MID
5 1997). From 1971 to 2007, hatchery production is estimated to have composed
6 about 29 percent of the returning adult fall-run Chinook Salmon in the
7 San Joaquin basin (PFMC 2008). Table 9B.8 provides a summary of estimated
8 escapement from 1990 to 2013 in the Sacramento and San Joaquin River systems.

Appendix 9B: Aquatic Species Life History Accounts

1 **Table 9B.8 Recent Fall-run Chinook Salmon Natural and Hatchery Escapement**

Year	Sacramento River System				San Joaquin River System			Sacramento and San Joaquin Combined		
	Hatch.	Main.	Trib.	Total	Hatch.	Trib.	Total	Hatch.	In-River	Total
1990	25,611	48,284	12,803	86,698	114	1,041	1,155	25,725	62,128	87,853
1991	28,528	30,631	72,296	131,455	83	917	1,000	28,611	103,844	132,455
1992	30,171	32,229	44,995	107,395	1,078	1,940	3,018	31,249	79,164	110,413
1993	30,234	46,231	82,975	159,440	2,573	3,410	5,983	32,807	132,616	165,423
1994	42,760	58,546	111,078	212,384	2,862	5,421	8,283	45,622	175,045	220,667
1995	45,324	63,934	211,025	320,283	3,925	5,960	9,885	49,249	280,919	330,168
1996	36,936	84,086	213,646	334,668	5,024	11,859	16,883	41,960	309,591	351,551
1997	71,448	119,296	185,484	376,228	7,440	19,129	26,569	78,888	323,909	402,797
1998	75,028	6,318	141,079	222,425	3,890	19,711	23,601	78,918	167,108	246,026
1999	49,657	161,192	180,501	391,350	4,787	18,122	22,909	54,444	359,815	414,259
2000	50,965	96,688	290,698	438,351	7,396	39,934	47,330	58,361	427,320	485,681
2001	61,318	75,296	453,323	589,937	7,391	27,303	34,694	68,709	555,922	624,631
2002	96,248	65,690	672,962	834,900	9,753	28,016	37,769	106,001	766,668	872,669
2003	118,097	89,229	362,161	569,487	8,666	12,839	21,505	126,763	464,229	590,992
2004	116,869	43,604	202,904	363,377	11,406	12,065	23,471	128,275	258,573	386,848
2005	187,427	57,012	172,457	416,896	5,984	14,813	20,797	193,411	244,282	437,693
2006	80,594	55,468	146,427	282,489	4,289	6,176	10,465	84,883	208,071	292,954
2007	22,511	17,061	54,767	94,339	1,130	1,699	2,829	23,641	73,527	97,168
2008	18,785	24,743	25,618	69,146	315	1,830	2,145	19,100	52,191	71,291
[2009]	20,904	5,827	22,842	49,573	1,799	1,757	3,556	22,703	30,426	53,129

Appendix 9B: Aquatic Species Life History Accounts

Year	Sacramento River System				San Joaquin River System			Sacramento and San Joaquin Combined		
	Hatch.	Main.	Trib.	Total	Hatch.	Trib.	Total	Hatch.	In-River	Total
[2010]	46,306	16,372	90,154	152,832	5,421	4,937	10,358	51,727	111,463	163,190
[2011]	87,679	11,957	105,460	205,096	16,293	6,500	22,793	103,972	123,917	227,889
[2012]	136,710	28,701	155,450	320,861	7,620	13,342	20,962	144,330	197,493	341,823
[2013]	107,001	40,084	279,871	426,956	6,279	14,668	20,947	113,280	334,623	447,903
[2014]	50,713	34,876	152,587	238,176	9,627	8,094	17,721	60,340	195,557	255,897

- 1 Source: DFW 2014
- 2 Note:
- 3 Data for years in brackets are preliminary.

1 **9B.5.5.4.2 Late Fall-run Chinook Salmon**

2 There is little information to evaluate the historical abundance of late fall-run
3 salmon in the Sacramento River basin. In fact, late fall-run salmon were first
4 recognized by fishery agencies as a distinct run only after the construction of
5 RBDD in 1966, which permitted more accurate counting of upstream migrants
6 and the timing of upstream migration (USFWS 1996). Between 1967 and 1976,
7 late fall-run salmon escapements averaged 22,000 adults (USFWS 1996);
8 however, between 1977 and 1985, escapements averaged only about 9,900 adults
9 (DFW 2014). Population estimates of late fall-run salmon after 1985 are
10 complicated by changes in RBDD gate operations, when Reclamation began
11 raising the dam gates during winter months to facilitate the upstream migration of
12 winter-run Chinook Salmon. Because the upstream migration of late fall-run
13 salmon overlaps with that of winter-run Chinook Salmon, late fall-run benefited
14 from improved upstream access, but the accuracy of escapement estimates
15 suffered (USFWS 1996). RBDD gate operations were revised again in 1994 so
16 that gates were raised between September 15 and May 15, encompassing the
17 entire upstream migration period of late fall-run salmon and further compromising
18 the calculation of escapements. Post-1985 escapement estimates are cruder
19 because of the change in RBDD gate operations. Table 9B.9 provides a summary
20 of estimated escapement from 1970 to 2013 in the mainstem Sacramento River,
21 Battle Creek, and Clear Creek.

1 **Table 9B.9 Recent Late Fall-run Chinook Salmon Natural and Hatchery Escapement**

Year	Sacramento River above RBDD	CNFH Transfers	Total above RBDD	Sacramento River below RBDD	Battle Creek	Battle Creek CNFH	Battle Creek Total	Clear Creek	Total
Nov 1990-Apr 1991	6,493	118	6,611	1,491	–	161	161	–	8,263
Nov 1991-Apr 1992	8,958	398	9,356	431	–	344	344	–	10,131
Nov 1992-Apr 1993	339	400	739	–	–	528	528	–	1,267
Nov 1993-Apr 1994	137	154	291	–	–	598	598	–	889
Nov 1994-Apr 1995	–	166	166	–	–	323	323	–	489
Nov 1995-Apr 1996	–	48	48	–	–	1,337	1,337	–	1,385
Nov 1996-Apr 1997	–	–	–	–	–	4,578	4,578	–	4,578
Nov 1997-Apr 1998	38,239	–	38,239	1,101	–	3,079	3,079	–	42,419
Nov 1998-Apr 1999	8,683	–	8,683	–	–	7,075	7,075	–	15,758
Nov 1999-Apr 2000	8,580	–	8,580	122	0	4,181	4,181	–	12,883
Nov 2000-Apr 2001	18,351	–	18,351	925	98	2,439	2,537	–	21,813
Nov 2001-Apr 2002	36,004	–	36,004	0	216	4,186	4,402	–	40,406
Nov 2002-Apr 2003	5,346	38	5,384	148	57	3,183	3,240	110	8,882
Nov 2003-Apr 2004	8,824	60	8,884	0	40	5,166	5,206	60	14,150
Nov 2004-Apr 2005	9,493	79	9,572	1,031	23	5,562	5,585	94	16,282
Nov 2005-Apr 2006	7,678	12	7,690	2,485	50	4,822	4,872	42	15,089
Nov 2006-Apr 2007	13,798	66	13,864	1,477	72	3,361	3,433	69	18,843
Nov 2007-Apr 2008	3,673	0	3,673	291	19	6,334	6,353	55	10,372

Appendix 9B: Aquatic Species Life History Accounts

Year	Sacramento River above RBDD	CNFH Transfers	Total above RBDD	Sacramento River below RBDD	Battle Creek	Battle Creek CNFH	Battle Creek Total	Clear Creek	Total
Nov 2008-Apr 2009	3,271	58	3,329	63	32	6,436	6,468	336	10,196
[Nov 2009-Apr 2010]	3,843	81	3,924	439	27	5,505	5,532	91	9,986
[Nov 2010-Apr 2011]	3,686	39	3,725	0	28	4,635	4,663	58	8,446
[Nov 2011-Apr 2012]	2,811	47	2,858	11	19	3,031	3,050	50	5,969
[Nov 2012-Apr 2013]	4,918	43	4,961	309	42	3,577	3,619	77	8,966
[Nov 2013-Apr 2014]	7,227	39	7,266	723	120	4,869	4,989	72	13,050

1 Source: DFW 2014

2 Note:

3 Data for years in brackets are preliminary.

1 **9B.5.5.4.3 Hybridization**

2 Historically, spring-run Chinook Salmon and fall-run Chinook Salmon both
 3 spawned during the fall, but they were separated spatially because spring-run
 4 Chinook Salmon spawned in upper tributaries that the fall-run Chinook Salmon
 5 could not access. Under current conditions, the Keswick and Shasta dams have
 6 prevented spring-run Chinook Salmon from accessing upper tributaries, and
 7 instead they spawn in the mainstem Sacramento River where the fall run spawns.
 8 The elimination of spatial segregation of fall-run Chinook Salmon and spring-run
 9 Chinook Salmon spawning contributed to hybridization on the spawning grounds
 10 (Yoshiyama et al. 1998). Also, hatchery practices have likely mixed fall-run and
 11 spring-run Chinook Salmon stocks, causing even greater hybridization. By
 12 hybridizing with spring-run Chinook Salmon, the peak spawning activity of fall-
 13 run Chinook Salmon has likely shifted to occur earlier than it did historically.

14 **9B.5.5.5 Hatchery Influence**

15 Fall-run Chinook Salmon have long been a focus of hatchery production in the
 16 Central Valley, and the artificial propagation of the fall run supports the
 17 commercial and recreational harvest of salmon in California. Within the
 18 Sacramento River basin, Coleman National Fish Hatchery on Battle Creek
 19 produces substantial numbers of fall-run salmon for release in the Sacramento
 20 River and Bay-Delta estuary. Using a mixed-stock model to estimate the
 21 contribution of wild fish from the Central Valley to the fall-run Chinook Salmon
 22 ocean fishery, Barnett-Johnson et al. (2007) found that the contribution of wild
 23 fish was about 10 percent, which suggests that hatchery supplementation is a
 24 substantial contributor to the population.

25 Late fall-run salmon have been artificially propagated at the Coleman National
 26 Fish Hatchery on Battle Creek for more than two decades. USFWS releases
 27 between 200,000 and 2.5 million late fall-run juveniles in the Sacramento basin
 28 each year, primarily in Battle Creek. Although hatchery strays likely compose a
 29 portion of the spawning population of late fall-run salmon in the Sacramento
 30 River, it is unclear what proportion of escapements that hatchery-origin fish
 31 constitutes. It is also unclear whether hatchery juveniles that are released in
 32 Battle Creek compete with naturally spawned juveniles for oversummering
 33 habitat in the mainstem Sacramento River.

34 **9B.5.6 Upper Klamath and Trinity Rivers Spring-Run Chinook**
 35 **Salmon**

36 **9B.5.6.1 Legal Status**

37 Federal: Not warranted

38 State: Species of Special Concern

39 Two Chinook Salmon ESUs are found in the Klamath basin, the Southern Oregon
 40 and Coastal (SOCC) ESU and the Upper Klamath and Trinity Rivers ESU. The
 41 former are fall-run fish that spawn in the mainstem of the lower Klamath River.
 42 The Upper Klamath and Trinity Rivers ESU contains fall-run, late fall-run, and

1 spring-run fish that spawn in the Klamath and Trinity rivers upstream of the
2 Trinity River's confluence with the Klamath. Although wild spring-run Chinook
3 Salmon in the Klamath River system differ from fall-run Chinook Salmon
4 genetically, as well as in terms of life history and habitat requirements (NRC
5 2004), all are included within this ESU (Myers et al. 1998). The following profile
6 pertains only to the spring-run, and focuses on the South Fork Trinity River
7 (SFTR), which is within the action area and supports one of the few remaining
8 stocks of wild spring-run Chinook Salmon within the greater Klamath Basin (Van
9 Kirk and Naman 2008). The SFTR is the largest undammed river remaining in
10 California.

11 A status review in 1999 concluded that neither ESU warranted listing (NMFS
12 1999b). A petition to list the Upper Klamath and Trinity Rivers ESU was
13 submitted to NMFS in January 2011 (CBD et al. 2011); in April 2011, NMFS
14 announced that listing was not warranted. Of primary importance in their
15 decision was their conclusion that the spring-run and fall-run Chinook Salmon in
16 the basin constitute a single ESU (NMFS 2012). The genetic structure of
17 Chinook Salmon populations in coastal basins (as opposed to the Central Valley)
18 indicates that the spring- and fall-run life histories have evolved multiple times in
19 different watersheds (Myers et al. 1998, Waples et al. 2004). Three hatchery
20 stocks from the Iron Gate and Trinity River hatcheries are considered part of the
21 ESU because they were founded using native, local stock in the watershed where
22 fish are released (NMFS 2012).

23 **9B.5.6.2 Distribution**

24 The Upper Klamath and Trinity Rivers ESU includes all naturally spawned and
25 hatchery populations of spring, fall, and late-fall runs of Chinook Salmon in the
26 Klamath and Trinity rivers upstream of the confluence of the Klamath and Trinity
27 rivers. Iron Gate Dam currently blocks upstream migration to historical spawning
28 habitat on the Klamath River, and Lewiston Dam is likewise a barrier to upstream
29 migration on the Trinity River.

30 **9B.5.6.3 Life History and Habitat Requirements**

31 General habitat requirements for Chinook Salmon are described earlier; the
32 following describes life-history strategies and habitat requirements unique to the
33 spring-run Chinook or of primary importance to its life history. Spring-run
34 Chinook Salmon display a stream-type life-history strategy—adults migrate
35 upstream while sexually immature, hold in deep cold pools over the summer, and
36 spawn in late summer and early fall. Juvenile outmigration is highly variable,
37 with some age 0+ juveniles outmigrating in their first spring, but others
38 overwintering and then emigrating as yearlings the following spring.

39 Table 9B.10 illustrates life-history timing for spring-run Chinook Salmon in the
40 South Fork Trinity River basin.

1 **Table 9B.10 Life History Timing of Spring-run Chinook Salmon in the South Fork Trinity River**

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Adult upstream migration in Klamath River ^a												
Spawning in SFTR ^b												
Incubation and alevin development												
Fry emergence ^c												
Age 0+ outmigration in SFTR ^{d, e}												
Age 1+ outmigration in SFTR ^{d, f}				?	?	?	?	?				
Ocean entry (yearlings)												

2 Sources:

3 a. Snyder 1931; Strange 2008


4 b. State Coastal Conservancy 2009

5 c. West et al. 1990

6 d. Dean 1994, 1995

7 e. It is not possible to differentiate between fall-run and spring-run juveniles; therefore, exact timing for the spring run is unknown and may differ
8 from the fall run.

9 f. Occurs in the spring after spawning; exact timing unknown.



10

1 **9B.5.6.3.1 Adult Upstream Migration, Holding, and Spawning**

2 Adults spawn from September through early November in the South Fork Trinity
3 River (State Coastal Conservancy 2009).

4 Within the SFTR watershed, spring-run Chinook Salmon spawning takes place
5 primarily between Hitchcock Creek and the East Fork of the SFTR on the
6 mainstem SFTR, in Plummer Creek, in the mainstem of Hayfork Creek and the
7 lower reaches of Salt and Tule creeks (USFS 2001a, Reclamation 1994), and
8 possibly Big Creek (Chilcote et al. 2012). The East Fork of Hayfork Creek is used
9 as summer holding habitat by adults, according to USFS (2001b), and adults have
10 been observed during August in the lower SFTR below Surprise Creek and below
11 Mule Bridge (USFS 2011).

12 **9B.5.6.3.2 Egg Incubation and Alevin Development**

13 Emergence takes place from March until early June (West et al. 1990).

14 **9B.5.6.3.3 Juvenile Rearing and Outmigration**


15 Rearing in the SFTR basin takes place in the mainstem SFTR between Hitchcock
16 Creek and the East Fork of the SFTR (USFS 2001a). This area was noted to be an
17 oversummering area by USFS (2001a). Rearing also takes place in Plummer
18 Creek (USFS 2001a).

19 Juvenile spring-run Chinook Salmon of the Upper Klamath and Trinity Rivers
20 ESU generally remain in fresh water for a year or more. On the South Fork
21 Trinity River, outmigration occurs in late April and May with a peak in May
22 (Dean 1994, 1995); however, it is not possible to differentiate between spring and
23 fall juveniles, so spring-run outmigration timing may differ somewhat from the
24 fall run. Age-1 juveniles (Type III) have been found to outmigrate from the South
25 Fork Trinity River during the following spring (Dean 1994, 1995).

26 **9B.5.6.4 Population Trends**

27 A review by Williams et al. (2011) of Myers et al. (1998) and DFG (1965)
28 estimates historical abundance of the entire ESU (both spring and fall runs) at
29 approximately 130,000 adults for 1912, evenly split between the Klamath and
30 Trinity rivers (NMFS 2012). Since the review by Myers et al. (1998) was
31 published, there apparently has been little change in abundance, population
32 trends, or population growth rates (Williams et al. 2011), except for two of the
33 three spring-run populations that were evaluated, one of which was the South
34 Fork Trinity River, where abundance is low relative to historical estimates
35 (NMFS 2012). The spring run likely dominated numbers of Chinook Salmon in
36 the South Fork Trinity River historically (Reclamation 1994). Declines in the
37 SFTR basin have been attributed to increased sediment delivery and destruction
38 of riparian vegetation from a history of logging and road-building in the
39 characteristically unstable soils found there (USFS 1996; Trinity County
40 Resource Conservation District 2003), effects of the 1964 flood (Reclamation
41 1994), major wildfire events (e.g., 1987, 2008), mining, and livestock grazing
42 (Chilcote et al. 2012), as well as water withdrawals and clearing of large woody

1 debris from stream channels (USFS 1994). Water withdrawals for domestic and
2 agricultural uses appear to be a major factor influencing fish production in
3 Hayfork Creek (Reclamation 1994), a major tributary to the SFTR that is located
4 in more stable soils. Temperatures in the SFTR and Hayfork Creek are believed
5 to be limiting spring-run populations in the SFTR and Hayfork Creek (Chilcote
6 et al. 2012), thus climate change could result in future declines (Van Kirk and
7 Naman 2008). NMFS suspects that dams on the mainstem Klamath and Trinity
8 rivers caused as much as 90 percent of the spring-run Chinook Salmon decline
9 (USFS 2001b). These dams may affect Chinook Salmon populations by altering
10 natural seasonal flow patterns and temperatures, which affects habitat as well as
11 behavioral cues for life-history transitions (USFS 1999). Escapement of spring-
12 run Chinook Salmon to the Trinity River is shown in Figure 9B.1.



1

2 **Figure 9B.1 Spring-run Chinook Salmon Escapement in the Trinity River, 1980–**
 3 **2010 (from Williams et al. 2011)**

1 **9B.5.6.5 Hatchery Influences**

2 Hatchery stocking using native Chinook Salmon began in 1917 and includes both
3 fall- and spring-run fish. There are two hatcheries in the basin: Iron Gate
4 Hatchery on the Klamath River and Trinity River Hatchery on the Trinity River.
5 Chinook Salmon released from Iron Gate Hatchery are all fall-run fish (NRC
6 2004), while the Trinity River Hatchery produces both spring- and fall-run
7 Chinook Salmon. Approximately 10.3 million fingerling and yearling Chinook
8 Salmon are released annually from these two hatcheries (NMFS 2012). The
9 stocks from these hatcheries were founded from local, native fish and are
10 genetically similar to local, natural populations; they are considered part of the
11 same ESU by NMFS (NMFS 2012).

12 **9B.5.7 References**

- 13 AFS (American Fisheries Society). 1985. *Petition to List the Winter-run of*
14 *Chinook Salmon on the Sacramento River of California as a Threatened*
15 *Species*. Submitted by Cay Goude of the California-Nevada Chapter of the
16 American Fisheries Society to Dr. William Gordon, Director, National
17 Marine Fisheries Service as cited by National Marine Fisheries Service in
18 51 FR 5391-5392.. October 31, 1985.
- 19 Alderdice, D. F., and F. P. J. Velsen. 1978. Relation between temperature and
20 incubation time for eggs of Chinook salmon (*Oncorhynchus tshawytscha*).
21 *Journal of the Fisheries Research Board of Canada* 35: 69-75.
- 22 Alderdice, D. F., W. P. Wickett, and J. R. Brett. 1958. Some effects of temporary
23 exposure to low dissolved oxygen levels on Pacific salmon eggs. *Journal*
24 *of the Fisheries Research Board of Canada* 15: 229-250.
- 25 Armour, C. L. 1990. *Guidance for evaluating and recommending temperature*
26 *regimes to protect fish*. Instream Flow Information Paper 28, Biological
27 Report 90 (22). U.S. Fish and Wildlife Service, National Ecology
28 Research Center, Fort Collins, Colorado.
- 29 Association of California Water Agencies and California Urban Water Agencies.
30 1996. *The Status of Late-fall and Spring-run Chinook Salmon in the*
31 *Sacramento River Basin Regarding the Endangered Species Act*. Special
32 Report. Submitted to National Marine Fisheries Service. Prepared by S. P.
33 Cramer and D. B. Demko, S.P. Cramer and Associates, Inc., Gresham,
34 Oregon.
- 35 Atwater, B. F., S. G. Conard, J. N. Dowden, C. W. Hedel, R. L MacDonald, and
36 W. Savage. 1979. History, landforms, and vegetation of the estuary's tidal
37 marshes. In *San Francisco Bay: the Urbanized Estuary*, pp. 347-385.
38 Edited by T. J. Conomos. Pacific Division of the American Association
39 for the Advancement of Science, San Francisco, California.
- 40 Azat, J. 2012. Central Valley Chinook salmon harvest and escapement.
41 *Interagency Ecological Program for the San Francisco Estuary* 25:13-15.

- 1 Azevedo, R. L., and Z. E. Parkhurst. 1958. *The upper Sacramento River salmon*
2 *and steelhead maintenance program, 1949-1956*. United States Fish and
3 Wildlife Service. As cited in Slater 1963.
- 4 Bams, R. A. 1970. Evaluation of a revised hatchery method tested on pink and
5 chum salmon fry. *Journal of the Fisheries Research Board of Canada* 27:
6 1429-1452.
- 7 Banks, J. L., L. G. Fowler, and J. W. Elliott. 1971. Effects of rearing temperature
8 on growth, body form, and hematology of fall Chinook fingerlings. *The*
9 *Progressive Fish-Culturist* 33: 20-26.
- 10 Barnett-Johnson, R., C. B. Grimes, C. F. Royer, and C. J. Donohoe. 2007.
11 Identifying the contribution of wild and hatchery Chinook salmon
12 (*Oncorhynchus tshawytscha*) to the ocean fishery using otolith
13 microstructure as natural tags. *Canadian Journal of Fisheries and Aquatic*
14 *Sciences* 64:1683-1692.
- 15 Beckman, B. R., D. A. Larsen, B. Lee-Pawlak, and W. W. Dickhoff. 1998.
16 Relation of fish size and growth rate to migration of spring Chinook
17 salmon smolts. *North American Journal of Fisheries Management* 18:
18 537-546.
- 19 Beechie, T. J., M. Liermann, E. M. Beamer, and R. Henderson. 2005. A
20 classification of habitat types in a large river and their use by juvenile
21 salmonids. *Canadian Journal of Fisheries and Aquatic Sciences* 134:717-
22 729.
- 23 Bell, M. C. 1986. *Fisheries Handbook of Engineering Requirements and*
24 *Biological Criteria*. Report No. NTIS AD/A167-877. Fish Passage
25 Development and Evaluation Program, U.S. Army Corps of Engineers,
26 North Pacific Division, Portland, Oregon.
- 27 Bell, R. 1958. Time, Size, and Estimated Numbers of Seaward Migrants of
28 Chinook Salmon and Steelhead Trout in the Brownlee-Oxbow Section of
29 the Middle Snake River. State of Idaho Department of Fish and Game,
30 Boise.
- 31 Bjornn, T. C. 1971. Trout and salmon movements in two Idaho streams as related
32 to temperature, food, stream flow, cover, and population density.
33 *Transactions of the American Fisheries Society* 100: 423-438.
- 34 Bjornn, T. C., and D. W. Reiser. 1991. Habitat requirements of salmonids in
35 streams. In *Influences of Forest and Rangeland Management on Salmonid*
36 *Fishes and their Habitats*, pp. 83-138. Edited by W. R. Meehan. Special
37 Publication No. 19. American Fisheries Society, Bethesda, Maryland.
- 38 Boles, G. L., S. M. Turek, C. D. Maxwell, and D. M. McGill. 1988. *Water*
39 *Temperature Effects on Chinook Salmon (Oncorhynchus tshawytscha)*
40 *with Emphasis on the Sacramento River: a Literature Review*. California
41 Department of Water Resources, Northern District, Red Bluff.

- 1 Bradford, M. J., J. A. Grout, and S. Moodie. 2001. Ecology of juvenile Chinook
2 salmon in a small non-natal stream of the Yukon River drainage and the
3 role of ice conditions on their distribution and survival. *Canadian Journal*
4 *of Zoology* 79: 2043-2054.
- 5 Brandes, P. L., and J. S. McLain. 2001. Juvenile Chinook salmon abundance,
6 distribution, and survival in the Sacramento-San Joaquin estuary.
7 *Contributions to the Biology of Central Valley Salmonids*, pp. 39-138.
8 Edited by R. L. Brown. Fish Bulletin 179: Volume 2. California
9 Department of Fish and Game, Sacramento.
- 10 Brett, J. R. 1952. Temperature tolerance in young Pacific salmon, genus
11 *Oncorhynchus*. *Journal of the Fisheries Research Board of Canada* 9:
12 265-323.
- 13 Brett, J. R., W. C. Clarke, and J. E. Shelbourn. 1982. Experiments on thermal
14 requirements for growth and food conversion efficiency of juvenile
15 Chinook salmon *Oncorhynchus tshawytscha*. *Canadian Technical Report*
16 *of Fisheries and Aquatic Sciences* 1127. Department of Fisheries and
17 Oceans, Fisheries Research Branch, Pacific Biological Station, Nanaimo,
18 British Columbia.
- 19 Burner, C. J. 1951. Characteristics of spawning nests of Columbia River salmon.
20 *U.S. Fish and Wildlife Service Fishery Bulletin* 52: 97-110.
- 21 Burnett, K. M., and G. H Reeves. 2001. Valley segment use by juvenile ocean-
22 type Chinook salmon (*Oncorhynchus tshawytscha*) in tributaries of the Elk
23 River, Oregon (1988–1994). Chapter 3 in *Relationships among Juvenile*
24 *Anadromous Salmonids, their Freshwater Habitat, and Landscape*
25 *Characteristics over Multiple Years and Spatial Scales in the Elk River,*
26 *Oregon*. Doctoral dissertation. Oregon State University, Corvallis.
- 27 CALFED Bay-Delta Program. n.d. *Ecosystem Restoration: Winter-run Chinook*
28 *Salmon in the Sacramento River*.
29 www.calwater.ca.gov/science/pdf/eco_restor_winter_chinook.pdf.
- 30 Calkins, R. D., W. F. Durand, and W. H. Rich. 1940. *Report of the Board of*
31 *Consultants on the Fish Problem of the Upper Sacramento River*. Stanford
32 University, Stanford, California. As cited in Myer et al. 1998.
- 33 CBD et al. (Center for Biological Diversity, Oregon Wild, Environmental
34 Protection Information Center, and The Larch Company). 2011. *Petition*
35 *to List Upper Klamath Chinook Salmon (Oncorhynchus tshawytscha) as a*
36 *Threatened or Endangered Species*.
- 37 CDFW (California Department of Fish and Wildlife). 2014. GrandTab
38 2014.04.22. California Central Valley Chinook Population Report.
39 Compiled April 22, 2014. Fisheries Branch.

Appendix 9B: Aquatic Species Life History Accounts

- 1 Chambers, J. S., R. T. Pressey, J. R. Donaldson, and W. R. McKinley. 1954.
2 *Research Relating to Study of Spawning Grounds in Natural Areas.*
3 Annual Report, Contract No. DA 35026-Eng-20572. Prepared by
4 Washington State Department of Fisheries, Olympia, Washington, for
5 U.S. Army Corps of Engineers, Fisheries-Engineering Research Program,
6 North Pacific Division, Portland, Oregon.
- 7 Chambers, J. S., G. H. Allen, and R. T. Pressey. 1955. *Research Relating to Study*
8 *of Spawning Grounds in Natural Areas.* Annual Report, Contract No. DA
9 35026-Eng-20572. Prepared by Washington State Department of
10 Fisheries, Olympia, Washington, for U.S. Army Corps of Engineers,
11 Fisheries-Engineering Research Program, North Pacific Division,
12 Portland, Oregon.
- 13 Chapman, D. W., D. E. Weitkamp, T. L. Welsh, M. B. Dell, and T. H. Schadt.
14 1986. Effects of river flow on the distribution of Chinook salmon redds.
15 *Transactions of the American Fisheries Society* 115: 537-547.
- 16 Chelan County Public Utility District. 1989. *Summer and Winter Ecology of*
17 *Juvenile Chinook Salmon and Steelhead Trout in the Wenatchee River,*
18 *Washington.* Prepared by Don Chapman Consultants for Chelan County
19 Public Utility District, Wenatchee, Washington.
- 20 Chilcote, S., A. Collins, A. Cousins, N. Hemphill, A. Hill, and J. Smith. 2013.
21 *Spring Chinook in the SFTR Rivers: Recommended Management Actions*
22 *and the Status of their Implementation.* Trinity River Restoration Program,
23 South Fork Trinity River Spring Chinook Subgroup.
- 24 Clark, G. H. 1943. Salmon at Friant Dam - 1942. *California Fish and Game* 29:
25 89-91
- 26 Clarke, W. C., and J. E. Shelbourn. 1985. Growth and development of seawater
27 adaptability by juvenile fall Chinook salmon (*Oncorhynchus tshawytscha*)
28 in relation to temperature. *Aquaculture* 45: 21-31.
- 29 Coble, D. W. 1961. Influence of water exchange and dissolved oxygen in redds
30 on survival of steelhead trout embryos. *Transactions of the American*
31 *Fisheries Society* 90: 469-474.
- 32 Combs, B. D. 1965. Effect of temperature on the development of salmon eggs.
33 *The Progressive Fish-Culturist* 27: 134-137.
- 34 Combs, B. D., and R. E. Burrows. 1957. Threshold temperatures for the normal
35 development of Chinook salmon eggs. *The Progressive Fish-Culturist* 19:
36 3-6.
- 37 Cooper, A. C. 1965. *The Effect of Transported Stream Sediments on the Survival*
38 *of Sockeye and Pink Salmon Eggs and Alevin.* Bulletin 18. International
39 Pacific Salmon Fisheries Commission, New Westminster, British
40 Columbia, Canada.

- 1 Dean, M. 1994. Life history, distribution, run size, and harvest of spring-run
2 Chinook salmon in the South Fork Trinity River Basin. Chapter VII - job
3 VII in *Trinity River Basin Monitoring Project 1991-1992*.
- 4 _____. 1995. Life history, distribution, run size, and harvest of spring-run
5 Chinook salmon in the South Fork Trinity River Basin. Chapter VII - job
6 VII in *Trinity River Basin Monitoring Project 1992-1993*.
- 7 DeHaven, R. W. 1989. Distribution, Extent, Replaceability and Relative Values to
8 Fish and Wildlife of Shaded Riverine Aquatic Cover of the Lower
9 Sacramento River, California. Part I: 1987-88 Study Results and
10 Recommendations. Prepared by U.S. Fish and Wildlife Service,
11 Sacramento, California, for U.S. Army Corps of Engineers, Sacramento
12 District, Sacramento, California. As cited by Fris and Dehaven 1993.
- 13 Del Rosario, R., Y. J. Redler, K. Newman, P. L. Brandes, T. Sommer, K. Reece,
14 and R. Vincik. 2013. Migration patterns of juvenile winter-run-sized
15 Chinook salmon (*Oncorhynchus tshawytscha*) through the Sacramento–
16 San Joaquin Delta. *San Francisco Estuary and Watershed Science* 11(1).
17 <http://www.escholarship.org/uc/item/36d88128>.
- 18 DFG (California Department of Fish and Game). 1965. *California Fish and*
19 *Wildlife Plan*. DFG, Inland Fisheries Division, Sacramento, California.
- 20 _____. 1982. Sacramento River and Tributaries Bank Protection and Erosion
21 Control Investigation--Evaluation of Impacts on Fisheries. Final Report.
22 CDFG, Bay-Delta Fishery Project, Sacramento, California.
- 23 _____. 1995. Fish Species of Special Concern in California, Spring-run Chinook
24 Salmon. Habitat Conservation Planning Branch.
- 25 _____. 1997. Central Valley Anadromous Fish-Habitat Evaluations: Sacramento
26 and American River Investigations, October 1995 through September
27 1996. Stream Evaluation Program, Technical Report No. 97-1. Prepared
28 by CDFG, Environmental Services Division, Stream Flow and Habitat
29 Evaluation Program for U.S. Fish and Wildlife Service, Central Valley
30 Anadromous Fish Restoration Program.
- 31 _____. 1998. *A Status Review of the Spring-run Chinook Salmon (Oncorhynchus*
32 *tshawytscha) in the Sacramento River Drainage*. Report to the Fish and
33 Game Commission, Candidate Species Status Report 98-01. CDFG,
34 Sacramento.
- 35 _____. 2002a. *Sacramento River Winter-run Chinook Salmon*. Biennial Report
36 2000-2001. Prepared by CDFG, Habitat Conservation Division, Native
37 Anadromous Fish and Watershed Branch for California Fish and Game
38 Commission.
- 39 _____. 2002b. *Sacramento River Spring-run Chinook Salmon*. Annual report.
40 Prepared by CDFG, Habitat Conservation Division, Native Anadromous
41 Fish and Watershed Branch for Fish and Game Commission.

- 1 _____ . 2004. *Sacramento River Winter-run Chinook Salmon*. Biennial Report
2 2002-2003. Prepared by CDFG, Habitat Conservation Division, Native
3 Anadromous Fish and Watershed Branch for California Fish and Game
4 Commission.
- 5 _____ . 2010. Letter from Terry Foreman, Chief Fisheries Branch to Rod McInnis,
6 Regional Administrator, NMFS concerning the Sacramento River winter-
7 run Chinook escapement estimate for 2010, dated December 8, 2010 As
8 cited in NMFS 2011.
- 9 DFW (California Department of Fish and Wildlife). 2014. *GrandTab*. California
10 Central Valley Sacramento and San Joaquin River systems Chinook
11 salmon escapement, hatcheries and natural areas. Fisheries Branch,
12 Anadromous Resources Assessment. Sacramento.
- 13 Edmundson, E., F. E., Everest, and D. W. Chapman. 1968. Permanence of station
14 in juvenile Chinook salmon and steelhead trout. *Journal of the Fisheries*
15 *Research Board of Canada* 25: 1453–1464.
- 16 Everest, F. H., and D. W. Chapman. 1972. Habitat selection and spatial
17 interaction by juvenile Chinook salmon and steelhead trout in two Idaho
18 streams. *Journal of the Fisheries Research Board of Canada* 29: 91-100.
- 19 Fisher, F. W. 1994. Past and present status of Central Valley Chinook salmon.
20 *Conservation Biology* 8: 870-873.
- 21 Fris, M. B., and R. W. DeHaven. 1993. A Community-Based Habitat Suitability
22 Index Model for Shaded Riverine Aquatic Cover, Selected Reaches of the
23 Sacramento River System. U.S. Fish and Wildlife Service, Sacramento
24 Field Office, Sacramento, California.
- 25 Hallock, R. J., D. H. Fry, Jr., and D. A. LaFaunce. 1957. The use of fyke traps to
26 estimate the runs of adult salmon and steelhead in the Sacramento River.
27 *California Fish and Game* 43: 271-296.
- 28 Hallock, R. J., and F. W. Fisher. 1985. *Status of the Winter-run Chinook Salmon,*
29 *Oncorhynchus tshawytscha, in the Sacramento River*. Anadromous
30 Fisheries Branch Office Report. California Department of Fish and Game.
- 31 Harvey, C. D. 1995. *Juvenile Spring-run Chinook Salmon Emergence, Rearing*
32 *and Outmigration Patterns in Deer Creek and Mill Creek, Tehama County*
33 *for the 1994 Broodyear*. California Department of Fish and Game,
34 Redding.
- 35 Harvey-Arrison, C. 2001. Re: Accounts of winter-run Chinook salmon in Battle
36 and Mill creeks. Internal memorandum to D. Hallock, California
37 Department of Fish and Game, Sacramento. 19 June.
- 38 Hatton, S. R. 1940. Progress report on the Central Valley fisheries investigations,
39 1939. *California Fish and Game* 26:334-369.

- 1 Hausle, D. A., and D. W. Coble. 1976. Influence of sand in redds on survival and
2 emergence of brook trout (*Salvelinus fontinalis*). *Transactions of the*
3 *American Fisheries Society* 105: 57-63.
- 4 Healey, M. C. 1980. *Utilization of the Nanaimo River Estuary by Juvenile*
5 *Chinook Salmon*, *Oncorhynchus tshawytscha*. U.S. Fisheries Bulletin 77:
6 653–668.
- 7 _____. 1983. Coastwide distribution and ocean migration patterns of stream- and
8 ocean-type Chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Field*
9 *Naturalist* 97: 427-433.
- 10 _____. 1991. Life history of Chinook salmon (*Oncorhynchus tshawytscha*). In
11 *Pacific Salmon Life Histories*, pp. 311-393. Edited by C. Groot and L.
12 Margolis. University of British Columbia Press, Vancouver, British
13 Columbia.
- 14 Healey, T. P. 1979. *The Effect of High Temperature on the Survival of*
15 *Sacramento River Chinook (King) Salmon*, *Oncorhynchus tshawytscha*,
16 *Eggs and Fry*. Administrative Report 79-10. California Department of
17 Fish and Game, Anadromous Fisheries Branch.
- 18 Heming, T. A. 1982. Effects of temperature on utilization of yolk by Chinook
19 salmon (*Oncorhynchus tshawytscha*) eggs and alevins. *Canadian Journal*
20 *of Fisheries and Aquatic Sciences* 39: 184-190.
- 21 Hill, K. A., and J. D. Webber. 1999. *Butte Creek Spring-run Chinook Salmon*,
22 *Oncorhynchus tshawytscha*, *Juvenile Outmigration and Life History 1995-*
23 *1998*. Inland Fisheries Administrative Report No. 99-5. California
24 Department of Fish and Game, Sacramento Valley and Central Sierra
25 Region, Rancho Cordova, California.
- 26 Hillman, T. W., J. S. Griffith, and W. S. Platts. 1987. Summer and winter habitat
27 selection by juvenile Chinook salmon in a highly sedimented Idaho stream.
28 *Transactions of the American Fisheries Society* 116: 185-195.
- 29 Hinze, J. A. 1959. *Annual Report, Nimbus Salmon and Steelhead Hatchery,*
30 *Fiscal Year of 1957-58*. Inland Fisheries Administrative Report 59-4.
31 California Department of Fish and Game.
- 32 Johnson, R., D. C. Weigand, and F. W. Fisher. 1992. *Use of Growth Data to*
33 *Determine the Spatial and Temporal Distribution of Four Runs of Juvenile*
34 *Chinook Salmon in the Sacramento River, California*. Report No. AFF1-
35 FRO-92-15. U.S. Fish and Wildlife Service. As cited by The Nature
36 Conservancy *Sacramento River Ecological Flows Study: State of System*
37 *Report*, November 2006.
- 38 Kano, B. 2006. *GrandTab; Central Valley Streams Chinook Salmon Escapement*
39 *Database*. California Department of Fish and Game. Native Anadromous
40 Fish and Watershed Branch. Red Bluff, California. As cited by
41 Department of Fish and Wildlife *Annual Report Chinook Salmon Spawner*
42 *Stocks in California's Central Valley*, 2004.

- 1 Kimmerer, W., L. Brown, S. Culberson, P. Moyle, M. Nobriga, and J. Thompson.
2 2008. Aquatic ecosystems. In *The State of Bay-Delta Science 2008*, pp.
3 55-72. Edited by M. Healey, M. Dettinger, and R. Norgaard. CALFED
4 Science Program, Sacramento, California.
- 5 Kjelson, M. A., P. F. Raquel, and F. W. Fisher. 1981. Influences of freshwater
6 inflow on Chinook salmon (*Oncorhynchus tshawytscha*) in the
7 Sacramento-San Joaquin Estuary. In *Proceedings of the National*
8 *Symposium on Freshwater Inflow to Estuaries*, pp. 88-108. Edited by R.
9 D. Cross and D. L. Williams. FWS/OBS-81/04. U.S. Fish and Wildlife
10 Service, Washington, D. C.
- 11 Kondolf, G. M. 2000. Assessing salmonid spawning gravel quality. *Transactions*
12 *of the American Fisheries Society* 129: 262-281.
- 13 Kondolf, G. M., M. J. Sale, and M. G. Wolman. 1993. Modification of fluvial
14 gravel size by spawning salmonids. *Water Resources Research* 29: 2265-
15 2274.
- 16 Koski, K. V. 1981. The survival and quality of two stocks of chum salmon
17 (*Oncorhynchus keta*) from egg deposition to emergence. *Rapports et*
18 *Proces-Verbaux des Reunions, Conseil International pour L'Exploration*
19 *de la Mer* 178: 330-333.
- 20 Levings, C. D., and R. B. Lauzier. 1991. Extensive use of the Fraser River basin
21 as winter habitat by juvenile Chinook salmon (*Oncorhynchus*
22 *tshawytscha*). *Canadian Journal of Zoology* 69: 1759-1767.
- 23 Lindley, S. T., R. Schick, B. P. May, J. J. Anderson, S. Greene, C. Hanson, A.
24 Low, D. McEwan, R. B. MacFarlane, C. Swanson, and J. G. Williams.
25 2004. *Population Structure of Threatened and Endangered Chinook*
26 *Salmon ESUs in California's Central Valley Basin*. Technical
27 Memorandum NOAA-TM-NMFS-SWFSC-360. National Marine
28 Fisheries Service, Southwest Fisheries Science Center.
- 29 Lindley, S. T., R. Schick, A. Agrawal, M. Goslin, T. E. Pearson, E. Mora, J. J.
30 Anderson, B. May, S. Greene, C. Hanson, A. Low, D. McEwan, R. B.
31 MacFarlane, C. Swanson, and J. G. Williams. 2006. Historical population
32 structure of Central Valley steelhead and its alteration by dams.
33 *San Francisco Estuary and Watershed Science* [online serial] 4(2).
- 34 Lindsay, R. B., W. J. Knox, M. W. Flesher, B. J. Smith, E. A. Olsen, and L. S.
35 Lutz. 1986. *Study of Wild Spring-run Chinook Salmon in the John Day*
36 *River System*. 1985 Final Report. Contract DE-AI79-83BP39796, Project
37 79-4. Prepared by Oregon Department of Fish and Wildlife, Portland for
38 Bonneville Power Administration, Portland, Oregon.
- 39 Lister, D. B., and H. S. Genoe. 1970. Stream habitat utilization of cohabiting
40 underyearlings of Chinook (*Oncorhynchus tshawytscha*) and coho (*O.*
41 *kisutch*) salmon in the Big Qualicum River, British Columbia. *Journal of*
42 *the Fisheries Research Board of Canada* 27: 1215-1224.

- 1 Marcotte, B. D. 1984. *Life History, Status, and Habitat Requirements of Spring-*
2 *run Chinook Salmon in California*. U.S. Forest Service, Lassen National
3 Forest, Chester, California.
- 4 Marine, K. R. 1992. *A Background Investigation and Review of the Effects of*
5 *Elevated Water Temperature on Reproductive Performance of Adult*
6 *Chinook Salmon (Oncorhynchus tshawytscha)*. Department of Wildlife
7 and Fisheries Biology, University of California, Davis. As cited by
8 Department of Water Resources *Matrix of Life History and Habitat*
9 *Requirements for Feather River Fish Species*, SP-F15 Task 1 and SP-F21
10 Task 1 Oroville Facilities Relicensing FERC Project No. 2100, April
11 2004.
- 12 _____. 1997. *Effects of Elevated Water Temperature on Some Aspects of the*
13 *Physiological and Ecological Performance of Juvenile Chinook Salmon*
14 *(Oncorhynchus tshawytscha): Implications for Management of*
15 *California's Central Valley Salmon Stocks*. Master's thesis. University of
16 California, Davis.
- 17 Marine, K. R., and J. J. Cech, Jr. 2004. Effects of high water temperature on
18 growth, smoltification, and predator avoidance in juvenile Sacramento
19 River Chinook salmon. *North American Journal of Fisheries Management*
20 24: 198-210.
- 21 Martin, C. D., P. D. Gaines, and R. R. Johnson. 2001. *Estimating the Abundance*
22 *of Sacramento River Juvenile Winter Chinook Salmon with Comparisons*
23 *to Adult Escapement*. Final Report, Report Series: Volume 5. July.
24 Prepared by U.S. Fish and Wildlife Service, Red Bluff, CA. Prepared for
25 U.S. Bureau of Reclamation, Red Bluff, CA.
- 26 Maslin, P., M. Lennox, J. Kindopp, and W. McKinney. 1997. *Intermittent*
27 *Streams as Rearing Habitat for Sacramento River Chinook Salmon*
28 *(Oncorhynchus tshawytscha)*. Department of Biological Sciences,
29 California State University, Chico.
- 30 McCain, M. E. 1992. Comparison of habitat use and availability for juvenile fall
31 Chinook salmon in a tributary of the Smith River, California. *FHR*
32 *Currents* No. 7. U.S. Forest Service, Region 5.
- 33 McCuddin, M. E. 1977. *Survival of Salmon and Trout Embryos and Fry in*
34 *Gravel-sand Mixtures*. Master's thesis. University of Idaho, Moscow. As
35 cited in Kondolf 2000.
- 36 McNeil, W. J. 1964a. Effect of the spawning bed environment on reproduction of
37 pink and chum salmon. *U.S. Fish and Wildlife Service Fishery Bulletin* 65:
38 495-523.
- 39 _____. 1964b. Redd superimposition and egg capacity of pink salmon spawning
40 beds. *Journal of the Fisheries Research Board of Canada* 21: 1385-1396.

- 1 McReynolds, T. R., C. E. Garman, P. D. Ward, and M. C. Schommer. 2005. *Butte*
2 *and Big Chico Creeks Spring-run Chinook Salmon, Oncorhynchus*
3 *tshawytscha, Life History Investigation 2003-2004*. Inland Fisheries
4 Administrative Report No. 2005-1. California Department of Fish and
5 Game, Sacramento Valley and Central Sierra Region, Rancho Cordova,
6 California.
- 7 Meyer, J. H. 1979. *A Review of the Literature on the Value of Estuarine and*
8 *Shoreline Areas to Juvenile Salmonids in Puget Sound, Washington*. U.S.
9 Fish and Wildlife Service, Fisheries Assistance Office, Olympia,
10 Washington.
- 11 Michny, F. 1987. *Sacramento River, Chico Landing to Red Bluff Project, 1986*
12 *Juvenile Salmon Study*. Prepared by U.S. Fish and Wildlife Service,
13 Sacramento for U.S. Army Corps of Engineers, Sacramento, California.
14 As cited by U.S. Fish and Wildlife Service *Shaded Riverine Aquatic Cover*
15 *of the Sacramento River System: Classification as Resources Category 1*
16 *Under the Fish and Wildlife Mitigation Policy*, October 1992.
- 17 _____. 1988. *Sacramento River Butte Basin Reach Pre-project Juvenile Salmon*
18 *Study*. Prepared by U.S. Fish and Wildlife Service, Sacramento for U.S.
19 Army Corps of Engineers, Sacramento, California. As cited by U.S. Fish
20 and Wildlife Service *Shaded Riverine Aquatic Cover of the Sacramento*
21 *River System: Classification as Resources Category 1 Under the Fish and*
22 *Wildlife Mitigation Policy*, October 1992.
- 23 _____. 1989. *Sacramento River, Chico Landing to Red Bluff Project, 1987*
24 *Juvenile Salmon Study*. Prepared by U.S. Fish and Wildlife Service,
25 Sacramento for U.S. Army Corps of Engineers, Sacramento, California.
26 As cited by U.S. Fish and Wildlife Service *Shaded Riverine Aquatic Cover*
27 *of the Sacramento River System: Classification as Resources Category 1*
28 *Under the Fish and Wildlife Mitigation Policy*, October 1992.
- 29 Michny, F., and R. Deibel. 1986. *Sacramento River, Chico Landing to Red Bluff*
30 *Project, 1985 Juvenile Salmon Study*. Draft report. Prepared by U.S. Fish
31 and Wildlife Service, Sacramento, California for U.S. Army Corps of
32 Engineers, Sacramento, California.
- 33 Michny, F., and M. Hampton. 1984. *Sacramento River, Chico Landing to Red*
34 *Bluff Project, 1984 Juvenile Salmon Study*. Draft report. Prepared by U.S.
35 Fish and Wildlife Service, Sacramento, California for U.S. Army Corps of
36 Engineers, Sacramento, California.
- 37 Mills, T. J., and F. Fisher. 1994. *Central Valley Anadromous Sport Fish Annual*
38 *Run-size, Harvest, and Population Estimates, 1967 through 1991*. Inland
39 Fisheries Technical Report. California Department of Fish and Game.
- 40 Moore, J. W., D. E. Schindler, and M. D. Scheuerell. 2004. Disturbance of
41 freshwater habitats by anadromous salmon in Alaska. *Oecologia* 139: 298-
42 308.

- 1 Moyle, P. B. 2000. Abstract 89. *CALFED Bay-Delta Program Science*
2 *Conference 2000*. Edited by R. L. Brown, F. H. Nichols and L. H. Smith.
3 CALFED Bay-Delta Program, Sacramento, California. As cited by
4 CALFED Bay-Delta Program Science Conference 2000.
- 5 _____. 2002. *Inland Fishes of California*. Revised edition. University of
6 California Press, Berkeley.
- 7 Moyle, P. B., R. M. Yoshiyama, J. E. Williams, and E. D. Wikramanayake. 1995.
8 *Fish Species of Special Concern in California*. Final Report. Prepared by
9 Department of Wildlife and Fisheries Biology, University of California,
10 Davis for California Department of Fish and Game, Inland Fisheries
11 Division, Rancho Cordova.
- 12 Murphy, M. L., J. Heifetz, J. F. Thedinga, S. W. Johnson, and K. V. Koski. 1989.
13 Habitat utilization by juvenile Pacific salmon (*Oncorhynchus*) in the
14 glacial Taku River, southeast Alaska. *Canadian Journal of Fisheries and*
15 *Aquatic Sciences* 46: 1677-1685.
- 16 Myers, J. M., R. G. Kope, G. J. Bryant, D. Teel, L. J. Lierheimer, T. C.
17 Wainwright, W. S. Grant, F. W. Waknitz, K. Neely, S. T. Lindley, and R.
18 S. Waples. 1998. *Status Review of Chinook Salmon from Washington,*
19 *Idaho, Oregon, and California*. NOAA Technical Memorandum NMFS-
20 NWFSC-35. National Marine Fisheries Service, Northwest Fisheries
21 Science Center, Seattle, Washington.
- 22 Myrick, C. A., and J. J. Cech, Jr. 2001. *Temperature Effects on Chinook Salmon*
23 *and Steelhead: a Review Focusing on California's Central Valley*
24 *Populations*. Prepared by Department of Fishery and Wildlife Biology,
25 Colorado State University, Fort Collins and Department of Wildlife, Fish,
26 and Conservation Biology, University of California, Davis for the Bay-
27 Delta Modeling Forum.
- 28 _____. 2002. Growth of American River fall-run Chinook salmon in California's
29 Central Valley: temperature and ration effects. *California Fish and Game*
30 88:35-44.
- 31 _____. 2004. Temperature effects on juvenile anadromous salmonids in
32 California's Central Valley: what don't we know? *Reviews in Fish Biology*
33 *and Fisheries* 14: 113–123.
- 34 Needham, P. R., H. A. Hanson, and L. P. Parker. 1943. *Supplementary Report on*
35 *Investigations of Fish-salvage Problems in Relation to Shasta Dam*.
36 Special Scientific Report No. 26. U.S. Fish and Wildlife Service.
- 37 Neilson, J. D., and C. E. Banford. 1983. Chinook salmon (*Oncorhynchus*
38 *tshawytscha*) spawner characteristics in relation to redd physical features.
39 *Canadian Journal of Zoology* 61:1524-1531.

- 1 Newman, K. B., and P. L. Brandes. 2010. Hierarchical modeling of juvenile
2 Chinook Salmon survival as a function of Sacramento–San Joaquin Delta
3 water exports. *North American Journal of Fisheries Management* 30:157–
4 169.
- 5 Nicholas, J. W., and D. G. Hankin. 1989. *Chinook Salmon Populations in Oregon*
6 *Coastal River Basins: Descriptions of Life Histories and Assessment of*
7 *Recent Trends in Run Strengths*. Report EM 8402. Oregon Department of
8 Fish and Wildlife, Research and Development Section, Corvallis.
- 9 NMFS (National Marine Fisheries Service). 1989. Endangered and threatened
10 species; critical habitat; winter-run Chinook salmon. *Federal Register* 54:
11 32085-32088
- 12 _____. 1993. Designated critical habitat; Sacramento River winter-run Chinook
13 salmon. *Federal Register* 58: 33212-33219.
- 14 _____. 1994. Endangered and threatened species; status of Sacramento River
15 winter-run Chinook salmon. *Federal Register* 59: 440-450.
- 16 _____. 1997. *NMFS Proposed recovery plan for the Sacramento River winter-run*
17 *Chinook salmon*. NMFS, Southwest Region, Long Beach, California.
- 18 _____. 1999a. Endangered and threatened species; threatened status for two
19 Chinook salmon evolutionarily significant units (ESUs) in California.
20 *Federal Register* 64: 50394-50415.
- 21 _____. 1999b. *Status Review Update for Deferred ESUs of West Coast Chinook*
22 *Salmon (Oncorhynchus tshawytscha) from Washington, Oregon,*
23 *California, and Idaho*. Report of West Coast Biological Review Team to
24 NMFS, Seattle, Washington.
25 [http://www.nwr.noaa.gov/Publications/Biological-Status-](http://www.nwr.noaa.gov/Publications/Biological-Status-Reviews/loader.cfm?csModule=security/getfile&pageid=21676)
26 [Reviews/loader.cfm?csModule=security/getfile&pageid=21676](http://www.nwr.noaa.gov/Publications/Biological-Status-Reviews/loader.cfm?csModule=security/getfile&pageid=21676).
- 27 _____. 2004a. Endangered and threatened species: proposed listing
28 determinations for 27 ESUs of west coast salmonids. *Federal Register* 69:
29 33102-33179.
- 30 _____. 2004a. *Biological Opinion on the Long-term Central Valley Project and*
31 *State Water Project Operations Criteria and Plan*. Endangered Species
32 Act Section 7 Consultation. NMFS, Southwest Region, Long Beach,
33 California.
- 34 _____. 2004b. Endangered and threatened species: establishment of Species of
35 Concern list, addition of species to Species of Concern list, description of
36 factors for identifying Species of Concern, and revision of Candidate
37 Species list Under the Endangered Species Act: notice. *Federal Register*
38 69: 19975-19979.
- 39 _____. 2005. Endangered and threatened species; final listing determinations for
40 16 ESUs of West Coast salmon, and final 4(d) protective regulations for
41 threatened salmonid ESUs. *Federal Register* 70: 37160-37204.

- 1 _____ . 2009. *Public Draft Recovery Plan for the Evolutionarily Significant Units*
2 *of Sacramento River Winter-Run Chinook Salmon and Central Valley*
3 *Spring-Run Chinook Salmon and the Distinct Population Segment of*
4 *Central Valley Steelhead*. October. Sacramento Protected Resources
5 Division, Sacramento, CA.
- 6 _____ . 2011. *Central Valley Recovery Domain 5-Year Review: Summary and*
7 *Evaluation of Sacramento River Winter-run Chinook Salmon ESU*. NMFS,
8 Southwest Region, Long Beach, California.
- 9 _____ . 2012. Listing Endangered and Threatened species; 12-month finding on a
10 petition to list Chinook salmon in the Upper Klamath and Trinity rivers
11 basin as Threatened or Endangered under the Endangered Species Act.
12 *Federal Register* 77: 19597-19605. [http://www.gpo.gov/fdsys/pkg/FR-](http://www.gpo.gov/fdsys/pkg/FR-2012-04-02/pdf/2012-7879.pdf)
13 [2012-04-02/pdf/2012-7879.pdf](http://www.gpo.gov/fdsys/pkg/FR-2012-04-02/pdf/2012-7879.pdf).
- 14 Northern California Water Association and Sacramento Valley Water Users.
15 2011. *Insights into the Problems, Progress, and Potential Solutions for*
16 *Sacramento River Basin Native Anadromous Fish Restoration*. Prepared
17 by D. Vogel for Northern California Water Association and Sacramento
18 Valley Water Users. Red Bluff, California.
- 19 NRC (National Research Council). 2004. *Endangered and Threatened Fishes in*
20 *the Klamath River Basin: Causes of Decline and Strategies for Recovery*.
21 The National Academies Press, Washington, D.C.
22 <http://www.nap.edu/openbook.php?isbn=0309090970>.
- 23 Oregon Department of Fish and Wildlife. 1987. *Abundance of Rogue River Fall*
24 *Chinook Salmon*. Annual Progress Report, Fish Research Project Contract
25 AFS-78-1. Prepared by S.P. Cramer for Oregon Department of Fish and
26 Wildlife, Portland.
- 27 Parker, L. P., and H. A. Hanson. 1944. Experiments on transfer of adult salmon
28 into Deer Creek, California. *Journal of Wildlife Management* 8: 192-198.
- 29 Perry, R. W., J. G. Romine, N. S. Adams, A. R. Blake, J. R. Burau, S. V.
30 Johnston, and T. L. Liedtke. 2012. Using a non-physical behavioural
31 barrier to alter migration routing of juvenile Chinook salmon in the
32 Sacramento–San Joaquin River delta. *River Research and Applications*,
33 n/a-n/a. doi: 10.1002/trr.2628
- 34 Perry, R. W., J. R. Skalski, P. L. Brandes, P. T. Sandstrom, A. P. Klimley, A.
35 Ammann, and B. MacFarlane. 2010. Estimating survival and migration
36 route probabilities of juvenile Chinook salmon in the Sacramento–
37 San Joaquin River delta. *North American Journal of Fisheries*
38 *Management* 30:142–156.
- 39 Peterson, D. P., and C. J. Foote. 2000. Disturbance of small-stream habitat by
40 spawning sockeye salmon in Alaska. *Transactions of the American*
41 *Fisheries Society* 129: 924-934.

- 1 PFMC (Pacific Fishery Management Council). 2008. *Review of 2007 Ocean*
2 *Salmon Fisheries*. Portland, Oregon. www.pcouncil.org.
- 3 Phillips, R. W., R. L. Lantz, E. W. Claire, and J. R. Moring. 1975. Some effects of
4 gravel mixtures on emergence of coho salmon and steelhead trout fry.
5 *Transactions of the American Fisheries Society* 104: 461-466.
- 6 Platts, W. S., M. A. Shirazi, and D. H. Lewis. 1979. *Sediment Particle Sizes Used*
7 *by Salmon for Spawning with Methods for Evaluation*. Ecological
8 Research Series EPA-600/3-79-043. U.S. Environmental Protection
9 Agency, Corvallis Environmental Research Laboratory, Corvallis, Oregon.
- 10 Reclamation (U.S. Bureau of Reclamation). 1991. *Guide to Upper Sacramento*
11 *River Chinook Salmon Life History*. Prepared by D. A. Vogel and K. R.
12 Marine, CH2M HILL, Redding, California, for U.S. Bureau of
13 Reclamation, Central Valley Project.
- 14 _____. 1994. *Action Plan for the Restoration of the SFTR Watershed and its*
15 *Fishes*. Prepared by Pacific Watershed Associates for U.S. Bureau of
16 Reclamation and Trinity River Task Force, Arcata, California. As cited by
17 Trinity River Restoration Program *Spring Chinook in the South Fork*
18 *Trinity River: Recommended Management Actions and the Status of their*
19 *Implementation*, January 29, 2013.
- 20 _____. 2004. *Long-term Central Valley Project and State Water Project*
21 *Operations Criteria and Plan Biological Assessment*. USDI Bureau of
22 Reclamation, Mid-Pacific Region, Sacramento, California.
- 23 Reiser, D. W., and R. T. Peacock. 1985. *A technique for assessing upstream fish*
24 *passage problems at small-scale hydropower developments*. Edited by F.
25 W. Olson, R. G. White, and R. H. Hamre. Pp. 423-432. Symposium on
26 Small Hydropower and Fisheries. American Fisheries Society, Bethesda,
27 Maryland.
- 28 Reiser, D. W., and R. G. White. 1988. Effects of two sediment size-classes on
29 survival of steelhead and Chinook salmon eggs. *North American Journal*
30 *of Fisheries Management* 8: 432-437.
- 31 Rich, A. A. 1987. Report on studies conducted by Sacramento County to
32 determine the temperatures which optimize growth and survival in
33 juvenile Chinook salmon (*Oncorhynchus tshawytscha*). Prepared for
34 McDonough, Holland and Allen, Sacramento, California, by A. A. Rich
35 and Associates, San Rafael, California.
- 36 Roper, B. R., D. L. Scarnecchia, and T. J. La Marr. 1994. Summer distribution of
37 and habitat use by Chinook salmon and steelhead within a major basin of
38 the South Umpqua River, Oregon. *Transactions of the American Fisheries*
39 *Society* 123: 298-308.
- 40 Rutter, C. 1908. The fishes of the Sacramento-San Joaquin basin, with a study of
41 their distribution and variation. *Bulletin of the U.S. Bureau of Fisheries*
42 27: 103-152.

- 1 Shirvell, C. S. 1994. Effect of changes in streamflow on the microhabitat use and
2 movements of sympatric juvenile coho salmon (*Oncorhynchus kisutch*)
3 and Chinook salmon (*O. tshawytscha*) in a natural stream. *Canadian*
4 *Journal of Fisheries and Aquatic Sciences* 51: 1644-1652.
- 5 Shumway, D. L., C. E. Warren, and P. Doudoroff. 1964. Influence of oxygen
6 concentration and water movement on the growth of steelhead trout and
7 coho salmon embryos. *Transactions of the American Fisheries Society* 93:
8 342-356.
- 9 Silver, S. J., C. E. Warren, and P. Doudoroff. 1963. Dissolved oxygen
10 requirements of developing steelhead trout and Chinook salmon embryos
11 at different velocities. *Transactions of the American Fisheries Society* 92:
12 327-343.
- 13 Slater, D. W. 1963. *Winter-run Chinook salmon in the Sacramento River,*
14 *California with Notes on Water Temperature Requirements at Spawning.*
15 Special Scientific Report—Fisheries 461. U.S. Fish and Wildlife Service.
- 16 Smith, A. K. 1973. Development and application of spawning velocity and depth
17 criteria for Oregon salmonids. *Transactions of the American Fisheries*
18 *Society* 102: 312-316.
- 19 Smith, S. H. 1950. *Upper Sacramento River Sport Fishery.* Special Scientific
20 Report – Fisheries. U.S. Fish and Wildlife Service.
- 21 Snider, B., B. Reavis, and S. Hill. 1998. *Upper Sacramento River Late-fall-run*
22 *Chinook Salmon Escapement Survey, December 1997-May 1998.* Stream
23 Evaluation Program Technical Report No. 98-4. California Department of
24 Fish and Game, Environmental Services Division.
- 25 _____. 1999. *Upper Sacramento River Late-fall-run Chinook Salmon Escapement*
26 *Survey, December 1998 April 1999.* Stream Evaluation Program Technical
27 Report No. 99-3. California Department of Fish and Game, Habitat
28 Conservation Division, Native Anadromous Fish and Watershed Branch.
- 29 _____. 2000. *Upper Sacramento River Late-fall-run Chinook Salmon Escapement*
30 *Survey, December 1999 April 2000.* Stream Evaluation Program Technical
31 Report No. 00-9. California Department of Fish and Game, Habitat
32 Conservation Division, Native Anadromous Fish and Watershed Branch.
- 33 _____. 2001. *Upper Sacramento River Winter-run Chinook Salmon Escapement*
34 *Survey, May-August 2000.* Stream Evaluation Program Technical Report
35 No. 01-1. California Department of Fish and Game, Habitat Conservation
36 Division, Native Anadromous Fish and Watershed Branch.
- 37 Snider, B., and R. G. Titus. 2000. *Timing, Composition, and Abundance of*
38 *Juvenile Anadromous Salmonid Emigration in the Sacramento River near*
39 *Knights Landing, October 1996–September 1997.* California Department
40 of Fish and Game, Habitat Conservation Division, Stream Evaluation
41 Program Technical Report No. 00-04.

- 1 Snyder, J. O. 1931. Salmon of the Klamath River, California. *California Fish and*
2 *Game Bulletin* 34:130.
- 3 Sommer, T. R., M. L. Nobriga, W. C. Harrell, W. Batham, and W. J. Kimmerer.
4 2001. Floodplain rearing of juvenile Chinook salmon: evidence of
5 enhanced growth and survival. *Canadian Journal of Fisheries and Aquatic*
6 *Sciences* 58: 325-333.
- 7 State Coastal Conservancy. 2009. *Effects of Sediment Release following Dam*
8 *Removal on the Aquatic Biota of the Klamath River*. Technical report.
9 Prepared by Stillwater Sciences, Arcata, California, for State Coastal
10 Conservancy, Oakland, California.
11 [http://www.usbr.gov/mp/kbao/kbra/docs/other/Klamath%20Dam%20Rem](http://www.usbr.gov/mp/kbao/kbra/docs/other/Klamath%20Dam%20Removal%20Biological%20Analysis_FINAL.pdf)
12 [oval%20Biological%20Analysis_FINAL.pdf](http://www.usbr.gov/mp/kbao/kbra/docs/other/Klamath%20Dam%20Removal%20Biological%20Analysis_FINAL.pdf).
- 13 State Water Contractors. 1990. *Laboratory Information on the Effect of Water*
14 *Temperature on Juvenile Chinook Salmon in the Sacramento and*
15 *San Joaquin Rivers: a Literature Review*. San Francisco Bay/Sacramento-
16 San Joaquin Delta, Water Quality Control Plan Hearings, WQCP-SWC
17 Exhibit 605. Prepared by C. H. Hanson, Tenera Environmental, Berkeley,
18 California, for State Water Contractors, Sacramento, California.
- 19 Steward, C. R., and T. C. Bjornn. 1987. The distribution of Chinook salmon
20 juveniles in pools at three discharges. *Proceedings of the Annual*
21 *Conference, Western Association of Fish and Wildlife Agencies* 67: 364-
22 374.
- 23 Strange, J. 2008. *Adult Chinook Salmon Migration in the Klamath River Basin,*
24 *2007 Biotelemetry Monitoring Study Final Report*. Yurok Tribal Fisheries
25 Program, Klamath, California, and University of Washington, School of
26 Aquatic and Fishery Science, Seattle, Washington, in collaboration with
27 Hoopa Valley Tribal Fisheries, Hoopa, California.
- 28 Stuehrenberg, L. C. 1975. *The Effects of Granitic Sand on the Distribution and*
29 *Abundance of Salmonids in Idaho Streams*. Master's thesis. University of
30 Idaho, Moscow.
- 31 Swales, S., R. B. Lauzier, and C. D. Levings. 1986. Winter habitat preferences of
32 juvenile salmonids in two interior rivers in British Columbia. *Canadian*
33 *Journal of Zoology* 64: 1506-1514.
- 34 Taylor, E. B., and P. A. Larkin. 1986. Current response and agonistic behavior in
35 newly emerged fry of Chinook salmon, *Oncorhynchus tshawytscha*, from
36 ocean- and stream-type populations. *Canadian Journal of Fisheries and*
37 *Aquatic Sciences* 43: 565-573.
- 38 The Nature Conservancy. 2003. *Contrasting Patterns of Juvenile Chinook Salmon*
39 *(Oncorhynchus tshawytscha) Growth, Diet, and Prey Densities in Off-*
40 *channel and Main Stem Habitats on the Sacramento River*. Prepared by
41 M. P. Limm and M. P. Marchetti for The Nature Conservancy, Chico,
42 California.

- 1 Thompson, K. 1972. Determining stream flows for fish life. *Proceedings of the*
2 *Instream Flow Requirement Workshop*, pp. 31-50. Pacific Northwest
3 River Basin Commission, Vancouver, Washington.
- 4 TID/MID (Turlock Irrigation District and Modesto Irrigation District). 1997.
5 *Lower Tuolumne River Annual Report 97-1*. Trinity County Resource
6 Conservation District. 2003. *South Fork Trinity River Water Quality*
7 *Monitoring Project*. Prepared for California Department of Fish and
8 Game, Redding, California.
- 9 ULEP (Umpqua Land Exchange Project). 1998. *Mapping Rules for Chinook*
10 *Salmon* (*Oncorhynchus tshawytscha*). Draft Report. ULEP, Roseburg,
11 Oregon. As cited by The Nature Conservancy *Linking Biological*
12 *Responses to River Processes: Implications for Conservation and*
13 *Management of the Sacramento River—A Focal Species Approach*,
14 November 2007.
- 15 USFS (U.S. Forest Service).1994. *Lower Hayfork Creek Watershed Analysis*.
16 Hayfork Ranger District, Shasta-Trinity National Forest.
- 17 _____. 1996. *Lower Hayfork Creek Watershed Analysis*. Shasta-Trinity National
18 Forest, Hayfork Ranger District.
- 19 _____. 1999. *Middle Hayfork Creek Watershed Analysis*. Hayfork Ranger
20 District, Shasta-Trinity National Forest.
- 21 _____. 2001a. *Hidden Valley, Plummer Creek and Rattlesnake Creek Watershed*
22 *Analysis*. Prepared by Foster Wheeler Environmental Corporation for U.S.
23 Forest Service, Shasta-Trinity National Forest, Redding, California.
- 24 _____. 2001b. *Middle Hayfork-Salt Creek Watershed Analyses*. Prepared by URS
25 Greiner Woodward Clyde for U.S. Forest Service, Shasta-Trinity National
26 Forest, Redding, California.
- 27 _____. 2011. *Snorkel Survey Counts of Spring-run Chinook Salmon on the*
28 *Salmon River, California*. Available from M. Meneks, U.S. Forest Service,
29 Fort Jones, California.
- 30 USFWS (U.S. Fish and Wildlife Service). 1995. *Working Paper on Restoration*
31 *Needs: Habitat Restoration Actions to Double Natural Production of*
32 *Anadromous Fish in the Central Valley of California*. Volume 2. Prepared
33 for the USFWS under direction of the Anadromous Fish Restoration
34 Program Core Group. Stockton, California.
- 35 _____. 1996. *Recovery Plan for the Sacramento-San Joaquin Delta Native*
36 *Fishes*. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon.
- 37 _____. 1999. *Effect of Temperature on Early-life Survival of Sacramento River*
38 *Fall- and Winter-run Chinook Salmon*. Final report. USFWS, Northern
39 Central Valley Fish and Wildlife Office, Red Bluff, California.
- 40 _____. 2003. *Flow-habitat Relationships for Steelhead and Fall, Late-fall and*
41 *Winter-run Chinook Salmon Spawning in the Sacramento River between*

- 1 *Keswick Dam and Battle Creek*. Final report. USFWS, Sacramento Fish
2 and Wildlife Office, Sacramento, California.
- 3 _____. 2004. *Flow-habitat Relationships for Spring-run Chinook Salmon*
4 *Spawning in Butte Creek*. USFWS, Sacramento, California.
- 5 _____. 2005. *Flow-habitat Relationships for Chinook Salmon Rearing in the*
6 *Sacramento River between Keswick Dam and Battle Creek*. USFWS,
7 Sacramento Fish and Wildlife Office, Sacramento, California.
- 8 Van Kirk, R. W., and S. W. Naman. 2008. Relative effects of climate and water
9 use on base-flow trends in the lower Klamath Basin. *Journal of the*
10 *American Water Resources Association* 44: 1-18.
- 11 Van Woert, W. 1958. *Time Pattern of Migration of Salmon and Steelhead into the*
12 *Upper Sacramento River during the 1957-1958 Season*. Inland Fisheries
13 Administrative Report 58-7. California Department of Fish and Game. As
14 cited by Natural Heritage Institute *Estimating Ecologically Based Flow*
15 *Targets for the Sacramento and Feather Rivers*, April 2008.
- 16 Vaux, W. G. 1968. Intragravel flow and interchange of water in a streambed.
17 *Fishery Bulletin* 66: 479-489.
- 18 Vernier, J. M. 1969. *Chronological Table of Embryonic Development of Rainbow*
19 *Trout*. Canada Fisheries and Marine Service Translation Series 3913.
- 20 Vogel, D. A. 1987a. *Estimation of the 1986 Spring-run Chinook Salmon Run in*
21 *Deer Creek, California*. Report No. FR1/FAO-87-3. U.S. Fish and
22 Wildlife Service.
- 23 _____. 1987b. *Estimation of the 1986 Spring-run Chinook Salmon Run in Mill*
24 *Creek, California*. Report No. FR1/FAO-87-12. U.S. Fish and Wildlife
25 Service. As cited by Natural Heritage Institute *Estimating Ecologically*
26 *Based Flow Targets for the Sacramento and Feather Rivers*, April 2008.
- 27 Vogel, D. A., and K. R. Marine. 1991. Guide to the Upper Sacramento River
28 Chinook Salmon Life History. Bureau of Reclamation Central Valley
29 Project.
- 30 Vronskiy, B. B. 1972. Reproductive biology of the Kamchatka River Chinook
31 salmon (*Oncorhynchus tshawytscha* [Walbaum]). *Journal of Ichthyology*
32 12: 259-273.
- 33 Waples, R. S., D. J. Teel, J. M. Myers, and A. R. Marshall. 2004. Life-history
34 divergence in Chinook salmon: historic contingency and parallel
35 evolution. *Evolution* 58: 386-403.
- 36 Ward, P. D., and T. R. McReynolds. 2001. *Butte and Big Chico Creeks Spring-*
37 *run Chinook Salmon, Oncorhynchus tshawytscha, Life History*
38 *Investigation 1998-2000*. Inland Fisheries Administrative Report No.
39 2001-2. California Department of Fish and Game, Sacramento Valley and
40 Central Sierra Region, Rancho Cordova, California.

- 1 Ward, P. D., T. R. McReynolds, and C. E. Garman. 2004. *Butte and Big Chico*
2 *Creeks Spring-run Chinook Salmon, Oncorhynchus tshawytscha, Life*
3 *History Investigation 2002–2003*. Inland Fisheries Administrative Report
4 No. 2004-6. California Department of Fish and Game, Sacramento Valley
5 and Central Sierra Region, Rancho Cordova, California.
- 6 West, J. R., O. J. Dix, A. D. Olson, M. V. Anderson, S. A. Fox, and J. H. Power.
7 1990. *Evaluation of Fish Habitat Conditions and Utilization in Salmon,*
8 *Scott, Shasta, and Mid-Klamath Sub-basin Tributaries*. Annual report for
9 Interagency Agreement 14-16-0001-89508. Prepared by U.S. Forest
10 Service, Klamath National Forest, Yreka, California, and Shasta-Trinity
11 National Forest, Weaverville, California.
- 12 Wickett, W. P. 1954. The oxygen supply to salmon eggs in spawning beds.
13 *Journal of the Fisheries Research Board of Canada* 11: 933-953.
- 14 Williams, J. G. 2006. Central Valley salmon: a perspective on Chinook and
15 steelhead in the Central Valley of California. *San Francisco Estuary and*
16 *Watershed Science* 4 (3).
- 17 _____. 2012. Juvenile Chinook salmon (*Oncorhynchus tshawytscha*) in and
18 around the San Francisco Estuary. *San Francisco Estuary and Watershed*
19 *Science* 10 (3).
- 20 Williams, T. H., J. C. Garza, N. Hetrick, S. T. Lindley, M. S. Mohr, J. M. Myers,
21 M. R. O'Farrell, R. M. Quinones, and D. J. Teel. 2011. *Upper Klamath*
22 *and Trinity River Chinook Salmon Biological Review Team Report*.
23 National Marine Fisheries Service, Southwest Fisheries Science Center,
24 La Jolla, California.
- 25 Yoshiyama, R. M., F. W. Fisher, and P. B. Moyle. 1998. Historical abundance
26 and decline of Chinook salmon in the Central Valley region of California.
27 *North American Journal of Fisheries Management* 18: 487-521.
- 28 Yoshiyama, R. M., E. R. Gerstung, F. W. Fisher, and P. B. Moyle. 1996.
29 *Historical and Present Distribution of Chinook Salmon in the Central*
30 *Valley Drainage of California*. In Volume III: *Assessments, Commissioned*
31 *Reports, and Background Information*, pp. 309-362. *Sierra Nevada*
32 *Ecosystem Project: Final Report to Congress*. University of California,
33 Center for Water and Wildland Resources, Davis.
- 34 _____. 2001. Historical and present distribution of Chinook salmon in the Central
35 Valley drainage of California. In *Contributions to the Biology of Central*
36 *Valley Salmonids*, pp. 71-176. Edited by R. L. Brown. Fish Bulletin 179,
37 Volume 1. California Department of Fish and Game, Sacramento.

1 **9B.6 Central Valley Steelhead (*Oncorhynchus***
2 ***mykiss*)**

3 **9B.6.1 Legal Status**

4 Federal: Threatened; Designated Critical Habitat

5 State: None

6 NMFS listed the Central Valley Steelhead ESU as threatened under the Federal
7 ESA in 1998 (NMFS 1998). In 2004, NMFS proposed that all west coast
8 steelhead ESUs be reclassified to DPSs and proposed to retain Central Valley
9 Steelhead as threatened. In January 2006, after a status review (Good et al. 2005),
10 NMFS issued its final decision to retain the status of Central Valley Steelhead as
11 threatened (NMFS 2006).

12 Designated critical habitat for Central Valley Steelhead includes stream reaches of
13 the American, Feather, Yuba, and Bear rivers and their tributaries and tributaries
14 of the Sacramento River including Deer, Mill, Battle, Antelope, and Clear creeks
15 in the Sacramento River basin; the Mokelumne, Calaveras, Stanislaus, Tuolumne,
16 and Merced rivers in the San Joaquin River basin; and portions of the Sacramento
17 and San Joaquin rivers. Designated critical habitat in the Delta includes portions
18 of the Delta Cross Channel Yolo Bypass, Ulati Creek, and portions of the
19 network of channels in the Sacramento River portion of the Delta as well as
20 portions of the San Joaquin, Cosumnes, and Mokelumne rivers and portions of the
21 network of channels in the San Joaquin portion of the Delta.

22 The DPS includes naturally spawned anadromous *O. mykiss* (steelhead)
23 populations below natural and manmade impassable barriers in the Sacramento
24 and San Joaquin rivers and their tributaries, excluding steelhead from
25 San Francisco and San Pablo bays and their tributaries and those from two
26 artificial propagation programs: the Coleman Nimbus Fish Hatchery and Feather
27 River Hatchery steelhead hatchery programs.

28 NMFS considered including resident *O. mykiss* in listed steelhead DPSs in certain
29 instances, including (1) where resident *O. mykiss* have the opportunity to
30 interbreed with anadromous fish below natural or artificial barriers, or (2) where
31 resident fish of native lineage once had the ability to interbreed with anadromous
32 fish but no longer do because they are above artificial barriers and are considered
33 essential for the recovery of the DPS (NMFS 1998). However, USFWS, which
34 under the ESA has authority over resident fish, concluded that behavioral forms
35 of *O. mykiss* can be regarded as separate DPSs and that lacking evidence that
36 resident Rainbow Trout need ESA protection, only anadromous forms should be
37 included in the DPS and listed under the ESA (NMFS 1998). USFWS also did
38 not believe that steelhead recovery would rely on the intermittent exchange of
39 genetic material between resident and anadromous forms. In the final rule, the
40 listing includes only the anadromous form of *O. mykiss*.

41 However, NMFS considers all *O. mykiss* that have access to the ocean (including
42 resident Rainbow Trout) to potentially be steelhead and will treat these fish as
43 steelhead because (1) resident fish can produce anadromous offspring, and (2) it is

1 difficult or impossible to distinguish between juveniles of the different forms.
2 Adult resident Rainbow Trout in Central Valley streams are often larger than
3 Central Valley Steelhead. Several sources indicate that resident trout in the
4 Central Valley commonly exceed 16 inches (406 mm) in length. Cramer et al.
5 (1995) reported that resident Rainbow Trout in Central Valley rivers grow longer
6 than 20 inches (508 mm). Hallock et al. (1961) observed resident trout in the
7 upper Sacramento River upstream of the Feather River that were 14 to 20 inches
8 (356 to 508 mm) in length. Also, at Coleman National Fish Hatchery, USFWS
9 found about 15 percent overlap in size distribution between resident and
10 anadromous *O. mykiss* at a length of 22.8 inches (579 mm) (Cramer et al. 1995).
11 Steelhead, therefore, have significant size overlap with resident Rainbow Trout in
12 Central Valley rivers, and many resident adult trout will be considered by NMFS
13 to be steelhead.

14 The following profiles focus on the anadromous form of the species because these
15 are the most likely to be affected by the proposed action, and several have special
16 status under the ESA.

17 **9B.6.2 Distribution**

18 Central Valley Steelhead are widely distributed throughout their range but are low
19 in abundance, particularly in the San Joaquin River basin, and they continue to
20 decline (NMFS 2003). Microchemical analyses of otoliths taken from *O. mykiss*
21 in the San Joaquin River basin have verified that the anadromous form of this
22 species occurs in low numbers in the San Joaquin River basin (Zimmerman et al.
23 2009).

24 **9B.6.2.1 Historical Distribution**

25 *O. mykiss* once occurred throughout the Central Valley, spawning in the upper
26 reaches of tributaries to the Sacramento and San Joaquin rivers. Lindley et al.
27 (2006) conducted geographic information system (GIS) habitat modeling to
28 estimate the amount of suitable habitat to support *O. mykiss* populations in the
29 Central Valley, and their results suggest that steelhead were widely distributed
30 throughout the Sacramento River basin, but relatively less abundant in the
31 San Joaquin River basin due to natural barriers to migration. Yoshiyama et al.
32 (1996) conducted a review of historical sources to document the historical
33 distribution of Chinook Salmon in the Central Valley, which can be used to infer
34 historical distribution of steelhead. The assumption that steelhead distribution in
35 the Sacramento River basin overlapped with, and was likely more extensive than,
36 spring-run Chinook distribution under historical conditions has been supported by
37 studies conducted in the Klamath-Trinity River basin (Bureau of Indian Affairs
38 1985, Voight and Gale 1998). Yoshiyama et al. (1996) concluded that, because
39 steelhead upstream migration occurs during high flows, their leaping abilities are
40 superior to those of Chinook Salmon, and they have less restrictive spawning
41 gravel criteria. Steelhead in the Sacramento River basin “could have used at least
42 hundreds of miles of smaller tributaries not accessible to the earlier-spawning
43 salmon.” The model created by Lindley et al. (2006) estimates that 80 percent of
44 historically accessible habitat for Central Valley Steelhead is now behind

1 impassable dams; this estimate is supported by other research into steelhead and
2 Chinook Salmon habitat loss in the Central Valley (Clark 1929; Yoshiyama et al.
3 1996, 2001).

4 **9B.6.2.2 Current Distribution**

5 Steelhead distribution in Central Valley drainages has been greatly reduced
6 (McEwan and Jackson 1996). Steelhead are now primarily restricted to a few
7 remaining free-flowing tributaries and to stream reaches below large dams,
8 although a few steelhead may also spawn in intermittent streams during wet years.
9 Naturally spawning steelhead populations have been found in the upper
10 Sacramento River and tributaries below Keswick Dam; Mill, Deer, and Butte
11 creeks; and the Feather, Yuba, American, and Mokelumne rivers (CMARP 1998).
12 However, the records of naturally spawning populations depend on fish
13 monitoring programs. Recent implementation of monitoring programs has found
14 steelhead in additional streams, such as Auburn Ravine, Dry Creek, and the
15 Stanislaus River. It is possible that naturally spawning populations exist in many
16 other streams but are undetected because of the lack of monitoring or research
17 programs. Although impassable dams prevent resident Rainbow Trout from
18 emigrating, populations with steelhead ancestry may still exist above some dams
19 (Reclamation 2008).

20 In the Sacramento River basin, populations of *O. mykiss* are known to spawn in
21 the upper Sacramento, Yuba, Feather, and American rivers and in Deer, Mill, and
22 Butte creeks. Saeltzer Dam was removed from Clear Creek in 2000, granting
23 easier access to habitats in the higher-elevation canyon reaches. Though
24 improved access may have opened up suitable spawning and rearing habitat for
25 steelhead, it is not clear if steelhead have colonized Clear Creek since removal of
26 the dam. A summary of recent distribution information for steelhead in
27 Sacramento River tributaries in Good et al. (2005) shows that steelhead are
28 widespread in accessible streams, if not abundant.

29 Research and monitoring on steelhead are limited in comparison with Chinook
30 Salmon, so there is little specific information about the status and trend of the
31 species and how adults and juveniles use habitats in the mainstem river and the
32 Bay-Delta estuary. Though the upper reaches of the Sacramento River support a
33 spawning population of resident Rainbow Trout, the mainstem river habitat used
34 by the species is atypical for steelhead, which usually spawn in higher elevation,
35 steeper, and narrower channels. Management of the species is also complicated
36 by its polymorphism, with individuals being capable of exhibiting either a
37 resident (Rainbow Trout) or an anadromous (steelhead) life history.

38 **9B.6.3 Life History and Habitat Requirements**

39 Steelhead generally exhibit a more flexible life history strategy than Chinook
40 Salmon, and the habitat requirements of juvenile steelhead differ from those of
41 juvenile Chinook Salmon. Unlike Chinook Salmon, steelhead can be
42 iteroparous—that is, they can survive spawning, return to the ocean, and migrate
43 into fresh water to spawn again. Post-spawning adults are known as kelts. In
44 general, there are two types of steelhead: winter steelhead and summer steelhead.

1 Winter steelhead are of the ocean-maturing reproductive ecotype, becoming
2 sexually mature during their ocean phase and spawning soon after their arrival at
3 the spawning grounds. Adult summer steelhead are of the stream-maturing type,
4 which enter their natal streams and spend several months holding and maturing in
5 fresh water before spawning. Central Valley Steelhead are predominantly winter
6 steelhead, and this section describes the life history and habitat requirements of
7 winter steelhead.



8 Table 9B.11 illustrates aspects of the life-history timing of Central Valley
9 Steelhead.

Appendix 9B: Aquatic Species Life History Accounts

1 **Table 9B.11 Life-History Timing of Central Valley Steelhead**

Life Stage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Adult Upstream Migration ^a												
Spawning in Mainstem Sacramento River Downstream of Keswick Dam ^b				?								?
Incubation and Alevin Development ^c												
Fry Emergence ^c												
Age 0+ Outmigration from Upper Sacramento River ^b												
Age 1+ Outmigration through the Delta ^d												

- 2 Notes:
 3 a. Bailey 1954, Hallock et al. 1961, McEwan 2001
 4 b. Reclamation 2004
 5 c. Based on timing of spawning
 6 d. Based on fish facility salvage data (Reclamation 2004)

 Period of activity
 Period of peak activity

7

1 **9B.6.3.1 Adult Migration and Spawning**

2 Central Valley Steelhead generally leave the ocean and migrate upstream from
3 August through March (Busby et al. 1996). In the Sacramento River, steelhead
4 migrate upstream nearly every month of the year, with the bulk of migration from
5 August through November and the peak in late September (Bailey 1954, Hallock
6 et al. 1961, McEwan 2001). Spawning in the upper Sacramento River generally
7 occurs from December through April (Newton and Stafford 2011).

8 The majority of steelhead in the mainstem Sacramento River spawn downstream
9 of Keswick Dam (RM 302), with peak spawning from January through March
10 when water temperatures throughout much of the Sacramento River are suitable
11 to support egg incubation and emergence. The highest-density spawning within
12 the mainstem is likely in the upstream portion of this area near Redding; however,
13 the downstream extent of spawning is likely determined by the location of
14 suitable water temperatures to support summer rearing of 0+ juveniles, which lack
15 the swimming ability to move significant distances upstream to follow the
16 upstream retreat of cold water in summer. Most Sacramento River steelhead are
17 believed to spawn in the tributary streams. The progeny of adults that construct
18 redds downstream of locations with suitable water temperatures in summer likely
19 suffer high rates of mortality and contribute little to the population.

20 Steelhead migrate and spawn during high flows when observations and sampling
21 are difficult (McEwan 2001). They may have a spawning distribution similar to
22 late fall-run Chinook Salmon in that the juveniles of both species oversummer at
23 least once before outmigration, so redds must be located where summer water
24 temperatures can support summer rearing. The downstream extent of late fall-run
25 Chinook Salmon spawning is generally near Ball's Ferry Bridge (RM 276) in
26 most years. Steelhead generally have higher thermal tolerances than Chinook
27 Salmon (Moyle 2002), so steelhead spawning may extend slightly farther
28 downstream.

29 Under historical conditions, steelhead likely spawned in much higher-gradient
30 reaches in the Sacramento River and its tributaries, as do steelhead in other
31 portions of their range. Steelhead are common in reaches with gradients of less
32 than 6 percent (Burnett 2001, Harvey et al. 2002, Hicks and Hall 2003) and occur
33 in some systems in reaches of up to 12 percent and more (Engle 2002). Though
34 steelhead will spawn in mainstem river channels, it is unlikely that they spawned
35 in the reach of the mainstem Sacramento River below Keswick Dam where they
36 currently spawn because summer water temperatures in this reach were likely too
37 high to support oversummering by juveniles.

38 As with Chinook Salmon, steelhead spawn in areas with suitable gravel and
39 hydraulics. Work by Bovee (1978) found that steelhead prefer water depths of
40 14 inches (36 cm) for spawning, with a range between 6 and 24 inches (15 and
41 61 cm), and water velocities of 2 feet/second (61 cm/second), with a range of 1 to
42 3.6 feet/second (30 to 110 cm/second), which is similar to the hydraulic
43 conditions preferred by Chinook Salmon in the Central Valley. Steelhead
44 generally prefer to spawn in gravels, with optimal grain sizes ranging between

1 0.6 and 10 cm (6 and 102 mm) (Bjornn and Reiser 1991). For comparison, grain
2 sizes used by spawning Chinook range from a D₅₀ of 0.43 inch (10.8 mm) (Platts
3 et al. 1979) to a D₅₀ of 3.1 inches (78.0 mm) (Chambers et al. 1954, 1955).

4 Research in more northerly populations suggests that optimal spawning
5 temperatures range from 39 to 52°F (4 to 11°C), with egg mortality at water
6 temperatures above 56°F (13°C) (Hooper 1973, Bovee 1978, Reiser and Bjornn
7 1979, Bell 1986). More research is needed to understand the specific temperature
8 tolerances of steelhead in the Central Valley and southern portions of their range.
9 There is evidence that different strains of *O. mykiss* may have different thermal
10 tolerances at the egg and embryo stage (Myrick and Cech 2001).

11 As stated above, steelhead can survive spawning, return to the ocean, and migrate
12 into fresh water to spawn again. Although some kelts have been documented in
13 the Sacramento River, there are probably few repeat spawners in the Sacramento
14 River population (Reclamation 2004).

15 **9B.6.3.2 Fry and Juvenile Rearing**

16 Fry emergence is influenced by water temperature, but hatching generally
17 requires 4 weeks, with another 4 to 6 weeks in the gravels before emergence.
18 After emerging, steelhead fry typically disperse to shallow (<14 inches [36 cm]),
19 low-velocity near-shore areas such as stream margins and low-gradient riffles and
20 will forage in open areas lacking instream cover (Hartman 1965, Everest et al.
21 1986, Fontaine 1988). Everest and Chapman (1972) found that juvenile steelhead
22 of all sizes most often chose territories over large-sized substrates. As they
23 increase in size in late summer and fall, they increasingly use areas with cover
24 and show a preference for higher-velocity, deeper mid-channel areas near the
25 thalweg (Hartman 1965, Everest and Chapman 1972, Fontaine 1988). Bovee
26 (1978) reports that fry prefer water depths ranging between 10 inches (25 cm) and
27 20 inches (51 cm) and water temperatures ranging between 45°F (7°C) and 60°F
28 (16°C). Age 0+ steelhead have been relatively abundant in backwater pools and
29 often live in the downstream ends of pools in late summer (Bisson et al. 1988,
30 Fontaine 1988).

31 Steelhead fry may establish and defend territories soon after emerging
32 (Shapovalov and Taft 1954). Fry and juvenile steelhead that are unsuccessful in
33 establishing a territory may be displaced downstream where they may suffer
34 higher rates of mortality from predation, entrainment, or elevated water
35 temperatures (Dambacher 1991, Peven et al. 1994, Reedy 1995). Keeley (2001)
36 found that increased competition between juvenile steelhead, caused by higher
37 fish densities or lower food densities, caused increased mortality, lower or more
38 variable growth rates, and emigration of smaller fish. Downstream dispersal due
39 to overcrowding or high flows in rearing habitat does not necessarily increase
40 mortality where there is suitable habitat downstream (Kahler et al. 2001).
41 Downstream dispersal to larger stream reaches for further rearing prior to
42 smolting appears common in many systems (Bjornn 1978, Loch et al. 1985,
43 Leider et al. 1986, Dambacher 1991).

9B.6.3.3 Summer Rearing

Summer habitat can generally be assumed to be more limiting for age 1+ and 2+ juvenile steelhead than for age 0+ in many streams. Older age classes of juvenile steelhead (ages 1+ and 2+) prefer deeper water in summer than fry and show a stronger preference for pool habitats, especially deep pools near the thalweg with ample cover, as well as higher-velocity rapid and cascade habitats (Bisson et al. 1982, 1988; Dambacher 1991). Dambacher (1991) observed that most 1+ steelhead in the Steamboat Creek watershed of the North Umpqua River in Oregon were concentrated in mainstem reaches with relatively deep riffles and large substrates. Age 1+ fish typically feed in pools, especially scour and plunge pools (Fontaine 1988, Bisson et al. 1988). Age 1+ steelhead appear to avoid secondary channel and dammed pools, glides, and low-gradient riffles with mean depths less than 7.8 inches (20 cm) (Fontaine 1988, Bisson et al. 1988, Dambacher 1991). Beecher et al. (1993) reported that juvenile steelhead longer than 3 inches (75 mm) avoided areas less than 6 inches (15 cm) deep. Reedy (1995) indicates that age 1+ steelhead especially prefer high-velocity pool heads, where food resources are abundant, and pool tails, which provide optimal feeding conditions in summer due to lower energy expenditure requirements than the more turbulent pool heads. Fast, deep water, in addition to optimizing feeding versus energy expenditure, provides greater protection from avian and terrestrial predators (Everest and Chapman 1972).

9B.6.3.4 Winter Rearing

For juvenile steelhead to survive winter, they must avoid predation and high flows. The higher-gradient reaches typically used for spawning by steelhead (generally >3 percent) are often confined and characterized by coarse substrate that is immobile at all but the highest flows. Juvenile steelhead often use the interstitial spaces between cobbles and boulders as cover from high water velocity and presumably to avoid predation (Bjornn 1971, Hartman 1965, Bustard and Narver 1975, Swales et al. 1986, Everest et al. 1986, Grunbaum 1996). Age 0+ steelhead can use shallower habitats and can find interstitial cover in gravel-size substrates, while age 1+ or 2+ steelhead, because of their larger size, need coarser cobble/boulder substrate for cover (Bustard and Narver 1975; Bisson et al. 1982, 1988; Fontaine 1988; Dambacher 1991). Bustard and Narver (1975) reported that 1+ steelhead prefer water deeper than 17.5 inches (45 cm) in winter, while age 0+ steelhead often occupy water less than 5.8 inches (15 cm) deep and are rarely found at depths over about 23.4 inches (60 cm). In winter, age 1+ steelhead typically stay within the area of streambed that remains inundated at summer low flows, while age 0+ fish frequently overwinter beyond the summer low flow perimeter along the stream margins (Everest et al. 1986). Consequently, winter rearing habitat for age 1+ and 2+ juvenile steelhead is assumed to be more limiting than for age 0+ juveniles.

9B.6.3.5 Length of Stream Residence

Juvenile steelhead typically rear in fresh water from 1 to 3 years before outmigrating (McEwan and Jackson 1996). The majority of returning adult steelhead in the Central Valley have spent 2 years in fresh water before

1 emigrating to the ocean (McEwan 2001). A scale analysis conducted by Hallock
2 et al. (1961) indicated that 70 percent emigrated after 2 years, 29 percent after
3 1 year, and 1 percent after 3 years in fresh water. Juvenile emigration from the
4 upper Sacramento River occurs between November and late June, with a peak
5 between early January and late March (Reclamation 2004).

6 **9B.6.3.6 Bay-Delta Residence**

7 The Delta serves as an adult and juvenile migration corridor, connecting inland
8 habitat to the ocean. The Delta may also serve as a nursery area for juvenile
9 steelhead (McEwan and Jackson 1996); however, much is unknown regarding
10 historical and current role of the Delta as steelhead nursery habitat. In coastal
11 populations of winter steelhead, it is common for juvenile steelhead to migrate
12 downstream at age 1+ and rear in the estuary for an additional year before
13 smolting. Based on fish facility salvage data, most steelhead move through the
14 Delta from November through June, with the peak salvage during February,
15 March, and April. The majority of steelhead salvaged range from 175 to 325 mm,
16 with the most common size ranging from 226 to 250 mm. Some of the age 1+
17 steelhead captured in rotary screw traps at RBDD, GCID, and Knights Landing
18 may continue rearing for another year before entering the ocean. There may be
19 some areas of the Bay-Delta estuary where summer water temperatures are
20 moderated by tidal action so that steelhead 1+ migrants are able to rear throughout
21 summer (Reclamation 2008).

22 **9B.6.4 Population Trends**

23 Construction of large dams in the Central Valley had great impact on *O. mykiss*
24 populations because it eliminated access to nearly 80 percent of historical
25 spawning and rearing habitat (Lindley et al. 2006). Construction of Shasta and
26 Keswick dams eliminated access to many upstream tributaries (e.g., McCloud
27 River, Pit River, and Sacramento River) that provided the cold water temperatures
28 required for year-round rearing by steelhead. Dam construction also landlocked
29 potentially anadromous *O. mykiss* populations in the upper watershed, forcing
30 them to adopt a resident life history strategy (McEwan 2001).

31 In general, the majority of Central Valley Steelhead are confined to nonhistorical
32 spawning and rearing habitat below impassable dams, but the existing spawning
33 and rearing habitat can sustain steelhead at current population levels. In addition,
34 monitoring data indicate that much of the anadromous form of the species is
35 hatchery supported. Also, a strong resident component to the population
36 (Rainbow Trout) interacts with and produces both resident and anadromous
37 offspring.

38 In general, steelhead stocks throughout California have declined substantially.
39 McEwan and Jackson (1996) reported that the adult population of steelhead in
40 California was approximately 250,000, less than half the population that existed
41 in the 1960s (McEwan and Jackson 1996). In the Central Valley, approximately
42 1 to 2 million adult steelhead may have returned annually prior to 1850, as based
43 on historical Chinook Salmon abundance (McEwan 2001, NMFS 2006). In the
44 Sacramento River basin, the average run size of steelhead in the 1950s was

1 estimated to be approximately 20,540 adults (McEwan and Jackson 1996). In
2 contrast, escapement estimates in 1991 and 1992 were less than 10,000 adults,
3 less than half of the run size in the 1950s (McEwan and Jackson 1996). Similarly,
4 counts of wild steelhead at RBDD declined from an average annual run size of
5 12,900 in the late 1960s to 1,100 adults in the 1993–94 season (McEwan and
6 Jackson 1996). The most recent 5-year average for steelhead spawning upstream
7 of RBDD is less than 2,000 adults (Good et al. 2005). NMFS (2006) notes that
8 escapement estimates have not been made for the area upstream of RBDD since
9 the mid-1990s and that estimates of abundance are derived from extrapolation of
10 incidental catch of outmigrating juvenile steelhead captured as part of the
11 midwater-trawl sampling for juvenile Chinook Salmon at Chipps Island,
12 downstream of the confluence of the Sacramento and San Joaquin rivers.

13 Populations of naturally spawned Central Valley Steelhead have declined and are
14 composed predominantly of hatchery fish. The California Fish and Wildlife Plan
15 of 1965 estimated the combined annual run size for Central Valley and
16 San Francisco Bay tributaries to be about 40,000 during the 1950s (DFG 1965).
17 The spawning population during the mid-1960s for the Central Valley basin was
18 estimated at about 27,000 (DFG 1965). These numbers likely consisted of both
19 hatchery and wild steelhead. McEwan and Jackson (1996) estimated the annual
20 run size for the Central Valley basin to be less than 10,000 adults by the early
21 1990s. Much of the abundance data since the mid-1960s were obtained by visual
22 fish counts at the RBDD fish ladders when gates were closed during much of the
23 steelhead migration season. Current abundance estimates are not available for
24 naturally spawned fish since RBDD gate operations were changed, so the extent
25 to which populations have changed following the 1987–94 drought is unknown.
26 NMFS' (2003) status review estimated the Central Valley Steelhead population at
27 less than 3,000 adults.

28 **9B.6.5 Hatchery Influence**

29 Reclamation funds the operation of Coleman Hatchery, Livingston Stone
30 Hatchery, Nimbus Hatchery, and Trinity River Hatchery. DWR funds the
31 operation of the Feather River Hatchery. USFWS operates Coleman and
32 Livingston Stone hatcheries, and DFW operates Feather River, Nimbus, and
33 Trinity hatcheries. These hatcheries are operated to mitigate for the anadromous
34 salmonids that would be produced by the habitat if not for the dams on each
35 respective river. Reclamation and DWR have discretion over how the hatcheries
36 are operated, but generally leave operational decisions on how to meet mitigation
37 goals to the operating agency (Reclamation 2008).

38 Hatchery production of steelhead is large compared to natural production, based
39 on the Chipps Island trawl data (Good et al. 2005). The bulk of hatchery releases
40 in the Central Valley occurs in the Sacramento River basin. An analysis of
41 steelhead captures from trawl data by Nobriga and Cadrett (2001) indicated that
42 hatchery steelhead composed 63 to 77 percent of the steelhead catch. Steelhead
43 stocks at the Mokelumne River Hatchery and Nimbus Hatchery on the American
44 River are not part of the Central Valley Steelhead DPS because of the source of
45 broodstock used and genetic similarities to Eel River stocks (Good et al. 2005).

1 Genetic analysis indicated steelhead from the American River (collected from
2 both the Nimbus Hatchery and the American River) are genetically more similar
3 to Eel River steelhead (Northern California ESU) than other Central Valley
4 Steelhead stocks. Eel River steelhead were used to found the Nimbus Hatchery
5 stock. Mokelumne River Rainbow Trout (hatchery produced and naturally
6 spawned) are genetically most similar to Mount Shasta Hatchery trout, but also
7 show genetic similarity to the Northern California ESU (Nielsen 1997). Nielsen
8 et al. (2005) found American River steelhead to be genetically different from
9 other Central Valley stocks.

10 **9B.6.6 References**

- 11 Bailey, E .D. 1954. *Time pattern of 1953-54 migration of salmon and steelhead*
12 *into the upper Sacramento River*. Unpublished report. California
13 Department of Fish and Game. As cited in McEwan 2006.
- 14 Beecher, H. A., T. H. Johnson, and J. P. Carleton. 1993. Predicting
15 microdistributions of steelhead (*Oncorhynchus mykiss*) parr from depth
16 and velocity preference criteria: test of an assumption of the Instream
17 Flow Incremental Methodology. *Canadian Journal of Fisheries and*
18 *Aquatic Sciences* 50: 2380–2387.
- 19 Bell, M. C., editor. 1986. *Fisheries handbook of engineering requirements and*
20 *biological criteria*. NTIS AD/A167-877. Fisheries-Engineering Research
21 Program, U.S. Army Corps of Engineers, North Pacific Division. Portland,
22 Oregon.
- 23 Bisson, P., J. L. Nielsen, R. A. Palmason, and L. E. Grove. 1982. *A system of*
24 *naming habitat types in small streams, with examples of habitat utilization*
25 *by salmonids during low streamflows*. Proceedings of the symposium on
26 acquisition and utilization of aquatic habitat inventory information. Edited
27 by N. B. Armantrout, 62–73. American Fisheries Society, Western
28 Division. Bethesda, Maryland.
- 29 Bisson, P. A., K. Sullivan, and J. L. Nielsen. 1988. Channel hydraulics, habitat
30 use, and body form of juvenile coho salmon, steelhead trout, and cutthroat
31 trout in streams. *Transactions of the American Fisheries Society* 117: 262–
32 273.
- 33 Bjornn, T. C. 1971. Trout and salmon movements in two Idaho streams as related
34 to temperature, food, stream flow, cover, and population density.
35 *Transactions of the American Fisheries Society* 100: 423–438.
- 36 _____. 1978. *Survival, production, and yield of trout and Chinook salmon in the*
37 *Lemhi River, Idaho, Bulletin No. 27*. Prepared by Idaho Cooperative
38 Fishery Research Unit, College of Forestry, Wildlife and Range Sciences,
39 University of Idaho, Moscow, for Idaho Department of Fish and Game.

- 1 Bjornn, T. C., and D. W. Reiser. 1991. *Habitat requirements of salmonids in*
2 *streams*. Influences of forest and rangeland management on salmonid
3 fishes and their habitats. Edited W. R. Meehan, 83-138. American
4 Fisheries Society Special Publication No. 19.
- 5 Bovee, K. D. 1978. *Probability-of-use-criteria for the family Salmonidae*.
6 Instream Flow Information Paper 4. FWS/OBS-78/07. U.S. Fish and
7 Wildlife Service.
- 8 Bureau of Indian Affairs. 1985. *Klamath River basin fisheries resource plan*. U.S.
9 Department of the Interior. Prepared by CH2M HILL, Redding,
10 California. As cited by Klamath River Basin Fisheries Task Force *Long*
11 *Range Plan for the Klamath River Basin Conservation Area Fishery*
12 *Restoration Program*, January 1991.
- 13 Burnett, K. M. 2001. *Relationships among juvenile anadromous salmonids, their*
14 *freshwater habitat, and landscape characteristics over multiple years and*
15 *spatial scales in the Elk River, Oregon*. Doctoral dissertation. Oregon
16 State University, Corvallis.
- 17 Busby, P. J., T. C. Wainwright, G. J. Bryant, L. Lierheimer, R. S. Waples, F. W.
18 Waknitz, and I. V. Lagomarsino. 1996. Status Review of West Coast
19 steelhead from Washington, Idaho, Oregon and California. U.S.
20 Department of Commerce. NOAA Technical Memo. NMFS-NWFSC-27.
- 21 Bustard, D. R., and D. W. Narver. 1975. Aspects of the winter ecology of juvenile
22 coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo*
23 *gairdneri*). *Journal of the Fisheries Research Board of Canada* 32: 667–
24 680.
- 25 Chambers, J. S., G. H. Allen, and R. T. Pressey. 1955. *Research relating to study*
26 *of spawning grounds in natural areas*. Annual report, Contract DA 35026.
27 Washington Department of Fisheries, Olympia.
- 28 Clark, G. H. 1929. Sacramento River salmon fishery. *California Fish and Game*
29 15: 1-11.
- 30 CMARP (Comprehensive Monitoring, Assessment and Research Program for the
31 CALFED Bay-Delta Program). 1999. *Monitoring, assessment, and*
32 *research on Central Valley steelhead: status of knowledge, review of*
33 *existing programs, and assessment of needs*. Draft Report.
- 34 Cramer, S. P., D. W. Alley, J. E. Baldrige, K. Barnard, D. B. Demko, D. H.
35 Dettman, B. Farrell, J. Hagar, T. P. Keegan, A. Laird, W. T. Mitchell, R.
36 C. Nuzum, R. Orton, J. J. Smith, T. L. Taylor, P. A. Unger, and E. S. Van
37 Dyke. 1995. *The status of steelhead populations in California in regards*
38 *to the Endangered Species Act*. Special report. Submitted to National
39 Marine Fisheries Service on behalf of Association of California Water
40 Agencies, S.P. Cramer & Associates, Gresham, Oregon.

- 1 Dambacher, J. M. 1991. *Distribution, abundance, and emigration of juvenile*
2 *steelhead (Oncorhynchus mykiss), and analysis of stream habitat in the*
3 *Steamboat Creek basin, Oregon.* Master's thesis. Oregon State University,
4 Corvallis.
- 5 DFG (California Department of Fish and Game). 1965. *California fish and*
6 *wildlife plan.* California Department of Fish and Game, Sacramento.
- 7 Engle, R. O. 2002. *Distribution and summer survival of juvenile steelhead trout*
8 *(Oncorhynchus mykiss) in two streams within King Range National*
9 *Conservation Area, California.* Master's thesis, Humboldt State
10 University, Arcata, California.
- 11 Everest, F. H., and D. W. Chapman. 1972. Habitat selection and spatial
12 interaction by juvenile Chinook salmon and steelhead trout in two Idaho
13 streams. *Journal of the Fisheries Research Board of Canada* 29: 91–100.
- 14 Everest, F. H., G. H. Reeves, J. R. Sedell, J. Wolfe, D. Hohler, and D. A. Heller.
15 1986. *Abundance, behavior, and habitat utilization by coho salmon and*
16 *steelhead trout in Fish Creek, Oregon, as influenced by habitat*
17 *enhancement.* Annual report, 1985 Project No. 84-11. Prepared by U.S.
18 Forest Service for Bonneville Power Administration, Portland, Oregon.
- 19 Fontaine, B. L. 1988. *An evaluation of the effectiveness of instream structures*
20 *for steelhead trout rearing habitat in the Steamboat Creek basin.*
21 *Master's thesis.* Oregon State University, Corvallis.
- 22 Good, T. P., R. S. Waples, and P. Adams. 2005. *Updated status of federally*
23 *listed ESUs of west coast salmon and steelhead.* NOAA Technical
24 Memorandum NMFSNWFSC-66. National Marine Fisheries Service,
25 Seattle, Washington.
- 26 Grunbaum, J. B. 1996. *Geographical and seasonal variation in diel habitat use*
27 *by juvenile (age 1+) steelhead trout (Oncorhynchus mykiss) in Oregon*
28 *coastal and inland streams.* Master's thesis. Oregon State University,
29 Corvallis.
- 30 Hallock, R. J., W. F. Van Woert, and L. Shapovalov. 1961. An evaluation of
31 stocking hatchery-reared steelhead rainbow trout (*Salmo gairdnerii*
32 *gairdnerii*) in the Sacramento River system. California Department of
33 Fish and Game. *Fish Bulletin* 114.
- 34 Hartman, G. F. 1965. The role of behavior in the ecology and interaction of
35 underyearling coho salmon (*Oncorhynchus kisutch*) and steelhead trout
36 (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada*
37 22: 1035–1081.
- 38 Harvey, B. C., J. L. White, and R. J. Nakamoto. 2002. Habitat relationships and
39 larval drift of native and nonindigenous fishes in neighboring tributaries of
40 a coastal California river. *Transactions of the American Fisheries Society*
41 131:159–170.

- 1 Hicks, B. J., and J. D. Hall. 2003. Rock type and channel gradient structure
2 salmonid populations in the Oregon Coast Range. *Transactions of the*
3 *American Fisheries Society* 132: 468–482.
- 4 Hooper, D. R. 1973. *Evaluation of the effects of flows on trout stream ecology.*
5 Pacific Gas and Electric Company, Emeryville, California.
- 6 Kahler, T. H., P. Roni, and T. P. Quinn. 2001. Summer movement and growth of
7 juvenile anadromous salmonids in small western Washington streams.
8 *Canadian Journal of Fisheries and Aquatic Sciences* 58: 1947-2637.
- 9 Keeley, E. R. 2001. Demographic responses to food and space competition by
10 juvenile steelhead trout. *Ecology* 82: 1247-1259.
- 11 Leider, S. A., M. W. Chilcote, and J. J. Loch. 1986. Comparative life history
12 characteristics of hatchery and wild steelhead trout (*Salmo gairdneri*) of
13 summer and winter races in the Kalama River, Washington. *Canadian*
14 *Journal of Fisheries and Aquatic Sciences* 43: 1398–1409.
- 15 Lindley, S. T., R. S. Schick, A. Agrawal, M. Goslin, T. E. Pearson, E. Mora, J. J.
16 Anderson, B. May, S. Greene, C. Hanson, A. Low, D. McEwan, R. B.
17 MacFarlane, C. Swanson, and J. G. Williams. 2006. Historical population
18 structure of Central Valley steelhead and its alteration by dams. *San*
19 *Francisco Estuary and Watershed Science* 4: 1-19.
- 20 Loch, J. J., M. W. Chilcote, and S. A. Leider. 1985. *Kalama River studies final*
21 *report: Part II. Juvenile downstream migrant studies.* Washington
22 Department of Game, Fisheries Management Division, Olympia.
- 23 McEwan, D. 2001. *Central Valley steelhead.* Contributions to the biology of
24 Central Valley salmonids. Edited by R. L. Brown, 1-44. *Fish Bulletin* 179.
25 California Department of Fish and Game, Sacramento.
- 26 McEwan, D., and T. A. Jackson. 1996. *Steelhead restoration and management*
27 *plan for California.* California Department of Fish and Game, Inland
28 Fisheries Division, Sacramento.
- 29 Moyle, P. B. 2002. *Inland fishes of California.* Revised edition. University of
30 California Press, Berkeley.
- 31 Myrick, C. A., and J. J. Cech, Jr. 2001. *Temperature effects on Chinook salmon*
32 *and steelhead: a review focusing on California's Central Valley*
33 *populations.* Technical Publication 01-1. Bay-Delta Modeling Forum.
- 34 Newton, J. M. and L. A. Stafford. 2011. Monitoring Adult Chinook Salmon,
35 Rainbow Trout, and Steelhead in Battle Creek, California, from March
36 through November 2009. Red Bluff, CA: U.S. Fish and Wildlife Service.
- 37 Nielsen, J. L. 1997. *Genetic variation in Mokelumne River trout (Oncorhynchus*
38 *mykiss) using mitochondrial DNA and ten nuclear microsatellite loci.*
39 Revised technical report. Prepared for East Bay Municipal Utility District,
40 Oakland, California.

- 1 Nielsen, J. L., S. Paver, T. Wiacek, and I. Williams. 2005. *Genetics of Central*
2 *Valley O. mykiss populations: drainage and watershed scale analyses.*
3 San Francisco Estuary and Watershed Science. As cited by West Coast
4 Steelhead Biological Review Team *Status Review Update for Deferred*
5 *and Candidate ESU's of West Coast Steelhead*, December 19, 1997.
- 6 NMFS (National Marine Fisheries Service). 1998. Endangered and threatened
7 species; threatened status for two ESUs of steelhead in Washington,
8 Oregon, and California. *Federal Register* 63: 13347–13371.
- 9 _____. 2003. *Updated status of federally listed ESUs of West Coast salmon and*
10 *steelhead.* National Marine Fisheries Service. Northwest and Southwest
11 Fisheries Science Centers.
- 12 _____. 2006. Endangered and threatened species; final listing determinations for
13 10 Distinct Population Segments of West Coast steelhead. *Federal*
14 *Register* 71: 834-862.
- 15 Nobriga, M. L. and P. Cadrett. 2001. Differences among hatchery and wild
16 steelhead: evidence from delta fish monitoring programs. *Interagency*
17 *Ecological Program Newsletter* 30-38.
- 18 Peven, C. M., R. R. Whitney, and K. R. Williams. 1994. Age and length of
19 steelhead smolts from the mid-Columbia River basin, Washington. *North*
20 *American Journal of Fisheries Management* 14: 77–86.
- 21 Platts, W. S., M. A. Shirazi, and D. H. Lewis. 1979. *Sediment particle sizes used*
22 *by salmon for spawning with methods for evaluation.* Ecological Research
23 Series EPA-600/3-79-043. U.S. Environmental Protection Agency,
24 Corvallis Environmental Research Laboratory, Corvallis, Oregon.
- 25 Reclamation (Bureau of Reclamation). 2004. *Long-term Central Valley Project*
26 *and State Water Project operations criteria and plan.* Biological
27 Assessment. Bureau of Reclamation, Sacramento, California.
- 28 _____. 2008. *Biological assessment on the continued long-term operations of the*
29 *Central Valley Project and the State Water Project.* Bureau of
30 Reclamation, Sacramento, California.
- 31 Reedy, G. D. 1995. *Summer abundance and distribution of juvenile Chinook*
32 *salmon (Oncorhynchus tshawytscha) and steelhead trout (Oncorhynchus*
33 *mykiss) in the Middle Fork Smith River, California.* Master's thesis.
34 Humboldt State University, Arcata, California.
- 35 Reiser, D. W., and T. C. Bjornn. 1979. *Habitat requirements of anadromous*
36 *salmonids.* Influence of forest and rangeland management on anadromous
37 fish habitat in western North America. Edited by W. R. Meehan, 1-54.
38 General Technical Report PNW-96. U.S. Forest Service, Pacific
39 Northwest Forest and Range Experiment Station. Portland, Oregon.
- 40 Shapovalov, L., and A. C. Taft. 1954. The life histories of the steelhead rainbow
41 trout (*Salmo gairdneri gairdneri*) and silver salmon (*Oncorhynchus*
42 *kisutch*) with special reference to Waddell Creek, California, and

- 1 recommendations regarding their management. California Department of
2 Fish and Game. *Fish Bulletin* 98.
- 3 Swales, S., R. B. Lauzier, and C. D. Levings. 1986. Winter habitat preferences
4 of juvenile salmonids in two interior rivers in British Columbia. *Canadian*
5 *Journal of Zoology* 64: 1506–1514.
- 6 Voight, H. N., and D. B. Gale. 1998. *Distribution of fish species in tributaries of*
7 *the lower Klamath River: an interim report, FY 1996*. Technical report,
8 No. 3. Yurok Tribal Fisheries Program, Habitat Assessment and
9 Biological Monitoring Division.
- 10 Yoshiyama, R. M., E. R. Gerstung, F. W. Fisher, and P. B. Moyle. 1996.
11 Historical and present distribution of Chinook salmon in the Central
12 Valley drainage of California, Sierra Nevada Ecosystem Project. Final
13 report to congress. Volume III: assessments, commissioned reports, and
14 background information, 309–362. University of California, Center for
15 Water and Wildland Resources, Davis.
- 16 _____. 2001. Historical and present distribution of Chinook salmon in the
17 Central Valley drainage of California. Contributions to the biology of
18 Central Valley salmonids. Edited by R. L. Brown, 71-176. *Fish Bulletin*
19 179. California Department of Fish and Game, Sacramento.
- 20 Zimmerman, C. E., G. W. Edwards, and K. Perry. 2009. Maternal origin and
21 migratory history of steelhead and rainbow trout captured in rivers of the
22 Central Valley, California. *Transactions of the American Fisheries*
23 *Society* 138: 280-291.

24 **9B.7 Klamath Mountains Province Steelhead** 25 **(*Oncorhynchus mykiss*)**

26 **9B.7.1 Legal Status**

27 Federal: Not warranted

28 State: Species of Special Concern

29 A status review in 2001 (NMFS 2001) concluded that the Klamath Mountains
30 Province Steelhead DPS was not in danger of extinction or likely to become so in
31 the foreseeable future; therefore, it was not warranted for listing as threatened or
32 endangered. This conclusion was based on population estimates and a finding
33 that the genetic risk from naturally spawning hatchery fish was lower than
34 estimated in previous reviews, as well as consideration of ongoing and proposed
35 conservation efforts for anadromous salmonids in the basin (NMFS 2001).

36 The Klamath Mountains Province Steelhead DPS contains both summer and
37 winter runs. Moyle (2002) describes steelhead in the Klamath Basin as having a
38 summer run and a winter run. Some divide the winter run into fall and winter
39 runs (Barnhart 1994, Hopelain 1998, USFWS 1998, Papa et al. 2007). In this
40 section, winter steelhead refers to steelhead returning from fall through winter,

1 except in cases when the distinction is pertinent to the discussion. The following
2 summary focuses on steelhead in the Trinity River, which is within the area
3 potentially affected by the proposed action, and on the mainstem Klamath in
4 terms of potential effects on its role as a migration corridor for the steelhead runs.

5 **9B.7.2 Distribution**

6 Based on escapement data, approximately 55 percent of the summer run spawn in
7 the Trinity River and other lower-elevation tributaries to the Klamath River. The
8 Trinity, Scott, Shasta, and Salmon rivers are important spawning streams for the
9 winter run.

10 Historically, steelhead probably ascended Clear Creek past the French Gulch area,
11 but access to the upper basin was blocked by Whiskeytown Dam in 1964
12 (Yoshiyama et al. 1996). Operation of Whiskeytown Dam can produce suitable
13 cold-water habitat downstream to Placer Road Bridge depending on flow releases
14 (DFG 1998). McCormick-Saeltzer Dam, which limited steelhead migrations
15 through ineffective fish ladders, was removed in 2000, allowing steelhead
16 potential access to good habitat up to Whiskeytown Dam. USFWS has conducted
17 snorkel surveys targeting spring-run Chinook (May through September) since
18 1999. Steelhead/rainbow are enumerated and separated into small, medium, and
19 large (>22 inches) during these surveys, but because the majority of the steelhead
20 run is unsurveyed, no spawner abundance estimates have been attempted
21 (Reclamation 2008). Redd counts conducted during the 2001-02 run found that
22 most spawning occurred upstream, near Whiskeytown Dam. Because of the large
23 resident rainbow population, no steelhead population estimate could be made
24 (Reclamation 2008). A remnant “landlocked” population of Rainbow Trout with
25 steelhead ancestry may exist in Clear Creek above Whiskeytown Dam
26 (Reclamation 2008).

27 **9B.7.3 Life History and Habitat Requirements**

28 General habitat requirements for steelhead are described in the Central Valley
29 Steelhead profile; the following describes life history strategies and habitat
30 requirements unique to steelhead of the Upper Klamath Mountains Province DPS
31 or of primary importance to its life history. Both winter and summer runs of
32 steelhead are included in the DPS. Winter steelhead become sexually mature
33 during their ocean phase and spawn soon after arriving at their spawning grounds.
34 Adult summer steelhead enter their natal streams and spend several months
35 holding and maturing in fresh water before spawning. Throughout the entire year,
36 at least one of the diverse life stages can be found present in the river (Israel
37 2003). As with the Central Valley DPS, this DPS is composed predominantly of
38 winter steelhead.

39 **9B.7.3.1 Winter Run**

40 Winter steelhead adults generally enter the Klamath River from July through
41 October (fall run) and from November through March (winter run) (USFWS
42 1998). Winter steelhead primarily spawn in tributaries from January through
43 April (USFWS 1998), with peak spawn timing in February and March (ranging

1 from January to April) (NRC 2004). Adults may repeat spawning in subsequent
2 years after returning to the ocean. Half-pounders typically use the mainstem
3 Klamath River until leaving the following March (NRC 2004), although they also
4 use larger tributaries such as the Trinity River (Dean 1994, 1995).

5 Fry emerge in spring (NRC 2004), with fry observed in outmigrant traps in Bogus
6 Creek and Shasta River from March through mid-June (Dean 1994). Age-0+ and
7 1+ juveniles have been captured in outmigrant traps in spring and summer in
8 tributaries to the Klamath River above Seiad Creek (DFG 1990a, 1990b). These
9 fish are likely rearing in the mainstem or non-natal tributaries before leaving as
10 age-2+ outmigrants.

11 Juvenile outmigration primarily occurs between May and September with peaks
12 between April and June, although smolts are captured in the estuary as early as
13 March and as late as October (Wallace 2004). Most adult returns (86 percent)
14 originate from fish that smolt at age 2+, in comparison with only 10 percent for
15 age-1 juveniles and 4 percent for age 3+ juveniles (Hopelain 1998).

16 Similar limiting factors listed for summer steelhead also affect winter steelhead
17 populations, including degraded habitats, decreased habitat access, fish passage,
18 predation, and competition (for more species information see USFWS 1998, NRC
19 2004, and Wallace 2004).

20 **9B.7.3.2 Summer Run**

21 Summer steelhead adults enter and migrate up the Klamath River from March
22 through June while sexually immature (Hopelain 1998), then hold in cooler
23 tributary habitat until spawning begins in December (USFWS 1998).

24 Juvenile summer steelhead in the Klamath Basin may rear in fresh water for up to
25 3 years before outmigrating. Although many juveniles migrate downstream at age
26 1+ (Scheiff et al. 2001), those that outmigrate to the ocean at age 2+ appear to
27 have the highest survival (Hopelain 1998). Juveniles outmigrating from
28 tributaries at age 0+ and age 1+ may rear in the mainstem or in non-natal
29 tributaries (particularly during periods of poor water quality) for 1 or more years
30 before reaching an appropriate size for smolting. Age-0 juvenile steelhead have
31 been observed migrating upstream into tributaries, off-channel ponds, and other
32 winter refuge habitat in the lower Klamath River. Juvenile outmigration can
33 occur from spring through fall. Smolts are captured in the mainstem and estuary
34 throughout fall and winter (Wallace 2004), but peak smolt outmigration normally
35 occurs from April through June, based on estuary captures (Wallace 2004).

36 Temperatures in the mainstem are generally suitable for juvenile steelhead, except
37 during summer, especially upstream of Seiad Valley.

38 **9B.7.4 Population Trends**

39 Long-term data are not available to evaluate Klamath River steelhead population
40 trends. DFG (1965) estimated a basinwide annual run size of 283,000 adult
41 steelhead (spawning escapement + harvest). Busby et al. (1994) reported winter
42 steelhead runs in the basin to be 222,000 during the 1960s. Steelhead spawning
43 surveys on tributaries to the mainstem Trinity River were conducted in 1964,

1 1971, 1972, and 1974 to monitor the effect of Lewiston Dam on steelhead
 2 populations. Hopelain (2001) used creel and gill net harvest data to estimate the
 3 winter-run steelhead population at 10,000 to 30,000 adults annually in the early
 4 1980s. Spawning surveys were also conducted in South Fork Trinity River
 5 tributaries from 1989 to 1995 under DFW's Trinity River Project (Garrison 2000).
 6 Population estimates of summer steelhead showed a steep decline during the
 7 1990s (Reclamation 2008), but Koch (2001) reported increasing runs on the
 8 Klamath and Trinity rivers following the late 1990s.

9 **9B.7.5 Hatchery Influence**

10 Reclamation funds the operation of Coleman Hatchery, Livingston Stone
 11 Hatchery, Nimbus Hatchery, and Trinity River Hatchery. DWR funds the
 12 operation of the Feather River Hatchery. USFWS operates Coleman and
 13 Livingston Stone hatcheries, and DFW operates Feather River, Nimbus, and
 14 Trinity hatcheries. These hatcheries are operated to mitigate for the anadromous
 15 salmonids that would be produced by the habitat if not for the dams on each
 16 respective river. Reclamation and DWR have discretion over how the hatcheries
 17 are operated, but generally leave operational decisions on how to meet mitigation
 18 goals to the operating agency (Reclamation 2008).

19 NMFS (2001) reported that the Trinity River population is thought to contain a
 20 large percentage of hatchery origin spawners of mostly fall-run fish
 21 (20-70 percent).

22 **9B.7.6 References**

23 Barnhart, R. A. 1994. *Salmon and steelhead populations of the Klamath-Trinity*
 24 *Basin, California*. Klamath Basin fisheries symposium. Edited by T. J.
 25 Hassler, 73-97. California Cooperative Fishery Research Unit, Humboldt
 26 State University, Arcata. As cited by National Marine Fisheries Service
 27 *Biological Opinion for Klamath Project Operations*, May 31, 2002.

28 Busby P. J., T. C. Wainwright, and R. S. Waples. 1994. *Status review for Klamath*
 29 *Mountains Province steelhead*. NOAA Technical Memorandum NMFS-
 30 NWFSC-19. National Marine Fisheries Service, Seattle, Washington.

31 Dean, M. 1994. *Life history, distribution, run size, and harvest of spring-run*
 32 *Chinook salmon in the south fork Trinity River Basin*. Chapter VII - job
 33 VII in Trinity River Basin monitoring project 1991-1992.

34 _____. 1995. *Life history, distribution, run size, and harvest of spring-run*
 35 *Chinook salmon in the south fork Trinity River Basin*. Chapter VII - job
 36 VII in Trinity River Basin monitoring project 1992-1993.

37 DFG (California Department of Fish and Game). 1965. *California fish and*
 38 *wildlife plan*. California Department of Fish and Game, Sacramento. As
 39 cited by U.S. Fish and Wildlife Service *Klamath River (Iron Gate Dam to*
 40 *Seiad Creek) Life Stage Periodicities for Chinook, Coho, and Steelhead*,
 41 July 1997.

- 1 _____ . 1990a. *Juvenile salmonid sampling within the Klamath-Trinity Basin,*
2 *1984.* Draft report. Inland Fisheries Division, Arcata, California. As cited
3 by U.S. Fish and Wildlife Service *Klamath River (Iron Gate Dam to Seiad*
4 *Creek) Life Stage Periodicities for Chinook, Coho, and Steelhead,* July
5 1997.
- 6 _____ . 1990b. *Distribution, abundance, fork length and coded-wire tag recovery*
7 *data for juvenile anadromous salmonids within the Klamath-Trinity Basin,*
8 *1985.* Draft report. Inland Fisheries Division, Arcata, California.
- 9 _____ . 1998. *Strategic plan for management of Klamath Mountains Province*
10 *steelhead trout.* Prepared for the National Marine Fisheries Service by the
11 Resources Agency.
- 12 Garrison, P. 2000. *Study 2d1 – Steelhead spawner surveys in Trinity River*
13 *tributaries.* California Department of Fish and Game [?], Steelhead
14 Research and Monitoring Program, Weaverville Remote Office.
- 15 Hopelain J. S. 1998. *Age, growth, and life history of Klamath River basin*
16 *steelhead trout (Oncorhynchus mykiss irideus) as determined from scale*
17 *analysis.* Inland Fisheries Administration Report 98-3. California
18 Department of Fish and Game, Sacramento.
- 19 _____ . 2001. *Lower Klamath River angler creel census with emphasis on*
20 *upstream migrating fall Chinook salmon, coho salmon, and steelhead*
21 *trout during July through October, 1983 through 1987.* Inland Fisheries
22 Administrative Report 01-1. California Department of Fish and Game,
23 Sacramento.
- 24 Israel, J. 2003. *Life history, ecology, and status of Klamath River steelhead.*
25 Report to University of California, Davis, Center for Watershed Sciences.
- 26 Koch, D. B. 2001. Letter from CDFG to J. Blum, National Marine Fisheries
27 Service, 16 February.
- 28 Moyle P. B. 2002. *Inland fishes of California (second edition).* University of
29 California Press, Berkeley.
- 30 NMFS (National Marine Fisheries Service). 2001. Endangered and threatened
31 species: final listing determination for Klamath Mountains Province
32 steelhead. Federal Register 66:17845-17856.
- 33 NRC (National Research Council). 2004. *Endangered and threatened fishes in the*
34 *Klamath River basin: causes of decline and strategies for recovery.* The
35 National Academies Press, Washington, D.C. Available at:
36 <http://www.nap.edu/openbook.php?isbn=0309090970>.
- 37 Papa R., J. A. Israel, F. Nonnis Marzano, and B. May. 2007. Assessment of
38 genetic variation between reproductive ecotypes of Klamath River
39 steelhead reveals differentiation associated with different run-timings.
40 *Journal of Applied Ichthyology* 23: 142-146.

- 1 Reclamation (Bureau of Reclamation). 2008. *Biological assessment on the*
2 *continued long-term operations of the Central Valley Project and the State*
3 *Water Project*. U.S. Bureau of Reclamation, Sacramento, California.
- 4 Scheiff A. J., J. S. Lang, and W. D. Pinnix. 2001. *Juvenile salmonid monitoring*
5 *on the mainstem Klamath River at Big Bar and mainstem Trinity River at*
6 *Willow Creek 1997-2000*. Annual report of the Klamath River Fisheries
7 Assessment Program. U.S. Fish and Wildlife Service, Arcata Fish and
8 Wildlife Office, Arcata, California. [Juvenile salmonid monitoring annual](#)
9 [report 2001](#)
- 10 USFWS (U.S. Fish and Wildlife Service). 1997. *Klamath River (Iron Gate Dam*
11 *to Seiad Creek) life state periodicities for Chinook, coho, and steelhead*.
12 Prepared by USFWS, Coastal California Fish and Wildlife Office, Arcata.
- 13 Wallace, M. 2004. *Natural vs. hatchery proportions of juvenile salmonids*
14 *migrating through the Klamath River estuary and monitor natural and*
15 *hatchery juvenile salmonid emigration from the Klamath River Basin. July*
16 *1, 1998 through June 30, 2003*. Final performance report. Federal Aid in
17 Sport Fish Restoration Act. Project no. F-51-R-6. Arcata, California.
- 18 Yoshiyama, R. M., E. R. Gerstung, F. W. Fisher, and P. B. Moyle. 1996.
19 *Historical and present distribution of Chinook salmon in the Central*
20 *Valley drainage of California*. Volume III: assessments, commissioned
21 reports, and background information. Sierra Nevada Ecosystem Project:
22 final report to Congress, 309-361. University of California, Centers for
23 Water and Wildlife Resources, Davis.

24 **9B.8 Southern Oregon/Northern California Coast** 25 **Coho Salmon ESU (*Oncorhynchus kisutch*)**

26 **9B.8.1 Legal Status**

27 Federal: Threatened

28 State: Threatened

29 Coho Salmon (*Oncorhynchus kisutch*) in the Trinity River are in the Southern
30 Oregon/Northern California Coast Coho Salmon ESU and were listed as
31 threatened under the ESA in 1997 (NMFS 1997) and threatened under the
32 California Endangered Species Act in 2002. This ESU includes naturally
33 spawning populations between Punta Gorda, California, and Cape Blanco,
34 Oregon, which encompasses the Trinity and Klamath basins (NMFS 1997).
35 Three artificial propagation programs are considered to be part of the ESU: the
36 Cole Rivers Hatchery, Trinity River Hatchery, and Iron Gate Hatchery Coho
37 Salmon programs. NMFS has determined that these artificially propagated stocks
38 are no more than moderately diverged from the local natural populations. In
39 addition, Coho Salmon in the Klamath Basin have been listed by the California
40 Fish and Game Commission as threatened under the California Endangered
41 Species Act (DFG 2002).

9B.8.2 Life History and Habitat Requirements

1 Coho Salmon exhibit a 3-year life cycle in the Trinity River and depend on
2 freshwater habitat conditions year-round because they spend a full year residing
3 in fresh water. Most Coho Salmon enter rivers between August and January, with
4 some more northerly populations entering as early as June. Coho Salmon river
5 entry timing is influenced by such factors as genetics, stage of maturity, river
6 discharge, and access past the river mouth. Spawning is concentrated in riffles or
7 in gravel deposits at the downstream end of pools with suitable water depth,
8 velocity, and substrate size. Spawning in the Trinity River occurs mostly in
9 November and December. Coho eggs incubate from 35 to more than 100 days
10 depending on water temperature and emerge from the gravel 2 to 7 weeks after
11 hatching. Coho eggs hatch after an accumulation of 400 to 500 temperature units
12 measured in degrees Celsius and emerge from the gravel after 700 to
13 800 temperature units. After emergence, fry move into areas out of the main
14 current. As Coho grow, they spread out from the areas where they were spawned.
15 During summer, juvenile Coho prefer pools and riffles with adequate cover such
16 as large woody debris with smaller branches, undercut banks, and overhanging
17 vegetation and roots.
18

19 Juvenile Coho Salmon overwinter in large mainstem pools, beaver ponds,
20 backwater areas, and off-channel pools with cover such as woody debris and
21 undercut banks. Most juvenile Coho Salmon spend a year in fresh water, with
22 northerly populations spending 2 full years in fresh water. Coho in the Trinity
23 River are thought to be exclusively 3-year-life-cycle fish (1 year in fresh water).
24 Because juvenile Coho remain in their spawning stream for a full year after
25 emerging from the gravel, they are exposed to the full range of freshwater
26 conditions. Most smolts migrate to the ocean between March and June, with most
27 leaving in April and May. Coho Salmon typically spend about 16 to 18 months in
28 the ocean before returning to their natal streams to spawn as 3- or 4-year-olds,
29 age 1.2 or 2.2. Trinity River Coho are mostly 3-year-olds. Some precocious
30 males, called jacks, return to spawn after only 6 months in the ocean.

31 Juvenile Coho Salmon in the Trinity River spend up to a full year in fresh water
32 before migrating to the ocean. Their habitat preferences change throughout the
33 year and are highly influenced by water temperature. During summer, when
34 Coho are most actively feeding and growing, they spend more time closer to main
35 channel habitats. Coho use slower water than steelhead or Chinook Salmon.
36 Coho juveniles are more oriented to submerged objects, such as woody debris,
37 while Chinook and steelhead select habitats in summer based largely on water
38 movement and velocities, although the species are often intermixed in the same
39 habitat. Juvenile Coho use the same habitats as pikeminnows, a possible reason
40 that Coho are not present in Central Valley watersheds. Juvenile Coho would be
41 vulnerable to predation from larger pikeminnows during warm-water periods.
42 Pikeminnow do not occur in Southern Oregon/Northern California Coast coho
43 streams. When the water cools in fall, juvenile Coho move farther into backwater
44 areas or into off-channel areas and beaver ponds if available. There is often no
45 water velocity in the areas inhabited by Coho during winter. These same

1 off-channel habitats are often dry or unsuitable during summer because
2 temperatures get too high.

3 Lewiston Dam blocks access to 109 miles of upstream habitat. Trinity River
4 Hatchery produces Coho Salmon with a production goal of 500,000 yearlings to
5 mitigate for the upstream habitat loss. Habitat in the Trinity River has changed
6 since flow regulation with the encroachment of riparian vegetation restricting
7 channel movement and limiting fry rearing habitat (Trush et al. 2000). According
8 to the Trinity River Restoration Plan, higher peak flows are needed to restore
9 attributes of a more alluvial river such as alternate bar features and more
10 off-channel habitats. These are projected in the restoration plan to provide better
11 rearing habitat for Coho Salmon than the dense riparian vegetation currently
12 present. A number of restoration actions have been completed. A new flow
13 schedule has provided higher spring releases to geomorphically maintain habitat.
14 Physical habitat manipulations have been implemented providing better juvenile
15 rearing in selected sites along the river.

16 **9B.8.3 Population Trends**

17 Coho Salmon were not likely the dominant species of salmon in the Trinity River
18 before dam construction. However, Coho were widespread in the Trinity Basin
19 ranging as far upstream as Stuarts Fork above Trinity Dam. Wild Coho in the
20 Trinity Basin today are not abundant, and the majority of the fish returning to the
21 river are of hatchery origin. An estimated 2 percent (200 fish) of the total Coho
22 Salmon run in the Trinity River were composed of naturally produced Coho from
23 1991 through 1995 at a point in the river near Willow Creek (USFWS 1998).
24 This, in part, prompted the threatened status listing in 1997. These estimates
25 included a combination of hatchery produced and wild Coho. About 10 percent
26 of the Coho were naturally produced since 1995.

27 **9B.8.4 Hatchery Influences**

28 The Trinity River portion of the Southern Oregon/Northern California Coast Coho
29 Salmon ESU is predominately of hatchery origin. Termination of hatchery
30 production of Coho Salmon at the Mad River and Rowdy Creek facilities has
31 eliminated further potential adverse risks associated with hatchery releases from
32 these facilities. Likewise, restrictions on recreational and commercial harvest of
33 Coho Salmon since 1994 likely have had a positive impact on Coho Salmon adult
34 returns.

35 **9B.8.5 References**

36 DFG (California Department of Fish and Game). 2002. *Status review of*
37 *California coho salmon north of San Francisco*. Candidate Species Status
38 Review Report 2002-3. Report to the California Fish and Game
39 Commission.

40 NMFS (National Marine Fisheries Service). 1997. Endangered and threatened
41 species: threatened status for southern Oregon/northern California coast
42 evolutionarily significant unit (ESU) of coho salmon. *Federal Register*
43 62: 24588-24609.

- 1 Trush, W. J., S. M. McBain, and L. B. Leopold. 2000. Attributes of an alluvial
 2 river and their relation to water policy and management. *PNAS* 97:
 3 11858-11863.
- 4 USFWS (U.S. Fish and Wildlife Service). 1997. *Klamath River (Iron Gate Dam
 5 to Seiad Creek) life history periodicities for Chinook, coho and steelhead.*
 6 U.S. Fish and Wildlife Service, Coastal California Fish and Wildlife
 7 Office, Arcata, California.

8 **9B.9 Sacramento Splittail (*Pogonichthys*** 9 ***macrolepidotus*)**

10 **9B.9.1 Legal Status**

11 Federal: None

12 State: Species of Special Concern

13 USFWS listed Sacramento Splittail as a threatened species on March 10, 1999,
 14 because of the reduction in its historical range and because of the large population
 15 decline during the 1987-93 drought (USFWS 1996, 1999). On June 23, 2000, the
 16 Federal Eastern District Court of California found the final rule to be unlawful
 17 and on September 22, 2000, remanded the determination back to USFWS for a
 18 reevaluation of the final decision. After a thorough review, USFWS removed the
 19 Sacramento Splittail from the list of threatened species (USFWS 2003) and
 20 reaffirmed this decision in 2010 (USFWS 2010).

21 **9B.9.2 Distribution**

22 Sacramento Splittail are endemic to the Sacramento and San Joaquin River
 23 systems of California, including the Delta and the San Francisco Bay.

24 Historically, splittail were found in the Sacramento River as far upstream as
 25 Redding, in the Feather River to Oroville, and in the American River upstream to
 26 Folsom. In the San Joaquin River, they were once documented as far upstream as
 27 Friant (Rutter 1908). Splittail are thought to have originally ranged throughout
 28 the San Francisco estuary, with catches reported by Snyder (1905) from southern
 29 San Francisco Bay and at the mouth of Coyote Creek.

30 In wet years, Sacramento Splittail have been found in the San Joaquin River as far
 31 upstream as Salt Slough (Saiki 1984, Baxter 1999, Brown and Moyle 1993,
 32 Baxter 2000) and in the Tuolumne River as far upstream as Modesto (Moyle
 33 2002), where the presence of both adults and juveniles during wet years in the
 34 1980s and 1990s indicated successful spawning.

35 When spawning, splittail can be found in the lower reaches of rivers and flooded
 36 areas. Otherwise they are primarily confined to the Delta, Suisun Bay, Suisun
 37 Marsh, the lower Napa River, the lower Petaluma River, and other parts of the
 38 San Francisco estuary (Meng et al. 1994, Meng and Moyle 1995). In general,
 39 splittail are most abundant in Suisun Marsh, especially in drier years (Meng and
 40 Moyle 1995), and reportedly rare in southern San Francisco Bay (Leidy 1984).
 41 Splittail abundance appears to be highest in the northern and western Delta when

1 population levels are low, and they are more evenly distributed throughout the
2 Delta during successful year classes (Sommer et al. 1997, Moyle 2002).
3 Splittail are largely absent from the upper river reaches where they formerly
4 occurred, residing primarily in the lower parts of the Sacramento and San Joaquin
5 rivers and tributaries and in Central Valley lakes and sloughs (Moyle 2002, Moyle
6 et al. 2004). In wet years, however, they have been known to ascend the
7 Sacramento River as far as RBDD and into the lower Feather and American rivers
8 (Baxter et al. 1996; Sommer et al. 1997; Baxter 1999, 2000). The Sutter and Yolo
9 bypasses along the lower Sacramento River appear to be important splittail
10 spawning areas (Sommer et al. 1997). Splittail now migrate into the San Joaquin
11 River only during wet years, and use of the Sacramento River and its tributaries is
12 likely more important (Moyle 2002).

13 **9B.9.3 Life History and Habitat Requirements**

14 **9B.9.3.1 Non-Breeding**

15 Non-reproductive adult splittail are most abundant in moderately shallow,
16 brackish areas, but can also be found in freshwater areas with tidal or riverine
17 flow (Moyle et al. 2004). Non-breeding splittail are found in temperatures
18 ranging from 5 to 24°C, depending on the season, and acclimated fish can survive
19 temperatures up to 33°C for short periods (Young and Cech 1996). Juveniles and
20 adult splittail demonstrate optimal growth at 20°C and signs of physiological
21 distress only above 29°C (Young and Cech 1995).

22 Because splittail are adapted for living in brackish waters with fluctuating
23 conditions, they are tolerant of high salinities and low dissolved oxygen (DO)
24 levels. Splittail are often found in salinities of 10 to 18 parts per thousand (ppt),
25 although lower salinities may be preferred (Meng and Moyle 1995) and can
26 survive low DO levels (0.6 to 1.2 milligrams per liter for young-of-the-year,
27 juveniles, and subadults) (Young and Cech 1995, 1996). Because splittail have a
28 high tolerance for variable environmental conditions (Young and Cech 1996) and
29 are generally opportunistic feeders (prey includes mysid shrimp, clams, copepods,
30 amphipods, and terrestrial invertebrates), reduced prey abundance will not likely
31 have major population-level impacts. Year class success appears dependent on
32 access and availability of floodplain spawning and rearing habitats, high outflow,
33 and wet years (Sommer et al. 1997).

34 **9B.9.3.2 Spawning**

35 Adults typically migrate upstream from brackish areas in January and February
36 and spawn in fresh water on inundated floodplains in March and April (Moyle
37 et al. 2004). Foraging in flooded areas along the main rivers, bypasses, and tidal
38 freshwater marsh areas of Montezuma and Suisun sloughs and San Pablo Bay
39 before the onset of spawning may contribute to spawning success and survival of
40 adults after spawning (Moyle et al. 2004). Splittail are adapted to the wet-dry
41 climatic cycles of Northern California and thus concentrate their reproductive
42 effort in wet years when potential success is enhanced by the availability of
43 inundated floodplain (Meng and Moyle 1995, Sommer et al. 1997). Splittail are

1 thought to be fractional spawners, with individuals spawning over a protracted
2 period—often as long as several months (Wang 1995). Older fish are believed to
3 begin spawning first (Caywood 1974).

4 Splittail eggs are deposited in flooded areas among submerged vegetation, to
5 which they adhere until hatching. Rising flows appear to be the major trigger for
6 splittail spawning, but increases in water temperature and day length may also be
7 factors (Moyle et al. 2004). Spawning typically occurs on inundated floodplains
8 from February through June, with peak spawning in March and April.

9 Information indicates that splittail spawn in open areas with moving, turbid water
10 less than 5 feet (1.5 m) deep, among dense annual vegetation and where water
11 temperatures are below 15°C (Moyle et al. 2004). Perhaps the most important
12 spawning habitat in the eastern Delta is the Cosumnes River floodplain, where
13 ripe splittail have been observed in flooded fields with cool temperatures below
14 15°C, turbid water, and submerged terrestrial vegetation (Crain et al. 2004).

15 Females are typically highly fecund, with the largest individuals potentially
16 producing 100,000 or more eggs (Daniels and Moyle 1983, Feyrer and Baxter
17 1998). Fecundity has been found to be variable, however, and may be influenced
18 by food supplies in the year before spawning (Moyle et al. 2004). The adhesive
19 eggs are released by the female, fertilized by one or more attendant males, and
20 adhere to vegetation until hatching (Moyle 2002). Splittail eggs, which are 0.4 to
21 0.6 inch (1.0 to 1.6 mm) in diameter (Wang 1986, Feyrer and Baxter 1998), begin
22 to hatch within 3 to 7 days, depending on temperature (Bailey 1994). Eggs laid in
23 clumps hatch more quickly than individual eggs (Moyle et al. 2004). Within 5 to
24 7 days after hatching, swim bladder inflation occurs, and larvae begin active
25 swimming and feeding (Moyle 2002). Little is known regarding the tolerance of
26 splittail eggs and developing larvae to DO, temperature, pH, or other water
27 quality parameters, or to other factors such as physical disturbance or desiccation.

28 **9B.9.3.3 Larvae**

29 Juveniles are strong swimmers and are usually found in shallow (less than 6.6 feet
30 [2 m] deep), turbid water (Young and Cech 1996). As their swimming ability
31 increases, juveniles move away from the shallow areas near spawning sites into
32 faster, deeper water (Moyle 2002). Floodplain habitat offers high food quality
33 and production and low predator densities to increase juvenile growth.

34 After emergence, most larval splittail remain in flooded riparian areas for 10 to
35 14 days, most likely feeding among submerged vegetation before moving off
36 floodplains into deeper water as they become stronger swimmers (Sommer et al.
37 1997, Wang 1986). Although juvenile splittail rear in upstream areas for a year or
38 more (Baxter 1999), most move to tidal waters after only a few weeks, often in
39 response to flow pulses (Moyle et al. 2004). The majority of juveniles move
40 downstream into shallow, productive bay and estuarine waters from April to
41 August (Meng and Moyle 1995). Growth likely depends on the availability of
42 high-quality food, especially in the first year of life (Moyle et al. 2004).

1 **9B.9.4 Population Trends**

2 A variety of surveys have compiled splittail abundance data. None of these,
3 however, was specifically designed to systematically sample splittail abundance,
4 and definitive conclusions are therefore not possible (Moyle et al. 2004).

5 Combined, the survey data indicate that successful reproduction occurs on a
6 yearly basis, but large numbers of juvenile splittail are produced only when
7 outflow is relatively high. Thus, the majority of adult fish in the population
8 probably result from spawning in wet years (Moyle et al. 2004). The stock-
9 recruitment relationship in splittail is apparently weak, indicating that given the
10 right environmental conditions, a small number of large females can produce
11 many young (Sommer et al. 1997, Meng and Moyle 1995).

12 Accounts of early fisheries suggested that splittail had large seasonal migrations
13 (Walford 1931). Splittail migration now appears closely tied to river outflow. In
14 wet years with increased river flow, adult splittail will still move long distances
15 upstream to spawn, allowing juvenile rearing in upstream habitats. The upstream
16 migration is smaller during dry years, although larvae and juveniles are often
17 found upstream of Sacramento to Colusa or Ord Bend on the Sacramento River
18 (Moyle et al. 2004). The tidal upper estuary, including Suisun Bay, provides most
19 juvenile rearing habitat, although young-of-the-year may rear over a broader area,
20 including the lower Sacramento River. Brackish water provides optimal rearing
21 habitat for splittail.

22 DFW estimates that splittail during most years are only 35 to 60 percent as
23 abundant as they were in 1940 (DFG 1992). DFW midwater trawl data indicate
24 considerable fluctuations in splittail numbers since the mid-1960s, with
25 abundance often tracking river and Delta outflow conditions. The overall trends
26 include a decline from the mid-1960s to the late 1970s, somewhat of a resurgence
27 through the mid-1980s, and another decline from the mid-1980s through 1994
28 (Moyle 2002). In 1995 and 1998, the population increased dramatically,
29 demonstrating the extreme short- and long-term variability of splittail recruitment
30 success and the apparent correlation with river outflow (Sommer et al. 1997). In
31 2006, when spring outflows were the highest since 1998, beach seine surveys
32 conducted by USFWS in the lower portion of the estuary recorded the highest
33 number of 0+ fish individuals since the surveys began in 1992 (Greiner et al.
34 2007). Surveys in the upper portions of the estuary showed a decline in catches of
35 splittail and many other Delta fish. These declines were coupled with declines in
36 zooplankton, which are the primary food source for splittail (Hieb et al. 2004).
37 Pesticide use in the Central Valley may contribute to reduced zooplankton
38 abundance in the Delta and thus to the POD (Oros and Werner 2005).

39 Splittail may also be negatively affected by the introduction of the overbite clam
40 (*Potamocorbula amurensis*) in the 1980s, which resulted in a collapse of opossum
41 shrimp (*Neomysis mercedis*) populations, which were a primary source of food for
42 splittail. The recent introduction of the Siberian prawn may similarly pose a
43 threat to splittail food sources, as the Siberian prawns prey on mysid shrimp,
44 which make up a large portion of splittail diets (Moyle et al. 2004). River outflow
45 in February through May can explain between 55 and 69 percent of the variability

1 in abundance of splittail young, depending on the abundance measure. Age -0
 2 abundance of splittail declined in the estuary during most dry years, particularly
 3 in the drought that began in 1987 (Sommer et al. 1997). However, not all wet
 4 years result in high splittail recruitment because recruitment success largely
 5 depends on the availability of flooded spawning habitat. In 1996, for example,
 6 most high river flows occurred in December and January, before the onset of the
 7 splittail spawning season (Moyle 2002).

8 **9B.9.5 References**

- 9 Bailey, H. C. 1994. Sacramento splittail work continues. *Interagency Ecological*
 10 *Program Newsletter 7*: Article 3.
- 11 Baxter, R. D. 1999. Status of splittail in California. *California Fish and Game 85*:
 12 28-30.
- 13 _____. 2000. Splittail and longfin smelt. *IEP Newsletter 13*: 19-21.
- 14 Baxter, R. D., W. Harrell, and L. Grimaldo. 1996. 1995 Splittail spawning
 15 investigations. *Interagency Ecological Program Newsletter 9*: 27–31.
- 16 Brown, L. R., and P. B. Moyle. 1993. Distribution, ecology, and status of fishes of
 17 the San Joaquin River drainage, California. *California Fish and Game*
 18 *Bulletin 79*: 96-113.
- 19 Caywood, M. L. 1974. *Contributions to the life history of the splittail*
 20 *(Pogonichthys macrolepidotus) (Ayres)*. Master's thesis. California State
 21 University, Sacramento, California.
- 22 Crain, P. K., K. Whitener, and P. B. Moyle. 2004. Use of a restored central
 23 California floodplain by larvae of native and alien fishes. *American*
 24 *Fisheries Society Symposium 39*: 125–140.
- 25 Daniels, R. A., and P. B. Moyle. 1983. Life history of splittail (Cyprinidae:
 26 *Pogonichthys macrolepdotus*) in the Sacramento-San Joaquin Estuary.
 27 *Fishery Bulletin 84*: 105–117.
- 28 DFG (California Department of Fish and Game). 1992. *Impact of water*
 29 *management on splittail in the Sacramento-San Joaquin estuary*. WRINT-
 30 CDFG-Exhibit 5. State Water Resources Control Board hearing for setting
 31 interim standards for the Delta.
- 32 Feyrer, F. V., and R. D. Baxter. 1998. Splittail fecundity and egg size. *California*
 33 *Fish and Game 84*: 119–126.
- 34 Greiner, T., M. Fish, S. Slater, K. Hieb, J. Budrick, J. DuBois, and D. Contreras.
 35 2007. 2006 Fishes: Annual status and trends report for the San Francisco
 36 Estuary. *Interagency Ecological Program Newsletter 20*(2).
- 37 Hieb, K., T. Greiner, and S. Slater. 2004. San Francisco Bay species: 2003 Status
 38 and trends report. *Interagency Ecological Program Newsletter 17*:17-28.
- 39 Leidy, R. A. 1984. Distribution and ecology of stream fishes in the San Francisco
 40 Bay drainage. *Hilgardia 52*: 1–175.

- 1 Meng, L., P. B. Moyle, and B. Herbold. 1994. Changes in abundance and
 2 distribution of native and introduced fishes of Suisun Marsh. *Transactions*
 3 *of the American Fisheries Society* 123: 498–507.
- 4 Meng, L., and P. B. Moyle. 1995. Status of splittail in the Sacramento-
 5 San Joaquin Estuary. *Transactions of the American Fisheries Society* 124:
 6 538–549.
- 7 Moyle, P. B. 2002. *Inland fishes of California*. Revised edition. University of
 8 California Press, Berkeley.
- 9 Moyle, P. B., R. D. Baxter, T. Sommer, T. C. Foin, and S. A. Matern. 2004.
 10 Biology and population dynamics of Sacramento splittail (*Pogonichthys*
 11 *macrolepidotus*) in the San Francisco Estuary: a review. *San Francisco*
 12 *Estuary and Watershed Science*. 2: Article 3.
- 13 Oros, D. R., and I. Werner. 2005. *Pyrethroid insecticides: an analysis of use*
 14 *patterns, distributions, potential toxicity and fate in the Sacramento-*
 15 *San Joaquin Delta and Central Valley*. White Paper for the Interagency
 16 Ecological Program. SFEI Contribution 415. San Francisco Estuary
 17 Institute, Oakland, California.
- 18 Rutter, C. 1908. The fishes of the Sacramento-San Joaquin basin, with a study of
 19 their distribution and variation. *Bulletin of the U.S. Bureau of Fisheries*
 20 27: 103-152.
- 21 Saiki, M. K. 1984. Environmental conditions and fish faunas in low elevation
 22 rivers on the irrigated San Joaquin Valley floor, California. *California*
 23 *Fish and Game* 70: 145-157.
- 24 Snyder, J. O. 1905. Notes on the fishes of the streams flowing into San Francisco
 25 Bay. *United States Bureau of Fisheries* 5: 327–338.
- 26 Sommer, T., R. Baxter, and B. Herbold. 1997. Resilience of splittail in the
 27 Sacramento-San Joaquin estuary. *Transactions of the American Fisheries*
 28 *Society* 126: 961–976.
- 29 USFWS (U.S. Fish and Wildlife Services). 1996. *Recovery plan for the*
 30 *Sacramento-San Joaquin Delta native fishes*. U.S. Fish and Wildlife
 31 Service, Portland, Oregon.
- 32 _____. 1999. *Endangered and threatened wildlife and plants; determination of*
 33 *threatened status for the Sacramento splittail*. Federal Register 64: 5963–
 34 5981.
- 35 _____. 2003. *Endangered and Threatened Wildlife and Plants; Notice of*
 36 *Remanded Determination of Status for the Sacramento splittail*
 37 *(Pogonichthys macrolepidotus); Final Rule*. Federal Register 68: 55140.
- 38 _____. 2010. *Endangered and Threatened Wildlife and Plants; 12-month Finding*
 39 *on a Petition to list the Sacramento Splittail as Endangered or*
 40 *Threatened*. Federal Register 75: 62070-62095.

- 1 Walford, L. A. 1931. *Handbook of common commercial and game fishes of*
 2 *California*. California Department of Fish and Game Fish Bulletin 28.
- 3 Wang, J. C. S. 1986. *Fishes of the Sacramento-San Joaquin estuary and adjacent*
 4 *waters, California: a guide to the early life histories*. Technical Report 9.
 5 Prepared for the Interagency Ecological Study Program for the
 6 Sacramento-San Joaquin Estuary by California Department of Water
 7 Resources, California Department of Fish and Game, Bureau of
 8 Reclamation, and U.S. Fish and Wildlife Service.
- 9 _____ 1995. *Observations of early life stages of splittail (Pogonichthys*
 10 *macrolepidotus) in the Sacramento-San Joaquin estuary, 1988 to 1994*.
 11 Interagency Ecological Program Technical Report 43.
- 12 Young, P. S., and J. J. Cech, Jr. 1995. *Salinity and dissolved oxygen tolerance of*
 13 *young-of-the-year and juvenile Sacramento splittail*. Consensus building
 14 in resource management. American Fisheries Society, California-Nevada
 15 Chapter.
- 16 _____ 1996. Environmental tolerances and requirements of splittail. *Transactions*
 17 *of the American Fisheries Society* 125: 664–678.

18 **9B.10 Delta Smelt (*Hypomesus transpacificus*)**

19 **9B.10.1 Legal Status**

20 Federal: Threatened, Designated Critical Habitat
 21 State: Endangered

22 The USFWS listed the Delta Smelt as threatened in March 1993 (USFWS 1993),
 23 and critical habitat for this species was designated in 1994 (USFWS 1994). The
 24 Delta Smelt was one of eight fish species addressed in the Recovery Plan for the
 25 Sacramento–San Joaquin Delta Native Fishes (USFWS 1996). This recovery plan
 26 is currently under revision. The 2004 status review affirmed the need to retain the
 27 Delta Smelt as a threatened species (USFWS 2004). A 12-month finding on a
 28 petition to reclassify the Delta Smelt was completed in April 2010 and the
 29 USFWS determined that re-classifying the Delta Smelt from a threatened to an
 30 endangered species was warranted, but precluded by other higher-priority listing
 31 actions (USFWS 2010).

32 **9B.10.2 Distribution**

33 Delta Smelt are endemic to and resident in the Delta and San Francisco Bay.
 34 According to a recent review (Merz et al. 2011), the distribution of Delta Smelt
 35 includes an area from northern San Francisco Bay in the west, the confluence of
 36 the Sacramento and Feather rivers in the north, and the junction of Old and San
 37 Joaquin rivers in the south. The highest densities most frequently occur near the
 38 center of their range, which appears to extend from Suisun Marsh down through
 39 Grizzly Bay and east Suisun Bay through the confluence of the Sacramento and

1 San Joaquin rivers, and into the lower portions of the Sacramento River, Cache
2 Slough area, and the Sacramento Deepwater Ship Channel.

3 Delta Smelt abundance and geographic distribution are dependent upon
4 freshwater outflows and the salinity of the Bay and Delta (Herbold et al. 1992).
5 There is a close association between Delta Smelt abundance and surface salinity
6 of 0–18 practical salinity units (psu) (psu are roughly equivalent to ppt),
7 suggesting that their distribution is determined largely by the interaction with
8 salinity conditions as determined by tidal currents, freshwater outflow, and
9 diffusion, rather than by geography (Bennett 2000, 2005; Moyle 2002). For
10 instance, water clarity and salinity were found to be the most reliable abiotic
11 predictors of Delta Smelt abundance during the summer and fall (Feyrer et al.
12 2007, Nobriga et al. 2008). In addition, geographic distribution for particular life
13 stages can vary dramatically between dry and wet years. Thus, in low outflow
14 years, Delta Smelt occur primarily in the lower Sacramento River, with the area
15 near Decker Island consistently exhibiting greatest catch over time. In years of
16 very high outflow, however, their distribution extends into San Pablo Bay and the
17 Napa River (Bennett 2000).

18 **9B.10.3 Life History and Habitat Requirements**

19 Overall, the Delta Smelt life cycle is completed in the brackish and tidal
20 freshwater reaches of the upper San Francisco Estuary. However, salinity
21 requirements vary by life stage. Apart from spawning and egg-embryo
22 development, the distribution and movements of all life stages are influenced by
23 transport processes associated with water flows in the estuary, which also affect
24 the quality and location of suitable open water habitat (Dege and Brown 2004;
25 Feyrer et al. 2007; Nobriga et al. 2008).

26 **9B.10.3.1 Spawning**

27 Delta Smelt generally exhibit an annual, 1-year lifecycle. They are found at
28 0-18 psu surface salinity (Baxter et al. 1999), although most are caught at
29 salinities less than 6.0 psu, with older juveniles and adults being found at the
30 higher end of that gradient (Bennett 2005). Delta Smelt feed primarily on
31 planktonic copepods, cladocerans, and amphipods (Baxter et al. 2008). In recent
32 years, a small to moderate number of Delta Smelt have been observed in the Deep
33 Water Ship Channel during the late fall. The Deep Water Ship Channel can
34 provide suitable water temperatures for Delta Smelt year-round (Sommer and
35 Mejia 2013), which likely promotes freshwater residence in Delta Smelt in this
36 region of the Delta (Sommer and Mejia 2013).

37 Delta Smelt are weakly anadromous and undergo a spawning migration from the
38 low salinity zone to freshwater in most years (Grimaldo et al. 2009; Sommer et al.
39 2011). Spawning migrations occur between late December and late February,
40 typically during “first flush” periods when inflow and turbidity increase on the
41 Sacramento and San Joaquin Rivers (Grimaldo et al. 2009, Sommer et al. 2011).
42 Notably, spawning movements are not always upstream. Under high outflow
43 conditions, when total outflow exceeds 100,000 cubic feet per second (cfs), adult
44 smelt tend to concentrate and spawn in Suisun Bay, Cache Slough Complex, and

1 Napa River (Hobbs et al. 2007; Sommer et al. 2011). During drier years, when
 2 total outflow is less than 20,000 cfs, smelt tend to concentrate and spawn in the
 3 Cache Slough Complex and western Delta.

4 Adequate flows and suitable water quality are needed to attract migrating adults in
 5 the Sacramento and San Joaquin River channels and their associated tributaries,
 6 including Cache and Montezuma sloughs and their tributaries (USFWS 1996).
 7 Adult smelt do not spawn immediately after migration to freshwater, but appear to
 8 stage in upstream habitats (Sommer et al. 2011). Spawning typically commences
 9 when water temperatures reach 12°C, which typically occurs in early March.
 10 Spawning can continue into July (Wang 1986, Sweetnam and Stevens 1993),
 11 although most spawning takes place from early April to mid-May (Moyle 2002).

12 Delta Smelt are believed to spawn in shallow water along edges of rivers and
 13 sloughs subject to tidal influence (USFWS 2001). Based upon the occurrence of
 14 ripe females and yolk-sac larvae, spawning areas during dry and typical years are
 15 found in the north Delta reaches of the Sacramento River (Moyle 2002).
 16 Spawning locations in the Delta have not been identified and are inferred from
 17 larval catches (Bennett 2005). Larval fish have been observed in Montezuma
 18 Slough (Wang 1986), Suisun Slough in Suisun Marsh (Moyle 2002), the Napa
 19 River estuary (Stillwater Sciences 2006), the Sacramento River above Rio Vista,
 20 and Cache, Lindsey, Georgiana, Prospect, Beaver, Hog, Sycamore, and Barker
 21 sloughs (USFWS 1996). During wet years, Delta Smelt can be found spawning
 22 throughout most of the Delta, Suisun Marsh, and west to the Napa River (Herbold
 23 et al. 1992).

24 Although the specific substrates or habitats used for spawning by Delta Smelt are
 25 not known, spawning habitat preferences of closely related species (Bennett 2005)
 26 suggest that spawning may occur in shallow areas over sandy substrates.
 27 Although smelt can be found within a wide salinity range, from 0 to 18.4 ppt
 28 (Swanson et al. 2000), spawning occurs within in freshwater (Wang 1986).
 29 Spawning apparently can occur at temperatures ranging from 45-72°F (7-22°C)
 30 (Moyle 2002), but most often takes place between 45 and 59°F (7 and 15°C)
 31 (Wang 1986).

32 Spawning is thought to occur at night during new or full moons when the tide is
 33 low (Moyle 2002). Females (2.3-2.8 in [59-70 mm] SL) typically lay between
 34 1,200 and 2,600 eggs (Moyle et al. 1992) and the relationship between female size
 35 (FL) and fecundity has been determined to be: Number of eggs = $0.266FL^{2.089}$
 36 (Mager 1996). Most adults die after spawning, although a small number remain
 37 in the population for a second year (Moyle 2002) and may contribute
 38 disproportionately to the egg supply because of their increased size (3.5-4.7 in
 39 [90-120 mm] SL) (Moyle 2002).

40 **9B.10.3.2 Hatching and Larval Distribution**

41 No data are available on optimal temperature for survival of embryos, though
 42 some data suggest that high temperatures correspond to low hatching success and
 43 low embryo survival (R. Mager, unpubl. data; as cited in Winternitz and
 44 Wadsworth 1997). According to Moyle (2002), “it is likely that survival

1 decreases as temperature increases beyond 18°C [64°F].” At temperatures
2 between 59 and 62°F (14.8 and 16.5°C), embryonic development is reported to
3 take approximately 9-13 days (Mager 1996). Although hatching has been
4 detected from late February to June, peak hatching typically occurs in April.

5 Newly hatched smelt begin feeding on rotifers and other microscopic prey
6 approximately 4-5 days after hatching, maintaining a position just above the
7 bottom with the help of a large oil globule that makes them semi-buoyant (Mager
8 1996). The swim bladder and fins are fully developed several weeks later, and
9 larvae rise up into the water column (Moyle 2002). During high outflow periods,
10 larvae are distributed more widely as the spawning range extends further west
11 when Delta outflows are high (Hobbs et al. 2007). Dege and Brown (2004) found
12 that larvae less than 20 mm rear 5 to 20 km upstream of X2 (Dege and Brown
13 2004; Sommer and Mejia 2013). As larvae grow and water temperatures increase
14 in the Delta (to approximately 23°C), their distribution shifts towards the low
15 salinity zone (Dege and Brown 2004; Nobriga et al. 2008), where they circulate
16 with the abundant zooplankton (Moyle 2002). By fall, the centroid of Delta Smelt
17 distribution is tightly coupled with X2 (Sommer et al. 2011; Sommer and Mejia
18 2013).

19 Sommer and Mejia (2013) conducted a General Additive Model (GAM) analysis
20 of Delta Smelt catch data from the 20-mm survey to determine suitable habitat
21 parameters. They found larval Delta Smelt are more frequently captured in turbid
22 and low salinity water. The analysis also showed that larval smelt presence in the
23 survey peaked when water temperatures reach 20°C with low capture probability
24 below 10°C and above 25°C.

25 The abundance of suitable rearing habitat for larvae varies from year to year,
26 depending upon when peak spawning occurs. Peak larval density may occur as
27 late as July or August. Base flows and pulse flows that transport and provide
28 behavioral cues for Delta Smelt larvae and juveniles from February through June
29 may not be adequate if larval peaks occur in July or August.

30 **9B.10.3.3 Juvenile Rearing and Growth**

31 The specific geographic area critical to the maintenance of suitable rearing habitat
32 for Delta Smelt extends eastward from Carquinez Strait, up the Sacramento River
33 to its confluence with Three Mile Slough (at RM 9), and south along the
34 San Joaquin River including Big Break (USFWS 1996). Within this area, Delta
35 Smelt typically rear in shallow (less than 10 ft [3 m]), open estuarine waters
36 (Moyle 2002), in salinities ranging from 2-7 ppt (Swanson and Cech 1995) where
37 “fresh and brackish water mix and hydrodynamics are complex as a result of the
38 meeting of tidal and riverine currents” (Moyle 2002). These conditions are
39 typically most common in Suisun Bay, which provides vital nursery habitat for
40 Delta Smelt. When the mixing zone is located in Suisun Bay, it provides optimal
41 conditions for algal and zooplankton growth, an important food source for Delta
42 Smelt (Moyle 2002). When freshwater outflow is low, the mixing zone moves
43 further up into the deeper, narrow channels of the Delta and Sacramento River,
44 reducing food availability and total area available to the smelt (Moyle 2002).

1 Water quality preferences and thresholds for Delta Smelt are not well
2 documented. Winternitz and Wadsworth (1997) observed that fewer Delta Smelt
3 were collected in areas of higher temperatures than in areas of lower
4 temperatures. Because other factors were not controlled, it is not clear whether
5 temperature or other factors were driving Delta Smelt distribution. Nobriga et al.
6 (2000) reported that Delta Smelt tolerated slightly higher water temperatures at a
7 salinity of 4 ppt than in fresh water, but noted that further study is needed of these
8 potentially interacting factors. Similar to larvae, a GAM analysis of the tow net
9 survey data shows that suitable smelt habitat is best defined by water clarity,
10 specific conductance (salinity), water temperature (Nobriga et al. 2008). As
11 previously noted, some juvenile smelt will remain in the Sacramento Deep Water
12 Ship Channel during the summer and fall months. The channel is deep, turbid,
13 and offers some temperature refuge, which may explain why smelt remain in this
14 freshwater habitat when most other smelt at this life stage are in found in the low
15 salinity zone.

16 Planktonic copepods, cladocerans, amphipods, and, to a lesser extent, insect
17 larvae, are the primary prey items for Delta Smelt (Moyle 2002). Delta Smelt
18 larvae have more specific prey-size requirements for first feeding. In a study
19 conducted in the northern estuary and Delta, Lott (1998) found that smaller size
20 classes of Delta Smelt tended to consume more nauplii and juvenile copepods,
21 while larger size classes consumed more adult copepods. It appears that food
22 availability after yolk-sac absorption is critical in determining success of Delta
23 Smelt (Nobriga 1998). However, it is not known if a limited food supply
24 contributes to reduced year-class success and therefore has population-level
25 implications.

26 Juvenile Delta Smelt grow rapidly, typically reaching 1.6-2 inches (40-50 mm)
27 FL by early August (Radtke 1966, Moyle et al. 1992). Growth rate appears to be
28 dependent on the quality and abundance of food (Moyle 2002). Adult length
29 (2.2-2.8 inches [55-70 mm] SL) is typically reached by September, or
30 approximately 7-9 months after hatching (Moyle 2002). By fall, Delta Smelt are
31 fully capable of altering their distribution to suitable habitat. Using a GAM
32 approach, Feyrer et al. (2007) showed that Delta Smelt habitat is best defined by
33 turbidity and specific conductance (salinity). Unlike the other analyses, Feyrer
34 et al. (2010) converted the GAM model results to a habitat index for Delta Smelt,
35 showing that habitat improves and expands for Delta Smelt when X2 is in Suisun
36 Bay compared to when X2 is located at or above the confluence. The relationship
37 between the habitat index and X2 is asymptotic, whereby the index does not
38 increase for $X2 \leq 74$ km or decrease for $X2 \geq 81$ km. For the period 1967 – 2008,
39 relative abundance of juvenile delta smelt, as measured by the fall midwater trawl
40 index, was positively correlated with the fall habitat index (Feyrer et al. 2010).

41 The quantity and suitability of Delta Smelt habitat increases with higher outflow
42 (Bennett 2005). When the near-bottom mixing zone is contained within Suisun
43 Bay and when adequate outflow from both the Sacramento and San Joaquin rivers
44 have allowed downstream movement, young Delta Smelt are dispersed more
45 widely throughout a large expanse of shallow-water and marsh habitat than when

1 the isohaline is upstream in the narrower, deeper Delta sloughs and channels. If
2 smelt use this habitat and their distribution is wider and shifted downstream,
3 subsequent entrainment in the winter will be reduced. Habitat conditions suitable
4 for transport of larvae and juveniles are needed as early as February 1 and as late
5 as August 31, because the spawning season varies from year to year and starts as
6 early as December and extends until July (USFWS 1996). Adequate river flow is
7 necessary to provide this transport to Suisun Bay and to maintain rearing habitat
8 (USFWS 1996).

9 The abundance of many local estuarine taxa has tended to increase in years when
10 flows into the estuary are high and the X2 location is pushed seaward (Jassby
11 et al. 1995), implying that over the range of historical experience the quantity or
12 suitability of estuarine habitat increases when outflows are high. Feyrer et al.
13 (2007) reported that fall environmental quality has declined over the long-term in
14 the core range of Delta Smelt, including Suisun Bay and the Delta. This decline
15 was largely due to changes in salinity in Suisun Bay and the western Delta, and
16 changes in water clarity within the Delta. Baxter et al. (2008) reported the long-
17 term environmental quality declines for Delta Smelt and Striped Bass are defined
18 by a lowered probability of occurrence in samples based on changes in specific
19 conductance and Secchi depth.

20 Planktonic copepods, cladocerans, amphipods, and, to a lesser extent, insect
21 larvae, are the primary prey items for Delta Smelt (Moyle 2002). Delta Smelt
22 larvae have more specific prey-size requirements for first feeding. In a study
23 conducted in the northern estuary and Delta, Lott (1998) found that smaller size
24 classes of Delta Smelt tended to consume more nauplii and juvenile copepods,
25 while larger size classes consumed more adult copepods. It appears that food
26 availability after yolk-sac absorption is critical in determining success of Delta
27 Smelt (Nobriga 1998). However, it is not known if a limited food supply
28 contributes to reduced year-class success and therefore has population-level
29 implications.

30 The overbite clam has been associated with large changes in phytoplankton
31 abundance in San Francisco Bay and the western Delta (Carlton et al. 1990),
32 causing a decrease in abundance of other species that depend on phytoplankton
33 (zooplankton) for food. Due in part to its efficiency in filtering water, the clarity
34 of Suisun Bay and delta waters has increased. This has affected Delta Smelt by
35 reducing food supply and increasing its susceptibility to predation.

36 **9B.10.4 Population Trends**

37 California Department of Fish and Wildlife has conducted several long-term
38 monitoring surveys that have been used to index the relative abundance of Delta
39 Smelt. The 20-mm Survey has been conducted every year since 1995. This
40 survey targets late-stage Delta Smelt larvae. Most sampling has occurred from
41 April to June. The Summer Townt Survey (TNS) has been conducted nearly
42 every year since 1959. This survey targets 38-mm Striped Bass, but collects
43 similar-sized juvenile Delta Smelt. Most sampling has occurred from June to
44 August. The Fall Midwater Trawl Survey (FMWT) has been conducted nearly

1 every year since 1967. This survey also targets age-0 Striped Bass, but collects
2 Delta Smelt longer than 40 mm. The FMWT samples monthly from September to
3 December. These abundance index time series document the long-term decline of
4 the Delta Smelt.

5 Early statistical assessments of Delta Smelt population dynamics concluded that
6 the relative abundance of the adult Delta Smelt population had only a very weak
7 influence on subsequent juvenile abundance (Sweetnam and Stevens 1993).
8 Thus, early attempts looked for environmental variables that were directly
9 correlated with interannual abundance variation (e.g., Stevens and Miller 1983;
10 Moyle et al. 1992; Sweetnam and Stevens 1993; Jassby et al. 1995). Because
11 these analyses did not find strong support for an outflow-abundance linkage, the
12 prevailing conceptual model was that multiple interacting factors had caused the
13 Delta Smelt decline (Moyle et al. 1992; Bennett and Moyle 1995; Bennett 2005).
14 It has also recently been noted that Delta Smelt's FMWT index is partly
15 influenced by concurrent environmental conditions (Feyrer et al. 2007; 2010).

16 It is now recognized that Delta Smelt abundance plays an important role in
17 subsequent smelt abundance. Bennett (2005) examined (1) the influence of adult
18 stock (FMWT) on the next generation of juveniles (TNS); (2) the influence of the
19 juvenile stock (TNS) on the subsequent adult stock (FMWT); (3) the influence of
20 the FMWT on the following year's FMWT and on the FMWT two years later,
21 and (4) the influence of the TNS abundance on the following year's TNS and on
22 the TNS 2 years later. His conclusions were that (1) 2-year-old Delta Smelt might
23 play an important role in Delta Smelt population dynamics, (2) it was not clear
24 whether juvenile production was a density-independent or density dependent
25 function of adult abundance, and (3) adult production was a density-dependent
26 function of juvenile abundance and the carrying capacity of the estuary to support
27 this life-stage transition had declined over time. These conclusions are also
28 supported by Maunder and Deriso (2011).

29 Delta Smelt were historically one of the most common species in the
30 San Francisco Estuary, but exhibited significant declines during the 1980s (DFG
31 2000). Kimmerer (2002) and Thomson et al. (2010) reported a Delta Smelt step-
32 decline during 1981-1982. Prior to this decline, the stock-recruit data are
33 consistent with "Ricker" type density-dependence where increasing adult
34 abundance resulted in decreased juvenile abundance. Since the decline,
35 recruitment has been positively and essentially linearly related to prior adult
36 abundance, suggesting that reproduction has been basically density-independent
37 for about the past 30 years. In contrast to the transition among generations, the
38 weight of scientific evidence strongly supports the hypothesis that, at least over
39 the history of IEP fish monitoring, Delta Smelt has experienced density-
40 dependence during the juvenile stage of its life cycle (i.e., between the summer
41 and fall) (Bennett 2005; Maunder and Deriso 2011). The most relevant aspect of
42 this juvenile density dependence is that the carrying capacity of the estuary for
43 Delta Smelt has likely declined (Bennett 2005).

44 Therefore, the USFWS (2012) believes that the Delta Smelt population decline
45 has occurred for two basic reasons. First, the compensatory density-dependence

1 that historically enabled juvenile abundance to rebound from low adult numbers
 2 stopped happening. This change had occurred by the early 1980s as described
 3 above. The reason is still not known, but the consequence of the change is that
 4 for the past several decades, adult abundance has driven juvenile production in a
 5 largely density-independent manner (Kimmerer 2011). Second, because juvenile
 6 carrying capacity has declined, juvenile production hits a ‘ceiling’ at a lower
 7 abundance than it once did. This limits adult abundance and possibly per capita
 8 fecundity, which cycles around and limits the abundance of the next generation of
 9 juveniles. The mechanism causing carrying capacity to decline is likely due to the
 10 long-term accumulation of adverse changes in both physical and biological
 11 aspects of habitat during the summer to fall (Bennett et al. 2008; Feyrer et al.
 12 2007; 2010; Maunder and Deriso 2011).

13 **9B.10.5 References**

- 14 Baxter, R., K. Hieb, S. DeLeon, K. Fleming, and J. Orsi. 1999. *Report on the*
 15 *1980–1995 fish, shrimp, and crab sampling in the San Francisco Estuary,*
 16 *California.* Technical Report 63. Prepared by California Department of
 17 Fish and Game, Stockton for the Interagency Ecological Program for the
 18 Sacramento-San Joaquin Estuary.
- 19 Baxter, R., R. Breuer, L. Brown, M. Chotkowski, F. Feyrer, M. Gingras, B.
 20 Herbold, A. Mueller-Solger, M. Nobriga, T. Sommer, and K. Souza. 2008.
 21 *Pelagic organism decline progress report: 2007 synthesis of results.*
 22 Technical Report 227. Interagency Ecological Program for the
 23 San Francisco Estuary. Available at:
 24 [http://www.science.calwater.ca.gov/pdf/workshops/POD/2007_IEPPOD_s](http://www.science.calwater.ca.gov/pdf/workshops/POD/2007_IEPPOD_synthesis_report_031408.pdf)
 25 [ynthesis_report_031408.pdf](http://www.science.calwater.ca.gov/pdf/workshops/POD/2007_IEPPOD_synthesis_report_031408.pdf).
- 26 Bennett, W.A. and Moyle, P. B. 1996. Where have all the fishes gone? Interactive
 27 factors producing fish declines in the Sacramento San Joaquin Estuary.
 28 Pages 519–542 in Hollibaugh, J. T. (ed.), *San Francisco Bay: The*
 29 *Ecosystem.* San Francisco, CA: Pacific Division American Association for
 30 the Advancement of Science. Pages 519–542.
- 31 Bennett, W. A. 2000. *Delta smelt population structure and factors influencing*
 32 *dynamics: implications for the CALFED Ecosystem Restoration Program.*
 33 Draft white paper prepared for CALFED Bay-Delta Program. As cited by
 34 Sam Luoma (preparer) *Delta Smelt and CALFED’s Environmental Water*
 35 *Account, Summary of a Workshop held September 7, 2001, Putah Creek*
 36 *Lodge, University of California, Davis,* by Randall Brown and Wim
 37 Kimmerer.
- 38 Bennett, W. A. 2005. Critical assessment of the delta smelt population in the
 39 San Francisco Estuary, California. *San Francisco Estuary & Watershed*
 40 *Science* 3: Article 1.
- 41 California Resources Agency. 2007. *Pelagic fish action plan.* California
 42 Department of Water Resources and California Department of Fish and
 43 Game, Sacramento, California.

- 1 Carlton, J. T., J. K. Thompson, L. E. Schemel, and F. H. Nichols. 1990.
2 Remarkable invasion of San Francisco Bay (California, USA) by the
3 Asian clam *Potamocorbula amurensis*. I. Introduction and dispersal.
4 *Marine Ecology Progress Series* 66:81-94.
- 5 DFG (California Department of Fish and Game). 2000. *The status of rare,*
6 *threatened, and endangered animals and plants of California: delta smelt.*
7 DFG, Habitat Conservation Planning Branch.
- 8 Dege, M. and L. R. Brown. 2004. Effect of outflow on spring and summertime
9 distribution and abundance of larval and juvenile fishes in the upper
10 San Francisco estuary. In: F. Feyrer, L. R. Brown, R. L. Brown, and J. J.
11 Orsi (eds.), *Early Life History of Fishes in the San Francisco Estuary and*
12 *Watershed*. American Fisheries Society Symposium 39:49–66.
- 13 Feyrer, F., M. L. Nobriga, and T. R. Sommer. 2007. Multi-decadal trends for
14 three declining fish species: habitat patterns and mechanisms in the
15 San Francisco Estuary, California, USA. *Canadian Journal of Fisheries*
16 *and Aquatic Sciences* 64:723–734.
- 17 Feyrer, F., Newman, K., Nobriga, M., and Sommer, T. 2010. Modeling the effects
18 of future freshwater flow on the abiotic habitat of an imperiled estuarine
19 fish. *Estuaries and Coasts* 34:120–128.
- 20 Grimaldo, L. F., T. Sommer, N. Van Ark, G. Jones, E. Holland, P. B. Moyle, P.
21 Smith and B. Herbold. 2009. Factors Affecting Fish Entrainment into
22 Massive Water Diversion in a Tidal Freshwater Estuary: Can Fish Losses
23 Be Managed? *North America Journal of Fisheries Management* 29:1253–
24 1270.
- 25 Herbold, B., A. D. Jassby, and P. B. Moyle. 1992. *San Francisco Estuary Project:*
26 *Status and trends report on aquatic resources in the San Francisco*
27 *Estuary*. Prepared by University of California, Davis under Cooperative
28 Agreement #CE009519-01-1 with the U.S. Environmental Protection
29 Agency.
- 30 Hobbs, J. A., W. A. Bennett, J. Burton, and M. Gras. 2007. Classification of
31 Larval and Adult Delta Smelt to Nursery Areas by Use of Trace Elemental
32 Fingerprinting. *Transactions of the American Fisheries Society* 136:518–
33 527.
- 34 Jassby, A. D., W. J. Kimmerer, S. G. Monismith, C. Armor, J. E. Cloern, T. M.
35 Powell, J. R. Schubel, and T. J. Vendlinski. 1995. Isohaline position as a
36 habitat indicator for estuarine populations. *Ecological Applications* 5:
37 272–289.
- 38 Kimmerer, W. J. 2002. Effects of Freshwater Flow on Abundance of Estuarine
39 Organisms: Physical Effects of Trophic Linkages. *Marine Ecology*
40 *Progress Series* 243:39–55.

- 1 Kimmerer, W. J. 2008. Losses of Sacramento River Chinook salmon and delta
2 smelt (*Hypomesus transpacificus*) to entrainment in water diversions in
3 the Sacramento-San Joaquin Delta. *San Francisco Estuary and Watershed*
4 *Science* 6: Article 2.
- 5 Kimmerer, W. J. 2011. Modeling Delta Smelt Losses at the South Delta Export
6 Facilities. *San Francisco Estuary and Watershed Science* 9(1). Available
7 at: <http://www.escholarship.org/uc/item/Ord2n5vb>.
- 8 Lindberg, J., R. Mager, B. Bridges, S. Doroshov. 1997. Status of delta smelt
9 culture project. *Interagency Ecological Program for the Sacramento-*
10 *San Joaquin Estuary Newsletter* 10: 21–22.
- 11 Lott, J. 1998. Feeding habits of juvenile and adult delta smelt from the
12 Sacramento-San Joaquin river estuary. *Interagency Ecological Program*
13 *for the Sacramento-San Joaquin Estuary Newsletter* 11: 14–19.
- 14 Mager, R. C. 1996. *Gametogenesis, reproduction, and artificial propagation of*
15 *delta smelt, Hypomesus transpacificus*. Doctoral dissertation. University
16 of Davis, California. As cited in Moyle 2002.
- 17 Maunder, M. N., and R. B. Deriso. 2011. A State-Space Multistage Life Cycle
18 Model to Evaluate Population Impacts in the Presence Of Density
19 Dependence: Illustrated with Application to Delta Smelt (*Hypomesus*
20 *transpacificus*). *Canadian Journal of Fisheries and Aquatic Sciences*
21 68:1285–1306.
- 22 Merz., J.E., S. Hamilton, P.S. Bergman, and B. Cavallo. 2011. Spatial perspective
23 for delta smelt: a summary of contemporary survey data. *California Fish*
24 *and Game*, 97(4), pp. 164-189.
- 25 Moyle, P. B. 2002. *Inland fishes of California*. Revised edition. University of
26 California Press, Berkeley.
- 27 Moyle, P. B., B. Herbold, D. E. Stevens, and L. W. Miller. 1992. Life history and
28 status of delta smelt in the Sacramento-San Joaquin estuary, California.
29 *Transactions of the American Fisheries Society* 121: 67–77.
- 30 Nobriga, M. 1998. Evidence of food limitation in larval delta smelt. *Interagency*
31 *Ecological Program for the Sacramento-San Joaquin Estuary Newsletter*
32 11: 20–24.
- 33 Nobriga, M., Z. Hymanson, and R. Oltmann. 2000. Environmental factors
34 influencing the distribution and salvage of young delta smelt: a
35 comparison of factors occurring in 1996 and 1999. *Interagency Ecological*
36 *Program for the Sacramento-San Joaquin Estuary Newsletter* 13: 55–65.
- 37 Nobriga, M. L., T. R. Sommer, F. Feyrer, and K. Fleming. 2008. Long-term
38 trends in summertime habitat suitability for delta smelt (*Hypomesus*
39 *transpacificus*). *San Francisco Estuary and Watershed Science* 6.

- 1 Radtke, L. D. 1966. Distribution of smelt, juvenile sturgeon, and starry flounder
2 in the Sacramento-San Joaquin Delta. Edited by J. L. Turner and D. W.
3 Kelley, 115–119. Ecological studies of the Sacramento-San Joaquin Delta,
4 Part 2. Fish Bulletin 136. California Department of Fish and Game.
- 5 Sommer, T., and Mejia, F. 2013. A place to call home: a synthesis of delta smelt
6 habitat in the upper San Francisco Estuary. *San Francisco Estuary and*
7 *Watershed Science* 11(2). Available at:
8 <http://www.escholarship.org/uc/item/32c8t244>.
- 9 Sommer, T., F. Mejia, M. Nobriga, F. Feyrer, and L. Grimaldo. 2011. The
10 Spawning Migration of Delta Smelt in the Upper San Francisco Estuary.
11 *San Francisco Estuary and Watershed Science* 9(2):1–16.
- 12 Stevens, D. E., and L. W. Miller. 1983. Effects of river flow on abundance of
13 young Chinook salmon, American shad, Longfin Smelt, and delta smelt in
14 the Sacramento–San Joaquin river system. *North American Journal of*
15 *Fisheries Management* 3:425-437.
- 16 Stillwater Sciences. 2006. *Napa River fisheries monitoring program*. Final report.
17 (Contract DACW05-01-C-0015.). Prepared by Stillwater Sciences, Davis,
18 California, for U.S. Army Corps of Engineers, Sacramento District,
19 Sacramento, California.
- 20 Swanson, C., and J. J. Cech, Jr. 1995. *Environmental tolerances and requirements*
21 *of the delta smelt, Hypomesus transpacificus*. Final report. Department of
22 Wildlife, Fish and Conservation Biology, University of California, Davis.
23 As cited by the U.S. Army Corps of Engineers and The Reclamation
24 Board *Standard Assessment Methodology for the Sacramento River Bank*
25 *Protection Project*, August 2004.
- 26 Swanson, C., P. S. Young, and J. J. Cech, Jr. 1998. Swimming performance and
27 behavior of delta smelt: maximum performance and behavioral and
28 kinematic limitations of swimming at submaximal velocities. *Journal of*
29 *Experimental Biology* 201: 333–345.
- 30 Swanson, C., T. Reid, P. S. Young, and J. J. Cech, Jr. 2000. Comparative
31 environmental tolerances of threatened delta smelt (*Hypomesus*
32 *transpacificus*) and introduced wakasagi (*H. nipponensis*) in an altered
33 California estuary. *Oecologia* 123: 384–390.
- 34 Sweetnam, D. A., and D. E. Stevens. 1993. *Report to the Fish and Game*
35 *Commission: a status review of the delta smelt (Hypomesus*
36 *transpacificus) in California*. Candidate Species Status Report 93-DS.
37 California Department of Fish and Game.
- 38 Thomson, J. R., W. J. Kimmerer, L. Brown, K. B. Newman, R. Mac Nally, W. A.
39 Bennett, F. Feyrer, and E. Fleishman. 2010. Bayesian Change-Point
40 Analysis of Abundance Trends for Pelagic Fishes in the Upper
41 San Francisco Estuary. *Ecological Applications* 20:1431–1448.

- 1 USFWS (U.S. Fish and Wildlife Service). 1993. *Endangered and Threatened*
2 *Wildlife and Plants; Determination of Threatened Status for the Delta*
3 *Smelt*. Federal Register 58: 12854.
- 4 USFWS. 1994. *Endangered and Threatened Wildlife and Plants; Critical Habitat*
5 *Determination for the Delta Smelt*. Federal Register 59: 65256.
- 6 USFWS. 1996. *Recovery plan for the Sacramento-San Joaquin Delta native*
7 *fishes*. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon.
8 Available at:
9 http://www.ecos.fws.gov/docs/recovery_plans/1996/961126.pdf.
- 10 USFWS. 2001. *Final biological opinion on the Sacramento River Bank*
11 *Protection Project on the lower Sacramento River in Solano, Sacramento,*
12 *Yolo, Sutter, Colusa, Glenn, Butte, and Tehama counties, California.*
13 Revised File Number 1-1-00-F-0126. Sacramento, California.
- 14 USFWS. 2004. Five Year Status Review for the Delta Smelt. Sacramento, CA.
- 15 USFWS. 2010. Five Year Status Review for the Delta Smelt. Sacramento, CA.
- 16 USFWS. 2012. Technical Staff Comments to the State Water Resources Control
17 Board re: the Comprehensive (Phase 2) Review and Update to the Bay-
18 Delta Plan. Written comments in response to the questions posed by the
19 State Water Resources Control Board (Board) for discussion at the low-
20 salinity zone and pelagic fish workshops that support the Comprehensive
21 (Phase 2) Review and Update to the Bay-Delta Plan. Dated August 17,
22 2012.
- 23 Wang, J. C. S. 1986. *Fishes of the Sacramento-San Joaquin estuary and adjacent*
24 *waters, California: a guide to the early life histories*. Technical Report 9.
25 Prepared for Interagency Ecological Study Program for the Sacramento-
26 San Joaquin Estuary by California Department of Water Resources,
27 California Department of Fish and Game, U.S. Bureau of Reclamation,
28 and U.S. Fish and Wildlife Service.
- 29 Winternitz, L., and K. Wadsworth. 1997. 1996 Temperature trends and potential
30 impacts to salmon, delta smelt, and splittail. *Interagency Ecological*
31 *Program for the Sacramento-San Joaquin Estuary Newsletter* 10: 14–17.

32 **9B.11 Longfin Smelt (*Spirinchus thaleichthys*)**

33 **9B.11.1 Legal Status**

34 Federal: Candidate for listing as Endangered

35 State: Threatened

36 Longfin Smelt is a state-listed threatened species throughout its range in
37 California (DFG 2009). USFWS denied a petition for Federal listing because the
38 population in California (and specifically the San Francisco Bay) was not
39 believed to be sufficiently genetically isolated from other populations (USFWS
40 2009). The Center for Biological Diversity challenged the merits of this

1 determination. In 2011, USFWS entered into a settlement agreement with the
2 Center for Biological Diversity and agreed to conduct a rangewide status review
3 and prepare a 12-month finding to be published by September 30, 2011. The
4 12-month finding on the petition to list the San Francisco Bay-Delta population of
5 the Longfin Smelt as endangered or threatened was completed in March 2012.
6 USFWS determined that listing the Longfin Smelt rangewide was not warranted
7 at the time, but that listing the Bay-Delta DPS of Longfin Smelt was warranted
8 but precluded by other higher priority listing actions (USFWS 2012).

9 **9B.11.2 Distribution**

10 Populations of the Longfin Smelt have been found in estuaries along the Pacific
11 coast from Prince William Sound, Alaska, to the Sacramento-San Joaquin estuary
12 (USFWS 2012). The largest population occupies the Sacramento-San Joaquin
13 estuary, with a smaller population in Humboldt Bay and the Eel River (Moyle
14 2002). They may occur throughout the year in the estuary and lowest reaches of
15 the Klamath River, but little is known of this population.

16 Merz et al. (2013) utilized recently available sampling data (~1959-2012) from
17 the Interagency Ecological Program and regional monitoring programs to provide
18 a comprehensive description of the range and temporal and geographic
19 distribution of Longfin Smelt (*Spirinchus thaleichthys*) by life stage within the
20 San Francisco Estuary. Observations occurred as far west as Tiburon in Central
21 San Francisco Bay and south as far as the Dumbarton Bridge in South San
22 Francisco Bay; north as far as the town of Colusa on the Sacramento River and
23 east as far as Lathrop on the San Joaquin River. Longfin smelt were also observed
24 in seasonally-inundated habitat of the Yolo Bypass and in tributaries like the Napa
25 and Petaluma rivers, Cache Slough, and the Mokelumne River (Merz et al. 2013).

26 **9B.11.3 Life History and Habitat Requirements**

27 Longfin Smelt typically live in bays and estuaries and make seasonal migrations.
28 During winter, they congregate for spawning in the upper reaches of the bays and
29 lower reaches of the river deltas. Juvenile and adult Longfin Smelt have been
30 found throughout the year in salinities ranging from pure fresh water to pure
31 seawater, although once past the juvenile stage, they are typically collected in
32 waters with salinities ranging from 14 to 28 ppt (Baxter 1999). Within the Delta,
33 adult Longfin Smelt occupy water at temperatures from 16 to 20°C (61 to 68°F)
34 and spawn in water with temperatures from 5.6 to 14.5°C (41 to 58°F) (Wang
35 1986).

36 Longfin Smelt have been observed in their winter and spring spawning period as
37 far upstream as Isleton in the Sacramento River, Santa Clara shoal in the
38 San Joaquin system, Hog Slough off the South-Fork Mokelumne River, and Old
39 River south of Indian Slough (DFG 2009). Merz et al. (2013) found that adults
40 were frequently detected in the central regions (from Carquinez Straight upstream
41 to the Confluence), adults were also detected relatively frequently upstream of the
42 Sacramento-San Joaquin confluence. Both adult and larval Longfin Smelt were
43 detected relatively frequently upstream of the confluence, unlike the juvenile and
44 subadult life stages, likely indicating that Longfin Smelt spawning habitat extends

1 further upstream into freshwater areas than rearing habitat. Spawning adults
2 appear to be able to disperse into upper Delta reaches and into San Francisco Bay
3 as well. The presence of adult Longfin Smelt in San Francisco Bay during the
4 spawning period likely relates to years with high Delta inflows, when low salinity
5 habitat shifted westward (Merz et al. 2013). Exact spawning locations in the
6 Delta are unknown and may vary from year to year, depending on environmental
7 conditions. However, it seems likely that spawning locations consist of the
8 overlap of appropriate conditions of flow, temperature, and salinity with
9 appropriate substrate (Rosenfield 2010). Most individuals die after spawning, but
10 occasionally a female may live to spawn a second time.

11 Longfin Smelt congregate in deep waters near the low salinity zone near X2
12 during the spawning period, and they likely make short runs upstream, possibly at
13 night, to spawn from these locations (DFG 2009, Rosenfield 2010). Longfin
14 Smelt in the Delta may spawn as early as November and as late as June, although
15 spawning typically occurs from January to April (DFG 2009, Moyle 2002). The
16 adhesive eggs are deposited on rocks or aquatic plants in the freshwater sections
17 of bays and river deltas. Baxter et al. (2010) found that female Longfin Smelt
18 produced between 1,900 and 18,000 eggs, with fecundity greater in fish with
19 greater lengths.

20 Larval Longfin Smelt less than 12 mm (0.5 inch) in length are buoyant because
21 they have not yet developed an air bladder; as a result, they occupy the upper one-
22 third of the water column. Longfin Smelt develop an air bladder at approximately
23 12 to 15 mm (0.5 to 0.6 inch) in length and are able to migrate vertically in the
24 water column. At this time, they shift habitat and live in the bottom two-thirds of
25 the water column (DFG 2009). Longfin Smelt are dispersed broadly in the Delta
26 by high flows and currents, which facilitate transport of larvae and juveniles long
27 distances. Longfin Smelt larvae are dispersed farther downstream during high
28 freshwater flows (Dege and Brown 2004). Longfin Smelt larvae were detected
29 relatively frequently upstream of the Sacramento-San Joaquin confluence; greater
30 than 73 percent of the time in the Lower Sacramento, Upper Sacramento, Cache
31 Slough and Ship Channel, and Lower San Joaquin regions, and greater than 31
32 percent of the time in the East Delta and South Delta regions during the smelt
33 larval surveys (Merz et al. 2013).

34 Longfin Smelt spend approximately 21 months of their 24-month life cycle in
35 brackish or marine waters (Baxter 1999, Dege and Brown 2004). In the Bay-
36 Delta, most Longfin Smelt spend their first year in Suisun Bay and Marsh. The
37 remainder of their life is spent in the San Francisco Bay or the Gulf of Farallones
38 (Moyle 2008). Based on monthly survey results, Rosenfield and Baxter (2007)
39 inferred that the majority of Longfin Smelt from the Bay-Delta migrate out of the
40 estuary after the first winter of their life cycle and return during late fall to winter
41 of their second year. They noted that migration out of the estuary into nearby
42 coastal waters is consistent with captures of Longfin Smelt in the coastal waters
43 of the Gulf of Farallones and hypothesized that the movement is a behavioral
44 response to warm water temperatures during summer and early fall in the
45 shallows of south San Francisco Bay and San Pablo Bay. Some Longfin Smelt

1 may stay in the ocean and not re-enter fresh water to spawn until the end of their
2 third year.

3 In the Bay-Delta, calanoid copepods such as *Pseudodiaptomus forbesi* and
4 *Eurytemora* sp., as well as the cyclopoid copepod *Acanthocyclops vernalis*, are the
5 primary prey of Longfin Smelt during the first few months of their lives
6 (approximately January through May) (Slater 2008). The Longfin Smelt's diet
7 shifts to include mysids such as opossum shrimp (*Neomysis mercedis*) and other
8 small crustaceans (*Acanthomysis* sp.) as soon as they are large enough (20 to
9 30 mm [0.78 to 1.18 inches]) to consume these larger prey items (DFG 2009).

10 Longfin Smelt numbers in the Bay-Delta have declined significantly since the
11 1980s (Rosenfield and Baxter 2007, Baxter et al. 2010). Rosenfield and Baxter
12 (2007) confirmed the positive correlation between Longfin Smelt abundance and
13 freshwater flow that had been previously documented by others (Stevens and
14 Miller 1983, Baxter 1999, Kimmerer 2002), noting that abundances of both adults
15 and juveniles were significantly lower during the 1987–94 drought than during
16 either the pre- or post-drought periods. Abundance of Longfin Smelt has
17 remained low since 2000, even though freshwater flows increased during several
18 of these years (Baxter et al. 2010). Abundance indices derived from the FMWT,
19 Bay Study Midwater Trawl, and Bay Study Otter Trawl show marked declines in
20 Longfin Smelt populations from 2002 to 2009. Longfin Smelt abundance over
21 the last decade is the lowest recorded in the 40-year history of DFG's FMWT
22 monitoring surveys (USFWS 2012).

23 Research on declines of Longfin Smelt and other pelagic fish species in the
24 Bay-Delta since 2002 (referred to as pelagic organism decline) have most recently
25 been summarized in the Interagency Ecological Program 2010 Pelagic Organism
26 Decline Work Plan and Synthesis of Results (Baxter et al. 2010). Although there
27 is substantial uncertainty about the causal mechanisms underlying the pelagic
28 organism decline, reduced Delta freshwater flows have been identified as one of
29 several key factors believed to contribute to recent declines in the abundance of
30 Longfin Smelt (Baxter et al. 2010).

31 **9B.11.4 References**

32 Baxter, R. D. 1999. Osmeridae. Pages 179-216 in J. Orsi, editor. Report on the
33 1980–1995 fish, shrimp, and crab sampling in the San Francisco Estuary,
34 California. Technical Report 63. Interagency Ecological Program.
35 California Department of Fish and Game, Stockton, USA. Available at:
36 http://www.bepress.com/archive/orsi_1999.

37 Baxter, R. D., R. Breuer, L. R. Brown, L. Conrad, F. Feyer, S. Fong, K. Gehrts, L.
38 Grimaldo, B. Herbold, P. Hrodey, A. Mueller-Solger, T. Sommer, and K.
39 Souza. 2010. Interagency Ecological Program 2010 Pelagic Organism
40 Decline Work Plan and Synthesis of Results. Interagency Ecological
41 Program for the San Francisco Estuary.

- 1 Dege, M. and L. R. Brown. 2004. Effect of outflow on spring and summertime
2 distribution and abundance of larval and juvenile fishes in the upper
3 San Francisco estuary. In: F. Feyrer, L. R. Brown, R. L. Brown, and J. J.
4 Orsi (eds.), *Early Life History of Fishes in the San Francisco Estuary and*
5 *Watershed*. American Fisheries Society Symposium 39:49–66.
- 6 DFG (California Department of Fish and Game). 2009. *A status review of the*
7 *longfin smelt (Spirinchus thaleichthys) in California*. Report to California
8 Fish and Game Commission.
- 9 Kimmerer, W. J. 2002. Effects of freshwater flow on abundance of estuarine
10 organisms: physical effects or trophic linkages. *Marine Ecology Progress*
11 *Series* 243:39-55.
- 12 Merz, J.E., P.S. Bergman, J.F. Melgo, and S. Hamilton. 2013. Longfin smelt:
13 spatial dynamics and ontogeny in the San Francisco estuary, California.
14 *California Fish and Game*, 99(3), pp. 122-148.
- 15 Moyle, P. B. 2002. *Inland fishes of California*. Revised edition. University of
16 California Press, Berkeley, California.
- 17 Moyle, Peter B. 2008. *The Future of Fish in Response to Large-Scale Change in*
18 *the San Francisco Estuary, California*. American Fisheries Society
19 Symposium 64:000–000.
- 20 Rosenfield, J.A. 2010. Life history conceptual model and sub-models for longfin
21 smelt, San Francisco Estuary population. Report for Delta Regional
22 Ecosystem Restoration Implementation Plan. California Department of
23 Fish and Wildlife, Sacramento, CA. Available at:
24 <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=28421>.
- 25 Rosenfield, Jonathan A., and Randall D. Baxter. 2007. Population Dynamics and
26 Distribution Patterns of Longfin Smelt in the San Francisco Estuary.
27 *Transactions of the American Fisheries Society* 136:1577–1592. DOI:
28 10.1577/T06-148.1.
- 29 Slater, Steven B. 2008. Feeding Habits of Longfin Smelt in the Upper
30 San Francisco Estuary. Longfin Smelt Diet Poster. California Department
31 of Fish and Game, Stockton, California.
- 32 Stevens, Donald E., and Lee W. Miller. 1983. Effects of River Flow on
33 Abundance of Young Chinook Salmon, American Shad, Longfin Smelt,
34 and Delta Smelt in the Sacramento-San Joaquin River System. *North*
35 *American Journal of Fisheries Management* 3:425-437.
- 36 USFWS (U.S. Fish and Wildlife Service). 2009. Endangered and threatened
37 wildlife and plants; 12-month finding on a petition to list the
38 San Francisco Bay-Delta population of the (*Spirinchus thaleichthys*) as
39 endangered. *Federal Register* 74: 16169-16175.
- 40 _____. 2012. Endangered and threatened wildlife and plants; 12-month finding on
41 a petition to list the San Francisco Bay-Delta population of the longfin
42 smelt as endangered or threatened. *Federal Register* 77: 19756.

- 1 Wang, Johnson C. S. 1986. *Fishes of the Sacramento-San Joaquin Estuary and*
2 *Adjacent Waters, California: A Guide to the Early Life Histories.*
3 Prepared for the Interagency Ecological Study Program for the
4 Sacramento-San Joaquin Estuary. Technical Report 9. January.

5 **9B.12 Eulachon (*Thaleichthys pacificus*)**

6 **9B.12.1 Legal Status**

- 7 Federal: Threatened
8 State: Species of Special Concern

9 **9B.12.2 Summary**

10 Eulachon are anadromous fish that occur in the lower portions of certain rivers
11 draining into the northeastern Pacific Ocean, ranging from northern California to
12 the southeastern Bering Sea in Bristol Bay, Alaska (Scott and Crossman 1973,
13 Willson et al. 2006).

14 The southern population of Pacific Eulachon consists of populations spawning in
15 rivers south of the Nass River in British Columbia, Canada, to and including the
16 Mad River in California (NMFS 2009). On March 18, 2010, NMFS listed the
17 southern DPS of Pacific Eulachon as threatened under the ESA (NMFS 2010);
18 critical habitat was designated in 2011 (NMFS 2011). The Klamath River is near
19 the southern limit of the range of Eulachon (Eulachon BRT 2010).

20 Spawning occurs in gravel riffles, with hatching about a month later. The larvae
21 generally move downstream to the estuary following hatching.

22 Large spawning aggregations of Pacific Eulachon used to regularly occur in the
23 Klamath River (Fry 1979), migrating in March and April to spawn, but they rarely
24 moved more than 8 miles inland (NRC 2004). DFW sampled in the Klamath
25 River from 1989 to 2003 with no Pacific Eulachon captures (USDI and DFG
26 2011). The Yurok Tribe sampled extensively for Pacific Eulachon in early 2011,
27 and although tribal fishermen did not capture Pacific Eulachon from the Klamath
28 River itself, they did recover Pacific Eulachon from the surf zone at the mouth of
29 the river (USDI and DFG 2011).

30 **9B.12.3 References**

- 31 Eulachon BRT (Eulachon Biological Review Team). 2010. *Status review update*
32 *for eulachon in Washington, Oregon, and California.*
33 [http://www.nwr.noaa.gov/Other-Marine-Species/upload/eulachon-review-](http://www.nwr.noaa.gov/Other-Marine-Species/upload/eulachon-review-update.pdf)
34 [update.pdf.](http://www.nwr.noaa.gov/Other-Marine-Species/upload/eulachon-review-update.pdf)
- 35 Fry, D. H., Jr. 1979. *Anadromous fishes of California.* California Department of
36 Fish and Game, Sacramento.
- 37 NMFS (National Marine Fisheries Service). 2009. Endangered and threatened
38 wildlife and plants; proposed threatened status for Southern Distinct
39 Population Segment of eulachon. *Federal Register* 75 13012-13024.

- 1 _____ . 2010. Endangered and threatened wildlife and plants; threatened status for
2 Southern Distinct Population Segment of eulachon. *Federal Register* 75
3 13012-13024.
- 4 _____ . 2011. Endangered and threatened species, designation of critical habitat
5 for Southern Distinct Population Segment of eulachon. *Federal Register*
6 76: 515-536.
- 7 NRC (National Research Council). 2004. *Endangered and threatened fishes in the*
8 *Klamath River basin: causes of decline and strategies for recovery*. The
9 National Academies Press, Washington, D.C.
10 <http://www.nap.edu/openbook.php?isbn=0309090970>.
- 11 Scott, W. B., and E. J. Crossman. 1973. Freshwater fishes of Canada. *Fisheries*
12 *Research Board of Canada Bulletin* No. 184.
- 13 USDI and DFG (U. S. Department of the Interior and California Department of
14 Fish and Game). 2012. *Klamath Facilities Removal environmental*
15 *impact statement/ environmental impact report*. State Clearinghouse
16 #2010062060. U.S. Department of the Interior, through the Bureau of
17 Reclamation and California Department of Fish and Game, Sacramento,
18 California.
- 19 Willson, M. F., R. H. Armstrong, M. C. Hermans, and K. Koski. 2006. *Eulachon:*
20 *a review of biology and an annotated bibliography*. AFSC Processed
21 Report 2006-12. National Marine Fisheries Service, Alaska Fisheries
22 Science Center, Juneau.

23 **9B.13 Striped Bass (*Morone saxatilis*)**

24 **9B.13.1 Legal Status**

25 Federal: None

26 State: None

27 Striped Bass are native to the Atlantic Coast of North America and were
28 introduced to California in 1879. Striped Bass are a large (>1 meter), long-lived
29 (>10 years) species. They are widespread in the San Francisco Estuary watershed
30 as juveniles and adults. Striped Bass move regularly from salt to fresh water.
31 They require a large body of water for foraging on fish (usually estuaries or large
32 reservoirs) and large cool rivers for spawning. Striped Bass spend most of their
33 lives in estuaries.

34 **9B.13.2 Distribution in Affected Area**

35 Adult Striped Bass are distributed mainly in the lower bays and ocean during the
36 summer, and in the Delta during fall and winter. Spawning takes place in the
37 spring (April–June), at which time Striped Bass swim upstream to spawning
38 grounds. In the Sacramento River, most spawning takes place between RM 77.7
39 and RM 121.2 (Moyle 2002). After spawning, adults move downstream into the
40 Delta and bays (Blunt 1962).

1 **9B.13.3 Life History and Habitat Requirements**

2 Female Striped Bass mature at between 4 and 6 years of age and can spawn every
3 year. In the Delta and Sacramento and San Joaquin rivers, spawning occurs from
4 April to June at temperatures between 14°C and 21°C. Eggs are free-floating and
5 negatively buoyant, and hatch in about two days as they drift downstream, with
6 larvae occurring in shallow and open waters of the lower reaches of the
7 Sacramento and San Joaquin rivers, the Delta, Suisun Bay, Montezuma Slough,
8 and Carquinez Strait. Location of spawning varies based on temperature, flow,
9 and salinity (Turner 1972). In the Yolo Bypass, Harrell and Sommer (2003)
10 observed that flow pulses immediately preceding floodplain inundation triggered
11 upstream movement of Striped Bass, resulting in successful spawning. During
12 low flow years, spawning occurs within the Delta itself.

13 Newly hatched Striped Bass feed off their yolk sac for up to 8 days (Wang 1986),
14 after which they start feeding on zooplankton. Larvae in the Sacramento River
15 migrate into the water column from April to mid-June (Stevens 1966). In the
16 Sacramento River, embryos and larvae are carried into the Delta and Suisun Bay
17 (Moyle 2002). In the San Joaquin River, embryos remain in the same general
18 area where spawning took place, as freshwater outflow is balanced by tidal
19 currents (Moyle 2002). When larval bass from both rivers begin to feed, they are
20 concentrated in the most productive part of the estuary—where freshwater and
21 salt water meet or near X2 (Moyle 2002).

22 Striped Bass are tolerant of a wide range of environmental conditions, surviving
23 temperatures up to 25°C (77°F) (and up to 34°C [93°F] for shorter periods), rapid
24 temperature swings, low oxygen levels between 3 and 5 milligrams per liter
25 (mg/L), and high turbidity (Moyle 2002). Hassler (1988), in a summary of
26 environmental tolerance studies, reported that Striped Bass could tolerate
27 dissolved oxygen concentrations ranging from 3 to 20 mg/L, and a pH range of
28 6 to 10, although the optimum level ranged from 6 to 12 mg/L and 7 to 9,
29 respectively. The information compiled by Hassler (1988) suggested juveniles
30 preferred rearing temperatures of 24 to 26°C (60.8 to 66.2°F). As Striped Bass
31 grow, their temperature preference shifts towards cooler water (Hill et al. 1989).
32 Adult Striped Bass appear to prefer water temperatures ranging from 20 to 24°C
33 (68 to 75.2°F) (Emmett et al. 1991).

34 Typical of an anadromous species, salinity tolerance of Striped Bass also changes
35 with age (Lal et al. 1977, Hill et al. 1989). Eggs and larvae reportedly thrive at
36 salinities less than 3 practical salinity units (psu) (Mansueti 1958, Dovel 1971),
37 and can tolerate salinities of 8 to 9 psu without ill effects (Morgan and Rasin
38 1973). Adults can apparently tolerate salinities from 0 to 34 psu or more (Rogers
39 and Westin 1978), with a range of 10 to 20 psu reported as optimal for larger
40 juveniles (Bogdanov et al. 1967).

41 **9B.13.4 Biotic Interactions**

42 Striped Bass are pelagic, opportunistic predators, feeding on invertebrates and
43 fishes. They tend to exhibit a roving school foraging strategy (Pickard et al.
44 1982). Larval and juvenile Striped Bass feed on invertebrates such as copepods

1 or opossum shrimp. In the San Francisco Bay area, juvenile bass form small
2 schools or feeding groups (Skinner 1962) with specific prey varying with fish
3 size, habitat, and season (Hill et al. 1989).

4 Striped Bass are a top predator in the Delta and are considered major predators on
5 fish (Thomas 1967). Fish become important in the diet of juveniles when they
6 reach a FL of 130 to 350 mm, especially late in the summer when young-of-the-
7 year Striped Bass and shad become available (Moyle 2002). Striped Bass are
8 primarily piscivorous as subadults, when they reach 250 to 470 mm FL
9 (approximately age 2+). Stevens (1966) found that the importance of fish in the
10 diet of subadult (260 to 470 mm FL) and adult (>380 mm FL) Striped Bass in the
11 Sacramento-San Joaquin estuary varied seasonally. Fish were most prevalent in
12 the diet of subadults in fall, and occurred most frequently in the diet of adults in
13 fall and winter. Adult Striped Bass feed primarily on smaller Striped Bass,
14 threadfin shad, and juvenile salmonids, as well as pelagic ocean fishes (Moyle
15 2002). Striped Bass can successfully switch to feeding on novel prey (Moyle
16 2002). Striped Bass are considered important predators on juvenile salmon in the
17 Sacramento River (Tucker et al. 1998, Moyle 2002). Average populations of
18 1.7 million adults during the late 1960s to early 1970s, and 1.25 million adults
19 during 1967-1991 (USFWS 1995), likely exerted considerable predation pressure
20 on outmigrating juvenile salmon (Yoshiyama et al. 1998). The impact of Striped
21 Bass on Delta Smelt and Sacramento Splittail is not known (Moyle 2002). Delta
22 Smelt were occasional prey fish for Striped Bass in the early 1960s (Turner and
23 Kelley 1966) but went undetected in a recent study of predator stomach contents
24 (Nobriga and Feyrer 2007). Striped Bass are likely the primary predator of
25 juvenile and adult Delta Smelt given their spatial overlap in pelagic habitats
26 (NMFS 2009).

27 Though Striped Bass may commonly exhibit a roving school foraging strategy
28 (Pickard et al. 1982), they appear to take advantage of prey that is concentrated at
29 screened diversions or pumps, and may be partially responsible for the decline of
30 some native fishes, including salmon, thicketail chub, and Sacramento perch
31 (Tucker et al. 1998). Striped Bass are considered to be a primary cause of
32 juvenile salmon mortality at the state water-export facility in the south Delta
33 (USFWS 1995). Tucker et al. (1998) observed Striped Bass preying heavily on
34 juvenile Chinook Salmon that passed through the diversion facilities at Red Bluff
35 Diversion Dam on the Sacramento River. Juvenile Chinook Salmon were found
36 by Thomas (1967) to be a major food item in the diet of Striped Bass in the spring
37 and early summer during smolt outmigration through the Sacramento and
38 San Joaquin rivers and Delta.

39 The introduction of the overbite clam in the 1980s has been associated with large
40 decreases in zooplankton and phytoplankton densities in San Francisco Bay and
41 the western Delta (Carlton et al. 1990), which has decreased the amount of food
42 available for larval and juvenile Striped Bass. The population responses of
43 juvenile Striped Bass to winter-spring outflows changed after the overbite clam
44 invasion as young Striped Bass relative abundance stopped responding to outflow
45 altogether (Sommer et al. 2007). In addition to decreased copepod densities, the

1 principal historic copepod food source, *Eurytemora affinis*, for larval and juvenile
 2 Striped Bass has largely been replaced by alien copepod species that may be
 3 energetically less desirable (Meng and Orsi 1991).

4 Within the Delta, adult Striped Bass feed primarily on Threadfin Shad and
 5 juvenile Striped Bass. Thus, when shortages of alternate prey exist, survival rates
 6 of juvenile bass may decrease as they become increasingly important to adult
 7 diets, resulting in an unusually high response to decreased productivity in the
 8 Delta (Moyle 2002).

9 **9B.13.5 References**

10 Blunt, C. E., Jr. 1962. *Striped Bass*. Delta Fish and Wildlife Protection Study.
 11 Annual Report 1, 61–86. California Department of Fish and Game. As
 12 cited by Environmental Defense Fund *A Focal Species and Ecosystem*
 13 *Functions Approach for Developing Public Trust Flows in the Sacramento*
 14 *and San Joaquin River Delta*, February 2010.

15 Bogdanov, A. S., S. I. Doroshev, and A. F. Karpevich. 1967. Experimental
 16 transfer of *Salmo gairdneri* and *Roccus saxatilis* from the USA for
 17 acclimatization in bodies of water of the USSR. Translated from Russian
 18 by R. M. Howland, Narragansett Marine Game Fish Research Laboratory,
 19 *R. I. Vopr. Ikhtiol* 42: 185–187. As cited Atlantic States Marine Fisheries
 20 Commission *Atlantic Coast Diadromous Fish Habitat, A Review of*
 21 *Utilization Threats, Recommendations for Conservation, and Research*
 22 *Needs, Habitat Management Series #9*, January 2009.

23 Carlton, J. T., J. K. Thompson, L. E. Schemel, and F. H. Nichols. 1990.
 24 Remarkable invasion of San Francisco Bay (California, USA) by the
 25 Asian clam *Potamocorbula amurensis*. I. Introduction and Dispersal.
 26 *Marine Ecology Progress Series* 66: 81-94.

27 Dovel, W. L. 1971. *Fish Eggs and Larvae of the Upper Chesapeake Bay*. Special
 28 Report 4. University of Maryland, Natural Resource Institute. As cited
 29 Atlantic States Marine Fisheries Commission *Atlantic Coast Diadromous*
 30 *Fish Habitat, A Review of Utilization Threats, Recommendations for*
 31 *Conservation, and Research Needs, Habitat Management Series #9*,
 32 January 2009.

33 Emmett, R. L., S. L. Stone, S. A. Hinton, and M. E. Monaco. 1991. *Distribution*
 34 *and Abundance of Fishes and Invertebrates in West Coast Estuaries.*
 35 *Volume 2: Species Life History Summaries*. ELMR Report No. 8.
 36 NOS/NOAA Strategic Environmental Assessment Division, Rockville,
 37 Maryland.

38 Harrell, W. C., and T. R. Sommer. 2003. Patterns of adult fish use on California's
 39 Yolo Bypass floodplain. *California Riparian Systems: Processes and*
 40 *Floodplain Management, Ecology, and Restoration*, pp. 88–93. 2001
 41 Riparian Habitat and Floodplains Conference Proceedings. Edited by P.
 42 M. Faber. Riparian Habitat Joint Venture, Sacramento, California.
 43 http://www.water.ca.gov/aes/docs/HarrellSommer_2003.pdf.

- 1 Hassler, T. J. 1988. *Species Profiles: Life Histories and Environmental*
2 *Requirements of Coastal Fishes and Invertebrates (Pacific Southwest):*
3 *Striped Bass*. Biological Report 82(11.82). U.S Army Corps of Engineers,
4 Vicksburg, Mississippi, and U.S. Fish and Wildlife Service, Washington,
5 DC.
- 6 Hill, J., J. W. Evans, and M. J. Van Den Avyle. 1989. *Species Profiles: Life*
7 *Histories and Environmental Requirements of Coastal Fishes and*
8 *Invertebrates (South Atlantic): Striped Bass*. U.S. Fish and Wildlife
9 Service Biological Report 82(11.118). U.S Army Corps of Engineers.
- 10 Lal, K., R. Lasker, and A. Kuljis. 1977. Acclimation and rearing of striped bass
11 larvae in seawater. *California Fish and Game* 63: 210–218.
- 12 Mansueti, R. 1958. *Eggs, Larvae and Young of the Striped Bass, Roccus saxatilis*.
13 Contribution 112. Maryland Department of Research and Education,
14 Solomans.
- 15 Meng, L., and J. J. Orsi. 1991. Selective predation by larval striped bass on native
16 and introduced copepods. *Transactions of the American Fisheries Society*
17 120: 187–192.
- 18 Morgan, R. P., and V. J. Rasin. 1973. Effects of salinity and temperature on the
19 development of eggs and larvae of striped bass and white perch. Appendix
20 X in *Hydrographic and Ecological Effects of Enlargement of the*
21 *Chesapeake and Delaware Canal*. Final Report DACW-61-71-C-0062.
22 U.S. Army Corps of Engineers, Philadelphia District. As cited by
23 Environmental Defense Fund *A Focal Species and Ecosystem Functions*
24 *Approach for Developing Public Trust Flows in the Sacramento and San*
25 *Joaquin River Delta*, February 2010.
- 26 Moyle, P. B. 2002. *Inland Fishes of California*. Revised edition. University of
27 California Press, Berkeley, California.
- 28 NMFS (National Marine Fisheries Service). 2009. Biological Opinion and
29 Conference Opinion on the Long-term Operations of the Central Valley
30 Project and State Water Project. Southwest Region.
31 [http://www.westcoast.fisheries.noaa.gov/central_valley/water_operations/](http://www.westcoast.fisheries.noaa.gov/central_valley/water_operations/ocap.html)
32 [ocap.html](http://www.westcoast.fisheries.noaa.gov/central_valley/water_operations/ocap.html).
- 33 Nobriga, M. L., and F. Feyrer. 2007. Shallow-water piscivore-prey dynamics in
34 California's Sacramento-San Joaquin Delta. *San Francisco Estuary and*
35 *Watershed Science* 5: Article 4.
- 36 Pickard, A., A. M. Grover, and F. A. Hall, Jr. 1982. *An Evaluation of Predator*
37 *Composition at Three Locations on the Sacramento River*. Technical
38 Report 2. Interagency Ecological Study Program for the Sacramento-
39 San Joaquin Estuary.

- 1 Rogers, B. A., and D. T. Westin. 1978. *A Culture Methodology for Striped Bass*.
2 Report No. 660/3-78-000. U.S. Environmental Protection Agency,
3 Ecological Research Series, Washington D.C. As cited Atlantic States
4 Marine Fisheries Commission *Atlantic Coast Diadromous Fish Habitat, A*
5 *Review of Utilization Threats, Recommendations for Conservation, and*
6 *Research Needs, Habitat Management Series #9*, January 2009.
- 7 Skinner, J. E. 1962. *A Historical Review of the Fish and Wildlife Resources of the*
8 *San Francisco Bay Area*. Report No. 1. California Department of Fish and
9 Game, Water Projects Branch.
- 10 Sommer, T., C. Armor, R. Baxter, R. Breuer, L. Brown, M. Chotkowski, S.
11 Culberson, F. Feyrer, M. Gingras, B. Herbold, W. Kimmerer, A. Mueller-
12 Solger, M. Nobriga, and K. Souza. 2007. The collapse of pelagic fishes in
13 the upper San Francisco Estuary. *Fisheries* 32: 270–277.
- 14 Stevens, D. E. 1966. *Food Habits of Striped Bass, Roccus saxatilis, in the*
15 *Sacramento-San Joaquin Delta*. Ecological Studies of the Sacramento-
16 San Joaquin Delta, Part II, pp. 68–96. Edited by J. L. Turner and D. W.
17 Kelley. Fish Bulletin 136. California Department of Fish and Game.
- 18 Thomas, J. L. 1967. The diet of juvenile and adult striped bass, *Roccus saxatilis*,
19 in the Sacramento-San Joaquin river system. *California Fish and Game*
20 53: 49–62.
- 21 Tucker, M. E., C. M. Williams, and R. R. Johnson. 1998. *Abundance, Food*
22 *Habits, and Life History Aspects of Sacramento Squawfish and Striped*
23 *Bass at the Red Bluff Diversion Complex, California, 1994–1996*. Red
24 Bluff Research Pumping Plant Report No. 4. U.S. Fish and Wildlife
25 Service, Red Bluff, California.
- 26 Turner, J. L. 1972. Striped bass. In *Ecological Studies of the Sacramento-*
27 *San Joaquin Estuary*, pp. 36-43. Edited by J. E. Skinner. California
28 Department of Fish and Game Delta Fish Wildlife Protection Studies
29 Report 8.
- 30 Turner, J. L., and D. W. Kelley. 1966. *Ecological Studies of the Sacramento-*
31 *San Joaquin Delta*. Fish Bulletin 136. California Department of Fish and
32 Game.
- 33 USFWS (U.S. Fish and Wildlife Service). 1995. *Working Paper on Restoration*
34 *Needs: Habitat Restoration Actions to Double Natural Production of*
35 *Anadromous Fish in the Central Valley of California*. Volume 3. Prepared
36 for USFWS under the direction of the Anadromous Fish Restoration
37 Program Core Group, Stockton, California.
- 38 Wang, J. C. S. 1986. *Fishes of the Sacramento-San Joaquin Estuary and Adjacent*
39 *Waters, California: a Guide to the Early Life Histories*. Technical Report
40 9. Prepared for Interagency Ecological Study Program for the Sacramento-
41 San Joaquin Estuary by California Department of Water Resources,
42 California Department of Fish and Game, U.S. Bureau of Reclamation,
43 and U.S. Fish and Wildlife Service.

1 Yoshiyama, R. M., F. W. Fisher, and P. B. Moyle. 1998. Historical abundance
2 and decline of Chinook salmon in the Central Valley region of California.
3 *North American Journal of Fisheries Management* 18: 487–521.

4 **9B.14 Southern Resident Killer Whale (*Orcinus orca*)**

5 **9B.14.1 Legal Status**

6 Federal: Endangered

7 State: None

8 Three distinct forms of Killer Whales, termed residents, transients, and offshores,
9 are recognized in the northeastern Pacific Ocean. Resident Killer Whales in U.S.
10 waters are distributed from Alaska to California, with four distinct communities
11 recognized: Southern, Northern, Southern Alaska, and Western Alaska (Krahn
12 et al. 2002, 2004). Resident Killer Whales are fish eaters and live in stable
13 matrilineal pods. Of these, only the Southern Resident Distinct Population
14 Segment (DPS) is listed as endangered.

15 The designated critical habitat does not overlap with the action area for this
16 consultation, nor are there any discernible changes to the physical environment
17 that occur within designated critical that could be correlated to project operations.
18 The only potential effects of project operations on the identified physical or
19 biological features essential to conservation would be to prey quantity, quality,
20 and availability. Project operations have the potential to affect only a portion of
21 juvenile salmon originating in California's Central Valley streams. As discussed
22 earlier, salmon originating in California streams are estimated to contribute
23 between 3 and 5 percent of the salmon population off the Washington coast based
24 on analysis of troll catches. These estimates were made based on data collected
25 during the time of year when the Southern Residents are present. As discussed
26 above, the majority of the fish attributed to California streams that are affected by
27 the project are expected to be hatchery fish.

28 **9B.14.2 Distribution**

29 The Southern Resident Killer Whale DPS is designated as endangered under the
30 ESA (NMFS 2005). This DPS primarily occurs in the inland waters of
31 Washington state and southern Vancouver Island, particularly during the spring,
32 summer, and fall, but members of the population have been observed off coastal
33 California in Monterey Bay, near the Farallon Islands, and off Point Reyes
34 (Heimlich-Boran 1988, Felleman et al. 1991, Olson 1998, Osborne 1999, NMFS
35 2005). The action area is outside of the DPS's designated Critical Habitat, which
36 is in Washington state (NMFS 2006a).

37 **9B.14.3 Life History and Habitat Requirements**

38 Southern Resident Killer Whales spend a significant portion of the year in the
39 inland waterways of the Strait of Georgia, Strait of Juan de Fuca, and Puget
40 Sound, particularly during the spring, summer, and fall, when all three pods are
41 regularly present in the Georgia Basin (defined as the Georgia Strait, San Juan

1 Islands, and Strait of Juan de Fuca) (Heimlich-Boran 1988, Felleman et al. 1991,
2 Olson 1998, Osborne 1999). The Southern Resident population consists of three
3 pods, identified as J, K, and L pods. Typically, K and L pods arrive in May or
4 June and spend most of their time in this core area until departing in October or
5 November. During this time, both pods also make frequent trips lasting a few
6 days to the outer coasts of Washington and southern Vancouver Island (Ford et al.
7 2000). J pod continues to spend intermittent periods of time in the Georgia Basin
8 and Puget Sound during late fall, winter, and early spring.

9 While the Southern Residents are in inland waters during the warmer months, all
10 of the pods concentrate their activities in Haro Strait, Boundary Passage, the
11 southern Gulf Islands, the eastern end of the Strait of Juan de Fuca, and several
12 localities in the southern Georgia Strait (Heimlich-Boran 1988, Felleman et al.
13 1991, Olson 1998, Ford et al. 2000). In general, they spend less time elsewhere,
14 including other sections of the Georgia Strait, Strait of Juan de Fuca, and San Juan
15 Islands, Admiralty Inlet west of Whidbey Island, and Puget Sound. Individual
16 pods are similar in their preferred areas of use (Olson 1998), although there are
17 some seasonal and temporal differences in certain areas visited by each pod
18 (Hauser 2006). For example, J pod visits Rosario Strait more frequently than K or
19 L pods (Hauser 2006). The movements of Southern Resident Killer Whales relate
20 to those of their preferred prey—salmon. Pods commonly seek out and forage in
21 areas where salmon occur, especially those associated with migrating salmon
22 (Heimlich-Boran 1986, 1988; Nichol and Shackleton 1996). Notable locations of
23 particularly high use include Haro Strait and Boundary Passage, the southern tip
24 of Vancouver Island, Swanson Channel off North Pender Island, and the mouth of
25 the Fraser River delta, which is visited by all three pods in September and
26 October (Felleman et al. 1991, Ford et al. 2000). These sites are major corridors
27 for migrating salmon.

28 Wild female Southern Resident Killer Whales give birth to their first surviving
29 calf between the ages of 12 and 16 years (mean = about 14.9 years) (Olesiuk et al.
30 1990, Matkin et al. 2003). Females produce an average of 5.4 surviving calves
31 during a reproductive life span lasting about 25 years (Olesiuk et al. 1990). Males
32 become sexually mature at body lengths ranging from 5.2 to 6.4 meters, which
33 corresponds to between the ages of 10 and 17.5 years (mean = about 15 years)
34 (Christensen 1984, Perrin and Reilly 1984, Duffield and Miller 1988, Olesiuk
35 et al. 1990), and are presumed to remain sexually active throughout their adult
36 lives (Olesiuk et al. 1990).

37 Southern Resident Killer Whales are known to consume 22 species of fish and
38 one species of squid (Scheffer and Slipp 1948; Ford et al. 1998, 2000; Ford and
39 Ellis 2005; Saulitis et al. 2000). Ford and Ellis (2005) found that salmon
40 represent over 96 percent of the prey consumed during the spring, summer, and
41 fall. Chinook Salmon were selected over other species, comprising over
42 70 percent of the identified salmonids taken. This preference occurred despite the
43 much lower abundance of Chinook in the study area in comparison to other
44 salmonids and is probably related to the species' large size, high fat and energy
45 content, and year-round occurrence in the area. Other salmonids eaten in smaller

1 amounts include chum (22 percent of the diet), pink (3 percent), coho (2 percent),
2 sockeye (less than 1 percent), and steelhead (less than 1 percent) (Ford and Ellis
3 2005). This work suggested an overall preference of these whales for Chinook
4 during the summer and fall, but also revealed extensive feeding on chum salmon
5 in the fall.

6 Southern Resident Killer Whale survival and fecundity are correlated with
7 Chinook Salmon abundance (Ward et al. 2009, Ford et al. 2009). Southern
8 Resident Killer Whales could potentially be affected by changes in salmon
9 populations caused by the Proposed Action, because their survival and fecundity
10 appear dependent on the abundance of Chinook Salmon (Ward et al. 2009, Ford
11 et al. 2009).

12 Chinook Salmon originating from the Fraser River are the dominant prey of
13 resident Killer Whales in the summer months when they are usually in inland
14 marine waters (Hanson et al. 2010). Less is known of their diet during the
15 remainder of the year (September through May), when they spend much of their
16 time in outer coastal waters, and may range from central California to northern
17 British Columbia (Hanson et al. 2010). However, it is believed likely that they
18 preferentially feed on Chinook Salmon when available, and roughly in proportion
19 to their relative abundance (Hanson et al. 2010). Hanson et al. (2010) found
20 Southern Resident stomachs to contain several different ESUs of salmon,
21 including Central Valley fall-run Chinook Salmon.

22 NMFS (2008) estimated the biological requirements of Southern Resident Killer
23 Whales including the diet composition and number of salmon the population
24 requires in their coastal range. NMFS estimated that the current population of
25 Southern Residents at the time (87) would be required to consume between
26 392,555 and 470,288 salmon based on diet compositions and bioenergetic needs
27 in their coastal range. These estimates were based on Chinook Salmon
28 comprising 70 to 88 percent of their diet.

29 Salmon originating in California streams are estimated to contribute 3 percent of
30 the salmon population off the Washington coast based on genetic stock
31 identification (GSI) of Washington troll catch in May of 1981 and 1982 (Utter
32 et al. 1983). Research in the mid-1970s estimated California's contribution at
33 5 percent (Wright 1976). More recent data from Collaborative Research on
34 Oregon Ocean Salmon using GSI estimate that 59 percent of salmon analyzed
35 from the Oregon commercial harvest (June–October 2006) were Central Valley
36 fall-run or spring-run Chinook Salmon (<https://fp.pacificfishtrax.org/portal/>). It is
37 important to note that these percentages could vary during different years or
38 seasons.

39 Reclamation funds the operation and maintenance of the Coleman, Livingstone,
40 and Nimbus hatcheries. These hatcheries have a combined yearly production goal
41 of 17,200,000 Chinook Salmon smolts. DWR funds the operation of the Feather
42 River hatcheries for production of approximately 8 million Chinook Salmon
43 smolts annually (yearly production goal).

1 Analysis of Chinook Salmon otoliths in 1999 and 2002 found that the contribution
2 of hatchery-produced fish (from the Sacramento and San Joaquin river system)
3 made up approximately 90 percent of the ocean fishery off the central California
4 coast from Bodega Bay to Monterey Bay (Barnett-Johnson et al. 2007). Similar
5 studies have not been completed to assess the percentage that Central Valley
6 hatcheries contribute to the salmon originating from California off the Oregon and
7 Washington coasts, but it suggests that hatchery fish would likely be the majority.

8 Based on observations of captive Killer Whales, studies have extrapolated the
9 energy requirements of wild Killer Whales and estimate an average size value for
10 the five salmon species combined. Osborne (1999) estimated that adult Killer
11 Whales would consume 28 to 34 adult salmon per day, and that younger Killer
12 Whales (less than 13 years of age) would consume about 15 to 17 salmon per day
13 to meet their daily energy requirements. Extrapolating these results, the Southern
14 Resident population (approximately 90 individuals) would consume about
15 750,000 to 850,000 adult salmon per year.

16 **9B.14.4 Population Trends**

17 Some evidence suggests that until the mid- to late-1800s, the Southern Resident
18 Killer Whale population may have numbered more than 200 animals (Krahn et al.
19 2002). This estimate was based, in part, on a recent genetic analysis of
20 microsatellite DNA, which found that the genetic diversity of the Southern
21 Resident population resembles that of the Northern Residents (Barrett-Lennard
22 2000, Barrett-Lennard and Ellis 2001), and concluded that the two populations
23 were possibly once similar in size. Recent efforts to assess the Killer Whale
24 population during the past century have been hindered by an absence of empirical
25 information prior to 1974 (NMFS 2006b). For example, a report by Scheffer and
26 Slipp (1948) is the only pre-1974 account of Southern Resident abundance in the
27 area, and it merely noted that the species was “frequently seen” during the 1940s
28 in the Strait of Juan de Fuca, northern Puget Sound, and off the coast of the
29 Olympic Peninsula, with smaller numbers along Washington’s outer coast.
30 Olesiuk et al. (1990) estimated the Southern Resident population size in 1967 to
31 be 96 animals. At about this time, marine mammals became popular attractions in
32 zoos and marine parks, which increased the demand for interesting and exotic
33 display animals. Between 1967 and 1973, it is estimated that 47 Killer Whales,
34 mostly immature, were taken from the Southern Resident population for public
35 display. The rapid removal of individual whales caused an immediate decline in
36 numbers (Ford et al. 2000). By 1971, the level of removal decreased the
37 population by about 30 percent, to approximately 67 whales (Olesiuk et al. 1990).
38 In 1993, two decades after the live capture of Killer Whales ended, the three
39 Southern Resident pods—J, K, and L—totaled 96 animals (Ford et al. 2000).

40 Over the past decade, the Southern Resident population has fluctuated. For
41 example, the population appeared to experience a period of recovery by
42 increasing to 99 whales in 1995, but then declined by 20 percent to 79 whales in
43 2001 (-3.3 percent per year) before another slight increase to 83 whales in 2003
44 (Ford et al. 2000, Carretta et al. 2004). NMFS (2008) estimated the 2007
45 population to be 87 whales. The population estimate in 2006 was approximately

1 90 animals (+3.5 percent per year since 2001); the decline in the 1990s, unstable
 2 population status, and population structure (e.g., few reproductive age males and
 3 non-calving adult females) continue to be causes for concern. Moreover, it is
 4 unclear whether the recent increasing trend will continue because these
 5 observations may represent an anomaly in the general pattern of survival or a
 6 longer-term shift in the survival pattern.

7 **9B.14.5 References**

- 8 Barnett-Johnson, R., C. B. Grimes, C. F. Royer, and C. J. Donohoe. 2007.
 9 Identifying the contribution of wild and hatchery Chinook salmon
 10 (*Oncorhynchus tshawytscha*) to the ocean fishery using otolith
 11 microstructure as natural tags. *Canadian Journal of Fisheries and Aquatic
 12 Sciences* 64: 1683-1692.
- 13 Barrett-Lennard, L. G. 2000. *Population Structure and Mating Patterns of Killer
 14 Whales as Revealed by DNA Analysis*. Doctoral dissertation. University of
 15 British Columbia, Vancouver, B.C.
- 16 Barrett-Lennard, L. G., and G. M. Ellis. 2001. *Population Structure and Genetic
 17 Variability in Northeastern Pacific Killer Whales: Towards an
 18 Assessment of Population Viability*. Research Document 2001/065.
 19 Department of Fisheries and Oceans Canada, Nanaimo, British Columbia.
- 20 Carretta, J. V., K. A. Forney, M. M. Muto, J. Barlow, J. Baker, and M. Lowry.
 21 2004. *U.S. Pacific Marine Mammal Stock Assessments: 2003*. NOAA-
 22 TM-NMFS-SWFSC-358. National Marine Fisheries Service.
- 23 Christensen, I. 1984. Growth and reproduction of killer whales, *Orcinus orca*, in
 24 Norwegian coastal waters. *Reports of the International Whaling
 25 Commission (Special Issue)* 6: 253–258.
- 26 Duffield, D. A., and K. W. Miller. 1988. Demographic features of killer whales in
 27 oceanaria in the United States and Canada, 1965-1987. In *North Atlantic
 28 Killer Whales*, pp. 297-306. Edited by J. Sigurjónsson, and S.
 29 Leatherwood. Workshop on North Atlantic Killer Whales. A special issue
 30 of *Journal of the Marine Research Institute Reykjavik* 11. As cited in
 31 <http://www.orcahome.de/growthrate.htm>.
- 32 Felleman, F. L., J. R. Heimlich-Boran, R. W. Osborne. 1991. Feeding ecology of
 33 the killer whale (*Orcinus orca*). In *Dolphin Societies*, pp. 113-147. Edited
 34 by K. Pryor and K. S. Norris. University of California Press, Berkeley.
- 35 Ford, J. K. B., and G. M. Ellis. 2005. *Prey Selection and Food Sharing by Fish-
 36 eating Resident Killer Whales (Orcinus orca) in British Columbia*.
 37 Canadian Science Advisory Secretariat Research Document 2005/041.
- 38 Ford, J. K. B., G. M. Ellis, L. G. Barrett-Lennard, A. B. Morton, R. S. Palm, and
 39 K. C. Balcomb, III. 1998. Dietary specialization in two sympatric
 40 populations of killer whales (*Orcinus orca*) in coastal British Columbia
 41 and adjacent waters. *Canadian Journal of Zoology* 76: 1456-1471.

- 1 Ford, J. K. B., G. M. Ellis, and K. C. Balcomb. 2000. *Killer Whales: the Natural*
2 *History and Genealogy of Orcinus orca in British Columbia and*
3 *Washington State*. Second edition. UBC Press, Vancouver, British
4 Columbia.
- 5 Ford, J. K. B., G. M. Ellis, P. F. Olesiuk, K. C. Balcomb, III. 2010. Linking killer
6 whale survival and prey abundance: food limitations in the oceans' apex
7 predator? *Biology Letters* doi:10.1098/rsbl.2009.0468.
- 8 Hanson, M. B., R. W. Baird, J. K. B. Ford, J. Hempelmann-Halos, D. M. Van
9 Doornik, J. R. Candy, C. K. Emmons, G. S. Schorr, B. Gisborne, K. L.
10 Ayres, S. K. Wasser, K. C. Balcomb, K. Balcomb-Bartok, J. G. Sneva, and
11 M. J. Ford. 2010. Species and stock identification of prey consumed by
12 endangered Southern Resident Killer Whales in their summer range.
13 *Endangered Species Research* 11: 69-82.
- 14 Hauser, D. D. W. 2006. *Summer Space Use of Southern Resident Killer Whales*
15 *(Orcinus orca) within Washington and British Columbia Inshore Waters*.
16 Master's thesis. University of Washington, Seattle.
- 17 Heimlich-Boran, J. R. 1988. Behavioral ecology of killer whales (*Orcinus orca*)
18 in the Pacific Northwest. *Canadian Journal of Zoology* 66: 565-578.
- 19 Krahn, M. M., P. R. Wade, S. T. Kalinowski, M. E. Dahlheim, B. L. Taylor, M.
20 B. Hanson, G. M. Ylitalo, R. P. Angliss, J. E. Stein, and R. S. Waples.
21 2002. *Status Review of Southern Resident Killer Whales (Orcinus orca)*
22 *under the Endangered Species Act*. NOAA Technical Memorandum
23 NMFS-NWFSC-54. National Marine Fisheries Service.
- 24 Krahn, M., M. J. Ford, W. F. Perrin, P. R. Wade, R. P. Angliss, M. B. Hanson, B.
25 L. Taylor, G. M. Ylitalo, M. E. Dahlheim, J. E. Stein, and R. S. Waples.
26 2002. *Status Review of Southern Resident Killer Whales (Orcinus orca)*
27 *under the Endangered Species Act*. NOAA Technical Memorandum
28 NMFS-NWFSC-62. National Marine Fisheries Service.
- 29 Matkin, C. O., G. Ellis, L. B. Lennard, H. Yurk, E. Saulitis, D. Scheel, P. Olesiuk,
30 and G. Ylitalo. 2003. *Photographic and Acoustic Monitoring of Killer*
31 *Whales in Prince William Sound and Kenai Fjords*. Exxon Valdez Oil
32 Spill Restoration Project. North Gulf Oceanic Society, Homer, Alaska.
- 33 Nichol, L. M., and D. M. Shackleton, 1996. Seasonal movements and foraging
34 behaviour of northern resident killer whales (*Orcinus orca*) in relation to
35 the inshore distribution of salmon (*Oncorhynchus* spp.) in British
36 Columbia. *Canadian Journal of Zoology* 74: 983-991.
- 37 NMFS (National Marine Fisheries Service). 2005. Endangered and threatened
38 wildlife and plants: endangered status for Southern Resident killer whales.
39 *Federal Register* 70: 69903-69912.
- 40 _____. 2006a. Endangered and threatened species; designation of critical habitat
41 for Southern Resident killer whale. *Federal Register* 71: 69054-69070.

- 1 _____ . 2006b. *Proposed Recovery Plan for Southern Resident Killer Whales*
2 (Orcinus orca). National Marine Fisheries Service, Northwest Region,
3 Seattle, Washington. As cited by *Reclamation Biological assessment on*
4 *the continued long-term operations of the Central Valley Project and the*
5 *State Water Project*, August 2008.
- 6 _____ . 2008. Chinook prey availability and biological requirements in coastal
7 range of Southern Residents, re: Supplemental comprehensive analysis of
8 Southern Resident killer whales. Memorandum to D. R. Lohn, NMFS,
9 from D. D. Darm, NMFS, Northwest Region, Seattle, Washington.
10 April 11.
- 11 Olesiuk, P. F., M. A. Bigg, and G. M. Ellis. 1990. Life history and population
12 dynamics of resident killer whales (*Orcinus orca*) in the coastal waters of
13 British Columbia and Washington State. *Rep. International Whaling*
14 *Commission (Special Issue) 12*: 209-244.
- 15 Olson, J. M. 1998. *Temporal and Spatial Distribution Patterns of Sightings of*
16 *Southern Community and Transient Orcas in the Inland Waters of*
17 *Washington and British Columbia*. Master's thesis, Western Washington
18 University, Bellingham. As cited in NMFS 2005.
- 19 Osborne, R. W. 1999. *A Historical Ecology of Salish Sea "Resident" Killer*
20 *Whales (Orcinus orca): with Implications for Management*. Doctoral
21 dissertation. University of Victoria, Victoria, British Columbia.
- 22 Perrin, W. F., and S. B. Reilly. 1984. Reproductive parameters of dolphins and
23 small whales of the family *Delphinidae*. In *Reproduction in Whales,*
24 *Dolphins and Porpoises*, pp. 97-134. Edited by W. F. Perrin, R. L.
25 Brownell Jr., and D. P. DeMaster. International Whaling Commission
26 (Special Issue 6), Cambridge, England.
- 27 Saulitis, E., C. Matkin, L. Barrett-Lennard, K. Heise, and G. Ellis. 2000. Foraging
28 strategies of sympatric killer whale (*Orcinus orca*) populations in Prince
29 William Sound, Alaska. *Marine Mammal Science* 16: 94-109.
- 30 Scheffer, V. B., and J. W. Slipp. 1948. The whales and dolphins of Washington
31 State with a key to the cetaceans of the west coast of North America.
32 *American Midland Naturalist* 39: 257-337.
- 33 Utter, F., D. Teel, and G. Milner. 1983. *Genetic Stock Identification Study, 1981-*
34 *1982*. Final report, Project No. 197900100. Bonneville Power
35 Administration, Portland, Oregon.
- 36 Ward, E. J., E. E. Holmes, and K. C. Balcomb. 2009. Quantifying the effects of
37 prey abundance on killer whale reproduction. *Journal of Applied Ecology*
38 46: 632-640.

- 1 Wright, S. G. 1976. *Status of Washington's Commercial Troll Fishery in the Mid-*
- 2 *1970s*. Technical Report No. 21. Washington Department of Fisheries,
- 3 Olympia. As cited by *Reclamation Biological assessment on the*
- 4 *continued long-term operations of the Central Valley Project and the State*
- 5 *Water Project*, August 2008.

This page left blank intentionally.

1 **Appendix 9C**

2 **Reclamation Salmon Mortality Model**
 3 **Analysis Documentation**

4 This appendix provides information about the methods and assumptions used for
 5 the Coordinated Long-Term Operation of the Central Valley Project (CVP) and
 6 State Water Project (SWP) Environmental Impact Statement (EIS) analysis using
 7 the Bureau of Reclamation (Reclamation) Salmon Mortality Model. It is
 8 organized in two main sections that are briefly described below:

- 9 • Section 9C.1: Reclamation Salmon Mortality Model Methodology and
 10 Assumptions
- 11 – The EIS Salmon Mortality analysis uses the Reclamation Salmon
 12 Mortality model to quantify salmon early life stage (pre-spawned eggs,
 13 fertilized eggs, and pre-emergent fry) losses on the Trinity, Sacramento,
 14 Feather, American, and Stanislaus Rivers. This section briefly describes
 15 the overall analytical approach and assumptions of the Reclamation
 16 Salmon Mortality model.
- 17 • Section 9C.2: Reclamation Salmon Mortality Model Results
- 18 – This section presents the salmon early life stage (pre-spawned eggs,
 19 fertilized eggs, and pre-emergent fry) mortality percentage of Trinity
 20 River Fall-Run, Sacramento River fall-run, late fall-run, spring-run, and
 21 winter-run, Feather River fall-run, American River fall-run, and Stanislaus
 22 River fall-run Chinook Salmon. Statistics are presented in tabular format.

23 **9.C.1 Reclamation Salmon Mortality Model**
 24 **Methodology and Assumptions**

25 **9.C.1.1 Reclamation Salmon Mortality Model Methodology**

26 The Reclamation Salmon Mortality Model simulates the early life stage mortality
 27 of Chinook Salmon along reaches of the Trinity (below Lewiston Dam to Burnt
 28 Ranch), Sacramento (below Keswick Dam to Princeton), Feather (below the Fish
 29 Dam to the Sacramento River confluence), American (below Nimbus Dam to the
 30 Sacramento River confluence), and Stanislaus Rivers (below Goodwin Dam to
 31 Riverbank). The model sets an initial spawning distribution along the different
 32 river reaches (as a percentage) and uses water temperature data to simulate egg
 33 development and mortality based on temperature relationships specified in the
 34 model. Daily water temperature results for the Sacramento, American, and
 35 Stanislaus rivers come from the HEC5Q models; and monthly water temperature
 36 results for the Trinity and Feather rivers come from the Reclamation Temperature
 37 Model are used as an input to Reclamation Salmon Mortality Model. The final
 38 output from the Reclamation Salmon Mortality Model used in this analysis is the
 39 resulting annual percent mortality. Operations Criteria and Plan (OCAP)

1 Biological Assessment (BA) Appendix L (Reclamation 2008) provides detailed
2 description of the Reclamation Salmon Mortality Model structure, assumptions,
3 and processes.

4 **9.C.1.2 Reclamation Salmon Mortality Model Analysis Scenario** 5 **Assumptions**

6 This section describes the assumptions for the Reclamation Salmon Mortality
7 Model analysis for the No Action Alternative, Second Basis of Comparison, and
8 other alternatives.

9 The following CalSim II model simulations were performed as the basis of
10 evaluating the impacts of Alternatives 1 through 5 as compared to the No Action
11 Alternative, and the No Action Alternative and Alternatives 1 through 5 as
12 compared to the Second Basis of Comparison:

- 13 • No Action Alternative
- 14 • Second Basis of Comparison
- 15 • Alternative 1 – for simulation purposes, considered the same as Second Basis
16 of Comparison
- 17 • Alternative 2 – for simulation purposes, considered the same as No Action
18 Alternative
- 19 • Alternative 3
- 20 • Alternative 4 – for simulation purposes, considered the same as Second Basis
21 of Comparison.
- 22 • Alternative 5

23 Assumptions for each of these alternatives were developed with the surface water
24 modeling tools and are described in Appendix 5A, Section B.

25 Alternative 1 modeling assumptions are the same as the Second Basis of
26 Comparison, and Alternative 2 modeling assumptions are the same as the No
27 Action Alternative; therefore, the assumptions for those alternatives are not
28 discussed separately in this document.

29 Assumptions for each of these alternatives are reflected to monthly CalSim II
30 flow data that are used in the HEC5Q and Reclamation Temperature Models to
31 generate flow and water temperature data that are then used in the Reclamation
32 Salmon Mortality Model. Table 9C.1 provides the assumed spawning
33 distributions for fall-, late fall-, winter-, and spring-Run Chinook Salmon on the
34 Sacramento River in simulating various scenarios in this EIS. The OCAP BA
35 Appendix L (Reclamation 2008) Tables L-2 to L-5 provide the assumed spawning
36 distributions for Trinity River, Feather River, American River, and Stanislaus
37 River fall-run Chinook Salmon.

1 **Table 9C.1 Upper Sacramento River Spawning Distributions**

Reach	No.	River Reach	Spawning Distribution (%)			
			Fall	Late Fall	Winter	Spring
UPPER	1	Keswick Dam – ACID Dam	16.28%	67.6%	45.03%	12.43%
	2	ACID Dam – Hwy 44	5.48%	5.0%	42.09%	32.77%
	3	Hwy 44 – Upper Anderson Bridge	12.26%	3.7%	12.23%	27.66%
	4	Upper Anderson Bridge – Balls Ferry	16.19%	7.9%	0.26%	10.90%
	5	Balls Ferry – Jellys Ferry	23.08%	8.0%	0.28%	8.75%
	6	Jellys Ferry – Bend Bridge	6.61%	1.0%	0.06%	2.58%
	7	Bend Bridge – Red Bluff Pumping Plant (previously Red Bluff Diversion Dam)	3.48%	0.5%	0.00%	0.83%
Total – Upper Salmon Reach			83.37%	93.8%	99.95%	95.92%
MIDDLE	8	Red Bluff Pumping Plant – Tehama Bridge	10.82%	3.1%	0.05%	4.08%
	9	Tehama Bridge – Woodson Bridge	3.07%	1.2%	0.00%	0.00%
	10	Woodson Bridge – Hamilton City	1.82%	1.1%	0.00%	0.00%
	Total – Middle Salmon Reach			15.71%	5.4%	0.05%
LOWER	11	Hamilton City – Ord Ferry	0.82%	0.6%	0.00%	0.0%
	12	Ord Ferry – Princeton	0.10%	0.2%	0.00%	0.0%
	Total – Lower Salmon Reach			0.92%	0.8%	0.0%

2 NOTE:

3 Sacramento River salmon spawning distributions were revised based on average
 4 2003-2014 redd survey data, provided by David Swank at National Marine Fisheries
 5 Service in April 2015.

6 **9.C.2 Reclamation Salmon Mortality Model Results**

7 Results are provided for each of the following runs separately:

- 8 • No Action Alternative
- 9 • Second Basis of Comparison
- 10 • Alternative 1
- 11 • Alternative 3
- 12 • Alternative 5

13 In addition, the same statistics are provided for the following comparisons to
 14 establish changes of the alternative with respect to one of the bases of
 15 comparison:

- 16 • Alternative 1 compared to No Action Alternative
- 17 • Alternative 3 compared to No Action Alternative
- 18 • Alternative 5 compared to No Action Alternative

- 1 • No Action Alternative compared to Second Basis of Comparison
- 2 • Alternative 1 compared to Second Basis of Comparison
- 3 • Alternative 3 compared to Second Basis of Comparison
- 4 • Alternative 5 compared to Second Basis of Comparison

5 Model results for Alternatives 1, 4, and Second Basis of Comparison are the
6 same, therefore Alternative 4 results are not presented separately. Model results
7 for Alternative 2 and No Action Alternative are the same, therefore Alternative 2
8 results are not presented separately.

9 The results are provided as tables summarizing the annual losses with long-term
10 averages over the 82-year CalSim II simulation period. Averages are also
11 provided by water year type.

12 The following results are presented in this section:

- 13 • B.1. Sacramento River Percent Salmon Loss Summary – Fall-Run Chinook
14 Salmon
- 15 • B.2. Sacramento River Percent Salmon Loss Summary – Late Fall-Run
16 Chinook Salmon
- 17 • B.3. Sacramento River Percent Salmon Loss Summary – Spring-Run Chinook
18 Salmon
- 19 • B.4. Sacramento River Percent Salmon Loss Summary – Winter-Run Chinook
20 Salmon
- 21 • B.5. Trinity River Percent Salmon Loss Summary – Fall-Run Chinook
22 Salmon
- 23 • B.6. American River Percent Salmon Loss Summary – Fall-Run Chinook
24 Salmon
- 25 • B.7. Feather River Percent Salmon Loss Summary – Fall-Run Chinook
26 Salmon
- 27 • B.8. Stanislaus River Percent Salmon Loss Summary – Fall-Run Chinook
28 Salmon

29 **9.C.3 References**

30 Reclamation (Bureau of Reclamation). 2008. *2008 Central Valley Project and*
31 *State Water Project Operations Criteria and Plan Biological Assessment,*
32 *Appendix L Reclamation Salmon Mortality Model.*

Table B-1. Sacramento River Percent Mortality - Fall-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	17.0	---	-0.1
Wet	10.7	---	-0.8
Above Normal	10.5	---	-1.3
Below Normal	15.3	---	0.1
Dry	17.3	---	-0.1
Critical	37.9	---	2.4
Second Basis of Comparison			
Long-term Average	17.1	0.1	
Wet	11.5	0.8	---
Above Normal	11.9	1.3	---
Below Normal	15.2	-0.1	---
Dry	17.4	0.1	---
Critical	35.5	-2.4	---
Alternative 3			
Long-term Average	16.8	-0.2	-0.3
Wet	11.3	0.6	-0.2
Above Normal	11.6	1.0	-0.3
Below Normal	14.7	-0.7	-0.6
Dry	16.9	-0.4	-0.5
Critical	35.6	-2.3	0.1
Alternative 5			
Long-term Average	16.9	-0.1	-0.2
Wet	10.6	0.0	-0.8
Above Normal	10.4	-0.1	-1.4
Below Normal	15.0	-0.3	-0.2
Dry	17.0	-0.3	-0.5
Critical	38.5	0.6	3.0

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-2. Sacramento River Percent Mortality - Late Fall-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	3.1	---	0.4
Wet	3.1	---	0.8
Above Normal	2.4	---	0.5
Below Normal	2.5	---	-0.1
Dry	2.7	---	0.1
Critical	4.8	---	0.2
Second Basis of Comparison			
Long-term Average	2.7	-0.4	
Wet	2.2	-0.8	---
Above Normal	1.9	-0.5	---
Below Normal	2.6	0.1	---
Dry	2.5	-0.1	---
Critical	4.6	-0.2	---
Alternative 3			
Long-term Average	2.7	-0.4	0.0
Wet	2.3	-0.8	0.0
Above Normal	1.8	-0.6	-0.1
Below Normal	2.6	0.1	0.0
Dry	2.6	-0.1	0.1
Critical	4.6	-0.2	-0.1
Alternative 5			
Long-term Average	3.1	0.0	0.4
Wet	3.0	0.0	0.8
Above Normal	2.4	0.0	0.5
Below Normal	2.4	-0.1	-0.1
Dry	2.7	0.0	0.2
Critical	4.9	0.1	0.2

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-3. Sacramento River Percent Mortality - Spring-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	21.9	---	0.7
Wet	6.3	---	-2.4
Above Normal	4.8	---	-2.4
Below Normal	13.3	---	0.8
Dry	19.4	---	0.7
Critical	84.8	---	10.4
Second Basis of Comparison			
Long-term Average	21.1	-0.7	
Wet	8.6	2.4	---
Above Normal	7.2	2.4	---
Below Normal	12.5	-0.8	---
Dry	18.6	-0.7	---
Critical	74.3	-10.4	---
Alternative 3			
Long-term Average	21.1	-0.7	0.0
Wet	8.4	2.1	-0.3
Above Normal	7.3	2.4	0.0
Below Normal	10.8	-2.5	-1.6
Dry	17.5	-1.9	-1.1
Critical	78.1	-6.6	3.8
Alternative 5			
Long-term Average	21.9	0.1	0.8
Wet	6.3	0.0	-2.4
Above Normal	4.9	0.0	-2.4
Below Normal	13.3	0.0	0.8
Dry	18.1	-1.3	-0.6
Critical	87.4	2.6	13.1

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-4. Sacramento River Percent Mortality - Winter-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	5.0	---	0.7
Wet	0.6	---	-0.1
Above Normal	0.1	---	0.0
Below Normal	0.2	---	-0.8
Dry	0.3	---	0.0
Critical	31.4	---	5.4
Second Basis of Comparison			
Long-term Average	4.3	-0.7	
Wet	0.6	0.1	---
Above Normal	0.1	0.0	---
Below Normal	1.0	0.8	---
Dry	0.3	0.0	---
Critical	26.0	-5.4	---
Alternative 3			
Long-term Average	4.2	-0.8	-0.1
Wet	0.6	0.1	0.0
Above Normal	0.1	0.0	0.0
Below Normal	1.0	0.7	0.0
Dry	0.3	-0.1	0.0
Critical	25.3	-6.0	-0.7
Alternative 5			
Long-term Average	4.6	-0.4	0.3
Wet	0.6	0.0	-0.1
Above Normal	0.1	0.0	0.0
Below Normal	0.3	0.0	-0.8
Dry	0.3	0.0	0.0
Critical	28.9	-2.5	2.9

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-5. Trinity River Percent Mortality - Fall-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	4.0	---	0.2
Wet	1.3	---	-0.6
Above Normal	1.5	---	0.2
Below Normal	3.8	---	0.5
Dry	2.5	---	0.2
Critical	14.8	---	1.8
Second Basis of Comparison			
Long-term Average	3.7	-0.2	
Wet	1.9	0.6	---
Above Normal	1.2	-0.2	---
Below Normal	3.4	-0.5	---
Dry	2.3	-0.2	---
Critical	13.0	-1.8	---
Alternative 3			
Long-term Average	3.7	-0.2	0.0
Wet	1.9	0.5	-0.1
Above Normal	1.2	-0.2	0.0
Below Normal	3.2	-0.6	-0.2
Dry	2.2	-0.3	-0.1
Critical	13.3	-1.5	0.3
Alternative 5			
Long-term Average	3.9	0.0	0.2
Wet	1.3	0.0	-0.6
Above Normal	1.4	0.0	0.2
Below Normal	3.6	-0.2	0.3
Dry	2.5	0.0	0.2
Critical	14.9	0.1	1.9

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-6. American River Percent Mortality - Fall-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	23.2	---	0.2
Wet	22.6	---	-0.6
Above Normal	23.2	---	0.6
Below Normal	23.5	---	2.0
Dry	22.9	---	-0.1
Critical	25.0	---	0.1
Second Basis of Comparison			
Long-term Average	23.1	-0.2	
Wet	23.2	0.6	---
Above Normal	22.7	-0.6	---
Below Normal	21.5	-2.0	---
Dry	23.0	0.1	---
Critical	24.9	-0.1	---
Alternative 3			
Long-term Average	23.2	-0.1	0.1
Wet	23.2	0.6	-0.1
Above Normal	22.6	-0.6	0.0
Below Normal	21.8	-1.7	0.3
Dry	22.9	0.0	-0.1
Critical	25.4	0.4	0.6
Alternative 5			
Long-term Average	23.0	-0.3	-0.1
Wet	22.7	0.1	-0.5
Above Normal	22.5	-0.7	-0.2
Below Normal	22.5	-1.0	1.0
Dry	22.9	0.0	-0.1
Critical	24.7	-0.3	-0.2

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-7. Feather River Percent Mortality - Fall Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	7.2	---	0.2
Wet	4.6	---	2.8
Above Normal	3.4	---	0.2
Below Normal	8.4	---	-0.9
Dry	7.7	---	-0.9
Critical	14.5	---	-3.0
Second Basis of Comparison			
Long-term Average	7.0	-0.2	
Wet	1.7	-2.8	---
Above Normal	3.1	-0.2	---
Below Normal	9.2	0.9	---
Dry	8.6	0.9	---
Critical	17.4	3.0	---
Alternative 3			
Long-term Average	6.0	-1.1	-0.9
Wet	1.9	-2.7	0.1
Above Normal	2.9	-0.4	-0.2
Below Normal	6.8	-1.6	-2.4
Dry	7.8	0.0	-0.8
Critical	14.6	0.2	-2.8
Alternative 5			
Long-term Average	6.9	-0.2	-0.1
Wet	4.5	0.0	2.8
Above Normal	3.2	-0.2	0.1
Below Normal	10.6	2.3	1.4
Dry	7.4	-0.3	-1.1
Critical	13.9	-0.6	-3.6

Notes: All results are based on the 82-year simulation period. The water year types are defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

Table B-8. Stanislaus River Percent Mortality - Fall-Run Chinook Salmon

	Percent Mortality	Difference from No Action Alternative	Difference from Second Basis of Comparison
	%	%	%
No Action Alternative			
Long-term Average	7.0	---	-0.4
Wet	1.6	---	0.1
Above Normal	5.3	---	-0.1
Below Normal	4.4	---	0.3
Dry	4.9	---	-0.3
Critical	14.4	---	-1.5
Second Basis of Comparison			
Long-term Average	7.4	0.4	
Wet	1.5	-0.1	---
Above Normal	5.4	0.1	---
Below Normal	4.1	-0.3	---
Dry	5.1	0.3	---
Critical	15.9	1.5	---
Alternative 3			
Long-term Average	6.2	-0.8	-1.2
Wet	1.6	0.0	0.1
Above Normal	4.0	-1.3	-1.4
Below Normal	3.8	-0.6	-0.3
Dry	4.2	-0.7	-0.9
Critical	13.4	-1.0	-2.5
Alternative 5			
Long-term Average	8.5	1.5	1.0
Wet	1.8	0.2	0.3
Above Normal	6.4	1.1	1.0
Below Normal	6.1	1.6	2.0
Dry	7.0	2.2	1.9
Critical	16.9	2.5	1.0

Notes: All results are based on the 82-year simulation period. The water year types are defined by the San Joaquin Valley 60-20-20 Index Water Year Hydrologic Classification (SWRCB D-1641, 1999); projected to Year 2030.

1 Appendix 9D

2 SALMOD Analysis Documentation

3 This appendix provides information about the methods and assumptions used for
4 the Remanded Biological Opinions on the Coordinated Long-Term Operation of
5 the Central Valley Project (CVP) and State Water Project (SWP) Environmental
6 Impact Statement (EIS) analysis using the SALMOD model. It is organized in
7 two main sections that are briefly described below:

- 8 • Section 9D.1: SALMOD Methodology and Assumptions
 - 9 – The analysis uses the SALMOD model to quantify fall-run, late fall-run,
10 spring-run, and winter-run Chinook Salmon survival and mortality for
11 different life-stages within the Sacramento River, specifically from below
12 Keswick Dam to the Red Bluff Pumping Plant (previously at Red Bluff
13 Diversion Dam). This section briefly describes the overall analytical
14 approach and assumptions of the SALMOD Model.
- 15 • Section 9D.2: SALMOD Model Results
 - 16 – This section presents the production (survival) and mortality by life-stages
17 and various causes of Sacramento River fall-run, late fall-run, spring-run,
18 and winter-run Chinook Salmon. Statistics are presented in exceedance
19 plots and in tabular format.

20 9D.1 SALMOD Methodology and Assumptions

21 9D.1.1 SALMOD Methodology

22 The SALMOD model simulates the life-stage dynamics of fall-run, late fall-run,
23 spring-run, and winter-run Chinook Salmon populations within the Sacramento
24 River, from below Keswick Dam to the Red Bluff Diversion Dam. The model
25 uses daily flow and temperature data from the Sacramento River HEC5Q model
26 to simulate the annual growth, movement, and mortality of the various riverine
27 life stages of the four Chinook Salmon populations based on an initial annual
28 adult population that resets each biological year. The dynamics simulated are
29 based on assumptions and relations specified in the model. The final output from
30 SALMOD used in this analysis is annual production (number of surviving
31 members of each life-stage) and annual mortality based on a variety of factors,
32 including temperature and habitat (flow) based mortality. The 2008 Operations
33 Criteria and Plan (OCAP) Biological Assessment (BA), Appendix P provides
34 detailed description of the SALMOD model structure, assumptions, and processes
35 (Reclamation 2008).

1 **9D.1.2 SALMOD Analysis Scenario Assumptions**

2 This section describes the assumptions for the SALMOD analysis for the
3 No Action Alternative, Second Basis of Comparison, and other alternatives.

4 The following CalSim II model simulations were performed as the basis of
5 evaluating the impacts of the Alternatives 1 through 5 as compared to the No
6 Action Alternative, and the No Action Alternative and Alternatives 1 through 5 as
7 compared to the Second Basis of Comparison:

- 8 • No Action Alternative
- 9 • Second Basis of Comparison
- 10 • Alternative 1 – for simulation purposes, considered the same as Second Basis
11 of Comparison
- 12 • Alternative 2 – for simulation purposes, considered the same as No Action
13 Alternative
- 14 • Alternative 3
- 15 • Alternative 4 – for simulation purposes, considered the same as Second Basis
16 of Comparison.
- 17 • Alternative 5

18 Assumptions for each of these alternatives were developed with the surface water
19 modeling tools and are described in Appendix 5A, Section B.

20 Alternative 1 modeling assumptions are the same as the Second Basis of
21 Comparison, and Alternative 2 modeling assumptions are the same as the
22 No Action Alternative; therefore, the assumptions for those alternatives are not
23 discussed separately in this document.

24 Assumptions for each of these alternatives are reflected in monthly CalSim II
25 flow data that are used in the Sacramento River HEC5Q Model to generate daily
26 flow and temperature data that are input to the SALMOD model. For this
27 analysis, the initial population of adult were assumed to be 23,356 for fall-run,
28 5,545 for late fall-run, 500 for spring-run, and 4,108 for winter-run based on
29 geometric mean of 2003-2014 GrandTab escapement data provided by David
30 Swank at the National Marine Fisheries Service (NMFS) in April 2015. For
31 spring-run, the number of adults in the mainstem Sacramento River are
32 significantly low (arithmetic mean of 69). Based on further discussion with
33 NMFS, 500 adults were assumed as the input in SALMOD. The assumed
34 spawning distribution by reach is shown in Table 9D.1. Assumptions of the
35 spawning distributions were based on average 2003-2014 Redd survey data,
36 provided by David Swank at NMFS in April 2015.

1 **Table 9D.1 Upper Sacramento River Spawning Distributions.**

River Reach	Spawning Distribution (%) Fall	Spawning Distribution (%) Late Fall	Spawning Distribution (%) Spring	Spawning Distribution (%) Winter
Keswick Dam – Anderson Cottonwood Irrigation District (ACID) Dam	19.50	71.30	12.80	45.10
ACID Dam – Highway 44 Bridge	6.60	5.20	33.90	42.10
Highway 44 Bridge – Airport Road Bridge	14.70	3.90	29.70	12.20
Airport Road Bridge – Balls Ferry	19.40	8.90	11.10	0.30
Balls Ferry – Battle Creek	12.50	5.90	7.40	0.10
Battle Creek – Jellys Ferry	15.20	3.10	1.50	0.10
Jellys Ferry – Bend Bridge	8.00	1.20	2.60	0.10
Bend Bridge – Red Bluff Pumping Plant (previously Red Bluff Diversion Dam)	4.20	0.60	0.80	0.00

2 **9D.2 SALMOD Results**

3 Results are provided for each of the following runs separately:

- 4 • No Action Alternative
- 5 • Second Basis of Comparison
- 6 • Alternative 1
- 7 • Alternative 3
- 8 • Alternative 5

9 In addition, the same statistics are provided for the following comparisons to
 10 establish changes of the alternative with respect to one of the bases of
 11 comparison:

- 12 • Alternative 1 compared to No Action Alternative
- 13 • Alternative 3 compared to No Action Alternative
- 14 • Alternative 5 compared to No Action Alternative
- 15 • No Action Alternative compared to Second Basis of Comparison
- 16 • Alternative 1 compared to Second Basis of Comparison
- 17 • Alternative 3 compared to Second Basis of Comparison
- 18 • Alternative 5 compared to Second Basis of Comparison

19 Model results for Alternatives 1, 4, and Second Basis of Comparison are the
 20 same, therefore Alternative 4 results are not presented separately. Model results

1 for Alternative 2 and No Action Alternative are the same, therefore Alternative 2
2 results are not presented separately.

3 The first set of results is provided as probability of exceedance curves of annual
4 production and mortality for the four Sacramento River salmonid populations.
5 For this analysis, exceedance plots for annual production and mortality were
6 generated based on the 82-year CalSim II time period for each of the alternatives
7 and basis of comparison. Differences among alternatives were evaluated using
8 the exceedance probability corresponding to varying levels of survival. The
9 results are provided at the end of this appendix in the following subsections:

- 10 • B.1. Fall-Run Chinook Salmon
- 11 • B.2. Late Fall-Run Chinook Salmon
- 12 • B.3. Spring-Run Chinook Salmon
- 13 • B.4. Winter-Run Chinook Salmon

14 The second set of results is provided as tables summarizing the comparison
15 between alternatives of annual production and mortality with long-term averages
16 over the entire CalSim II simulation period. Averages are also provided by water
17 year type.


18 **9D.3 References**

19 Reclamation (Bureau of Reclamation). 2008. *2008 Central Valley Project and*
20 *State Water Project Operations Criteria and Plan Biological Assessment,*
21 *Appendix P SALMOD Model.*

1 **B.1. Fall-Run Chinook Salmon**


2

Figure B-1-1. Annual Potential Production for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-2. Annual Mortality for Fall-Run Chinook Salmon - Eggs




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-3. Annual Mortality for Fall-Run Chinook Salmon - Fry




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-4. Annual Mortality for Fall-Run Chinook Salmon - Pre-Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-5. Annual Mortality for Fall-Run Chinook Salmon - Immature Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-6. Annual Mortality for Fall-Run Chinook Salmon - Pre- & Immature Smolts




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-7. Annual Mortality for Fall-Run Chinook Salmon - All Lifestages




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-8. Incubation - Habitat based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-9. Super-imposition - Habitat based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-10. Fry - Habitat based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-11. Pre-smolt - Habitat based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-12. Immature Smolt - Habitat based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-13. Total Habitat based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-14. Pre-Spawn Mortality - Temperature based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-15. Eggs - Temperature based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-16. Fry - Temperature based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-17. Pre-smolt - Temperature based Annual Mortality for Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-18. Immature Smolt - Temperature based Annual Mortality for Fall-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-1-19. Total Temperature based Annual Mortality for Fall-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Table B-1-1. Annual Potential Production for Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	16,838,069
Alternative 1	17,037,309
Difference	199,240
Percent Difference ³	1
Water Year Types²	
Wet (32.5%)	
No Action Alternative	16,537,313
Alternative 1	16,525,365
Difference	-11,948
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	15,696,855
Alternative 1	15,746,827
Difference	49,972
Percent Difference	0
Below Normal (17.5%)	
No Action Alternative	17,922,930
Alternative 1	17,847,310
Difference	-75,620
Percent Difference	0
Dry (22.5%)	
No Action Alternative	17,754,135
Alternative 1	17,934,726
Difference	180,590
Percent Difference	1
Critical (15%)	
No Action Alternative	15,800,949
Alternative 1	16,930,799
Difference	1,129,850
Percent Difference	7
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-1-2. Annual Mortality by Life Stage for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	7,894,954	4,684,028	272,676	47,521	320,197
Alternative 1	7,110,950	4,709,109	269,215	49,405	318,621
Difference	-784,003	25,081	-3,461	1,885	-1,576
Percent Difference ³	-10	1	-1	4	0
Water Year Types²					
Wet (32.5%)					
No Action Alternative	6,019,065	5,201,105	74,435	15,865	90,301
Alternative 1	6,023,551	5,129,591	71,744	16,838	88,581
Difference	4,486	-71,514	-2,692	973	-1,719
Percent Difference	0	-1	-4	6	-2
Above Normal (12.5%)					
No Action Alternative	11,831,604	5,007,353	161,828	32,005	193,834
Alternative 1	11,326,553	5,120,441	96,157	31,173	127,329
Difference	-505,051	113,088	-65,672	-833	-66,505
Percent Difference	-4	2	-41	-3	-34
Below Normal (17.5%)					
No Action Alternative	4,975,839	4,911,742	266,079	45,556	311,635
Alternative 1	4,943,736	4,895,243	284,538	50,880	335,418
Difference	-32,103	-16,499	18,459	5,324	23,783
Percent Difference	-1	0	7	12	8
Dry (22.5%)					
No Action Alternative	6,357,019	4,408,740	501,702	61,525	563,227
Alternative 1	5,846,335	4,371,799	440,615	59,727	500,342
Difference	-510,683	-36,940	-61,087	-1,798	-62,885
Percent Difference	-8	-1	-12	-3	-11
Critical (15%)					
No Action Alternative	14,391,374	3,441,525	458,729	110,322	569,051
Alternative 1	10,379,320	3,744,097	566,311	117,959	684,270
Difference	-4,012,054	302,572	107,582	7,638	115,220
Percent Difference	-28	9	23	7	20

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-1-3. Annual Mortality by Cause for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	5,949,693	6,949,486	12,899,179
Alternative 1	5,010,581	7,128,100	12,138,680
Difference	-939,112	178,614	-760,499
Percent Difference ³	-16	3	-6
Water Year Types²			
Wet (32.5%)			
No Action Alternative	927,546	10,382,925	11,310,471
Alternative 1	485,103	10,756,621	11,241,723
Difference	-442,443	373,695	-68,747
Percent Difference	-48	4	-1
Above Normal (12.5%)			
No Action Alternative	11,689,545	5,343,245	17,032,790
Alternative 1	11,136,551	5,437,771	16,574,323
Difference	-552,994	94,526	-458,468
Percent Difference	-5	2	-3
Below Normal (17.5%)			
No Action Alternative	4,200,054	5,999,162	10,199,216
Alternative 1	4,155,751	6,018,646	10,174,397
Difference	-44,304	19,484	-24,819
Percent Difference	-1	0	0
Dry (22.5%)			
No Action Alternative	5,983,150	5,345,836	11,328,986
Alternative 1	5,469,925	5,248,551	10,718,477
Difference	-513,224	-97,285	-610,509
Percent Difference	-9	-2	-5
Critical (15%)			
No Action Alternative	14,038,861	4,363,089	18,401,950
Alternative 1	10,019,091	4,788,596	14,807,687
Difference	-4,019,770	425,507	-3,594,263
Percent Difference	-29	10	-20

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-4. Annual Mortality by Cause and Life Stage for Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	5,139,812	1,955,690	799,452	154	4,683,874	10,275	309,922	12,899,179
Alternative 1	4,292,224	2,108,590	710,136	151	4,708,958	8,069	310,552	12,138,680
Difference	-847,588	152,900	-89,315	-3	25,084	-2,206	630	-760,499
Percent Difference ³	-16	8	-11	-2	1	-21	0	-6
Water Year Types²								
Wet (32.5%)								
No Action Alternative	213,200	5,097,346	708,520	428	5,200,677	5,398	84,903	11,310,471
Alternative 1	76,487	5,544,710	402,355	446	5,129,145	5,816	82,766	11,241,723
Difference	-136,713	447,364	-306,165	18	-71,532	417	-2,137	-68,747
Percent Difference	-64	9	-43	4	-1	8	-3	-1
Above Normal (12.5%)								
No Action Alternative	11,397,132	146,831	287,640	34	5,007,318	4,738	189,095	17,032,790
Alternative 1	10,875,176	194,605	256,772	9	5,120,432	4,595	122,734	16,574,323
Difference	-521,956	47,774	-30,868	-26	113,113	-144	-66,361	-458,468
Percent Difference	-5	33	-11	-74	2	-3	-35	-3
Below Normal (17.5%)								
No Action Alternative	4,050,002	780,040	145,797	60	4,911,682	4,196	307,440	10,199,216
Alternative 1	4,055,314	789,925	98,496	25	4,895,218	1,915	333,503	10,174,397
Difference	5,312	9,886	-47,300	-35	-16,465	-2,280	26,064	-24,819
Percent Difference	0	1	-32	-58	0	-54	8	0
Dry (22.5%)								
No Action Alternative	5,226,978	377,492	752,548	0	4,408,740	3,623	559,604	11,328,986
Alternative 1	4,603,020	378,293	865,023	0	4,371,799	1,883	498,459	10,718,477
Difference	-623,959	801	112,475	0	-36,940	-1,740	-61,145	-610,509
Percent Difference	-12	0	15	0	-1	-48	-11	-5
Critical (15%)								
No Action Alternative	11,740,400	395,039	2,255,935	0	3,441,525	42,525	526,526	18,401,950
Alternative 1	7,750,732	392,537	2,236,052	0	3,744,097	32,307	651,963	14,807,687
Difference	-3,989,668	-2,502	-19,884	0	302,572	-10,218	125,438	-3,594,263
Percent Difference	-34	-1	-1	0	9	-24	24	-20

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-5. Annual Mortality by All Factors for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super-imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
No Action Alternative	5,139,812	1,449,851	505,839	799,452	154	4,683,874	4,419	268,257	5,856	41,665	12,899,179
Alternative 1	4,292,224	1,473,372	635,217	710,136	151	4,708,958	3,312	265,903	4,757	44,648	12,138,680
Difference	-847,588	23,521	129,379	-89,315	-3	25,084	-1,106	-2,354	-1,099	2,984	-760,499
Percent Difference ³	-16	2	26	-11	-2	1	-25	-1	-19	7	-6
Water Year Types²											
Wet (32.5%)											
No Action Alternative	213,200	3,859,065	1,238,281	708,520	428	5,200,677	4,236	70,199	1,162	14,703	11,310,471
Alternative 1	76,487	3,907,496	1,637,214	402,355	446	5,129,145	4,203	67,541	1,613	15,225	11,241,723
Difference	-136,713	48,431	398,933	-306,165	18	-71,532	-33	-2,659	451	522	-68,747
Percent Difference	-64	1	32	-43	4	-1	-1	-4	39	4	-1
Above Normal (12.5%)											
No Action Alternative	11,397,132	67,263	79,569	287,640	34	5,007,318	3,300	158,529	1,438	30,567	17,032,790
Alternative 1	10,875,176	114,650	79,955	256,772	9	5,120,432	3,015	93,141	1,579	29,593	16,574,323
Difference	-521,956	47,387	386	-30,868	-26	113,113	-285	-65,387	141	-974	-458,468
Percent Difference	-5	70	0	-11	-74	2	-9	-41	10	-3	-3
Below Normal (17.5%)											
No Action Alternative	4,050,002	246,033	534,007	145,797	60	4,911,682	2,887	263,192	1,308	44,248	10,199,216
Alternative 1	4,055,314	257,762	532,163	98,496	25	4,895,218	1,115	283,424	801	50,079	10,174,397
Difference	5,312	11,729	-1,844	-47,300	-35	-16,465	-1,773	20,232	-508	5,832	-24,819
Percent Difference	0	5	0	-32	-58	0	-61	8	-39	13	0
Dry (22.5%)											
No Action Alternative	5,226,978	377,492	0	752,548	0	4,408,740	1,403	500,298	2,220	59,306	11,328,986
Alternative 1	4,603,020	378,293	0	865,023	0	4,371,799	423	440,192	1,460	58,267	10,718,477
Difference	-623,959	801	0	112,475	0	-36,940	-980	-60,107	-760	-1,038	-610,509
Percent Difference	-12	0	0	15	0	-1	-70	-12	-34	-2	-5
Critical (15%)											
No Action Alternative	11,740,400	395,039	0	2,255,935	0	3,441,525	12,058	446,671	30,467	79,854	18,401,950
Alternative 1	7,750,732	392,537	0	2,236,052	0	3,744,097	8,529	557,782	23,779	94,181	14,807,687
Difference	-3,989,668	-2,502	0	-19,884	0	302,572	-3,529	111,111	-6,689	14,327	-3,594,263
Percent Difference	-34	-1	0	-1	0	9	-29	25	-22	18	-20

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-6. Annual Potential Production for Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	16,838,069
Alternative 3	17,129,024
Difference	290,955
Percent Difference ³	2
Water Year Types²	
Wet (32.5%)	
No Action Alternative	16,537,313
Alternative 3	16,544,696
Difference	7,383
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	15,696,855
Alternative 3	15,897,563
Difference	200,708
Percent Difference	1
Below Normal (17.5%)	
No Action Alternative	17,922,930
Alternative 3	17,877,415
Difference	-45,515
Percent Difference	0
Dry (22.5%)	
No Action Alternative	17,754,135
Alternative 3	18,382,793
Difference	628,657
Percent Difference	4
Critical (15%)	
No Action Alternative	15,800,949
Alternative 3	16,667,512
Difference	866,563
Percent Difference	5
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-1-7. Annual Mortality by Life Stage for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	7,894,954	4,684,028	272,676	47,521	320,197
Alternative 3	6,873,719	4,709,136	258,786	47,224	306,009
Difference	-1,021,235	25,108	-13,891	-297	-14,187
Percent Difference ³	-13	1	-5	-1	-4
Water Year Types²					
Wet (32.5%)					
No Action Alternative	6,019,065	5,201,105	74,435	15,865	90,301
Alternative 3	5,981,293	5,099,805	75,392	16,365	91,757
Difference	-37,772	-101,300	957	500	1,457
Percent Difference	-1	-2	1	3	2
Above Normal (12.5%)					
No Action Alternative	11,831,604	5,007,353	161,828	32,005	193,834
Alternative 3	10,983,177	5,061,047	110,803	26,403	137,207
Difference	-848,427	53,694	-51,025	-5,602	-56,627
Percent Difference	-7	1	-32	-18	-29
Below Normal (17.5%)					
No Action Alternative	4,975,839	4,911,742	266,079	45,556	311,635
Alternative 3	4,905,579	4,909,824	267,778	50,091	317,869
Difference	-70,260	-1,918	1,699	4,535	6,234
Percent Difference	-1	0	1	10	2
Dry (22.5%)					
No Action Alternative	6,357,019	4,408,740	501,702	61,525	563,227
Alternative 3	4,403,331	4,450,665	464,033	59,943	523,976
Difference	-1,953,687	41,925	-37,668	-1,583	-39,251
Percent Difference	-31	1	-8	-3	-7
Critical (15%)					
No Action Alternative	14,391,374	3,441,525	458,729	110,322	569,051
Alternative 3	11,384,504	3,723,000	461,093	109,012	570,105
Difference	-3,006,871	281,476	2,364	-1,310	1,055
Percent Difference	-21	8	1	-1	0

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-1-8. Annual Mortality by Cause for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	5,949,693	6,949,486	12,899,179
Alternative 3	4,751,566	7,137,299	11,888,865
Difference	-1,198,127	187,813	-1,010,314
Percent Difference ³	-20	3	-8
Water Year Types²			
Wet (32.5%)			
No Action Alternative	927,546	10,382,925	11,310,471
Alternative 3	389,939	10,782,916	11,172,855
Difference	-537,606	399,991	-137,615
Percent Difference	-58	4	-1
Above Normal (12.5%)			
No Action Alternative	11,689,545	5,343,245	17,032,790
Alternative 3	10,788,099	5,393,332	16,181,431
Difference	-901,446	50,087	-851,359
Percent Difference	-8	1	-5
Below Normal (17.5%)			
No Action Alternative	4,200,054	5,999,162	10,199,216
Alternative 3	4,135,609	5,997,663	10,133,272
Difference	-64,445	-1,499	-65,944
Percent Difference	-2	0	-1
Dry (22.5%)			
No Action Alternative	5,983,150	5,345,836	11,328,986
Alternative 3	4,017,083	5,360,888	9,377,972
Difference	-1,966,066	15,053	-1,951,014
Percent Difference	-33	0	-17
Critical (15%)			
No Action Alternative	14,038,861	4,363,089	18,401,950
Alternative 3	10,991,653	4,685,957	15,677,609
Difference	-3,047,208	322,868	-2,724,340
Percent Difference	-22	7	-15

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-9. Annual Mortality by Cause and Life Stage for Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	5,139,812	1,955,690	799,452	154	4,683,874	10,275	309,922	12,899,179
Alternative 3	3,882,019	2,130,887	860,812	146	4,708,991	8,589	297,421	11,888,865
Difference	-1,257,793	175,198	61,360	-8	25,116	-1,686	-12,501	-1,010,314
Percent Difference ³	-24	9	8	-5	1	-16	-4	-8
Water Year Types²								
Wet (32.5%)								
No Action Alternative	213,200	5,097,346	708,520	428	5,200,677	5,398	84,903	11,310,471
Alternative 3	37,613	5,597,671	346,009	441	5,099,364	5,877	85,881	11,172,855
Difference	-175,587	500,325	-362,510	13	-101,313	478	978	-137,615
Percent Difference	-82	10	-51	3	-2	9	1	-1
Above Normal (12.5%)								
No Action Alternative	11,397,132	146,831	287,640	34	5,007,318	4,738	189,095	17,032,790
Alternative 3	10,309,394	196,462	477,321	0	5,061,047	1,384	135,823	16,181,431
Difference	-1,087,738	49,631	189,681	-34	53,729	-3,354	-53,273	-851,359
Percent Difference	-10	34	66	-100	1	-71	-28	-5
Below Normal (17.5%)								
No Action Alternative	4,050,002	780,040	145,797	60	4,911,682	4,196	307,440	10,199,216
Alternative 3	4,049,375	773,748	82,456	14	4,909,811	3,764	314,105	10,133,272
Difference	-627	-6,292	-63,341	-46	-1,871	-431	6,665	-65,944
Percent Difference	0	-1	-43	-77	0	-10	2	-1
Dry (22.5%)								
No Action Alternative	5,226,978	377,492	752,548	0	4,408,740	3,623	559,604	11,328,986
Alternative 3	3,355,934	388,784	658,614	0	4,450,665	2,536	521,440	9,377,972
Difference	-1,871,044	11,291	-93,934	0	41,925	-1,088	-38,164	-1,951,014
Percent Difference	-36	3	-12	0	1	-30	-7	-17
Critical (15%)								
No Action Alternative	11,740,400	395,039	2,255,935	0	3,441,525	42,525	526,526	18,401,950
Alternative 3	7,449,300	428,029	3,507,175	0	3,723,000	35,178	534,928	15,677,609
Difference	-4,291,101	32,990	1,251,240	0	281,475	-7,347	8,402	-2,724,340
Percent Difference	-37	8	55	0	8	-17	2	-15
1 Based on the 80-year simulation period								
2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.								
3 Relative difference of the Annual average								
4 Mortality values do not include base mortality								

Table B-1-10. Annual Mortality by All Factors for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super-imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	5,139,812	1,449,851	505,839	799,452	154	4,683,874	4,419	268,257	5,856	41,665	12,899,179
Alternative 3	3,882,019	1,491,155	639,732	860,812	146	4,708,991	3,342	255,443	5,247	41,977	11,888,865
Difference	-1,257,793	41,304	133,893	61,360	-8	25,116	-1,077	-12,814	-609	313	-1,010,314
Percent Difference ³	-24	3	26	8	-5	1	-24	-5	-10	1	-8
Water Year Types²											
Wet (32.5%)											
No Action Alternative	213,200	3,859,065	1,238,281	708,520	428	5,200,677	4,236	70,199	1,162	14,703	11,310,471
Alternative 3	37,613	3,945,868	1,651,803	346,009	441	5,099,364	4,272	71,120	1,605	14,761	11,172,855
Difference	-175,587	86,803	413,522	-362,510	13	-101,313	36	921	442	58	-137,615
Percent Difference	-82	2	33	-51	3	-2	1	1	38	0	-1
Above Normal (12.5%)											
No Action Alternative	11,397,132	67,263	79,569	287,640	34	5,007,318	3,300	158,529	1,438	30,567	17,032,790
Alternative 3	10,309,394	116,493	79,969	477,321	0	5,061,047	576	110,227	808	25,595	16,181,431
Difference	-1,087,738	49,230	401	189,681	-34	53,729	-2,724	-48,301	-630	-4,972	-851,359
Percent Difference	-10	73	1	66	-100	1	-83	-30	-44	-16	-5
Below Normal (17.5%)											
No Action Alternative	4,050,002	246,033	534,007	145,797	60	4,911,682	2,887	263,192	1,308	44,248	10,199,216
Alternative 3	4,049,375	242,891	530,857	82,456	14	4,909,811	2,116	265,663	1,649	48,442	10,133,272
Difference	-627	-3,142	-3,151	-63,341	-46	-1,871	-771	2,470	340	4,195	-65,944
Percent Difference	0	-1	-1	-43	-77	0	-27	1	26	9	-1
Dry (22.5%)											
No Action Alternative	5,226,978	377,492	0	752,548	0	4,408,740	1,403	500,298	2,220	59,306	11,328,986
Alternative 3	3,355,934	388,784	0	658,614	0	4,450,665	698	463,335	1,837	58,105	9,377,972
Difference	-1,871,044	11,291	0	-93,934	0	41,925	-705	-36,963	-382	-1,200	-1,951,014
Percent Difference	-36	3	0	-12	0	1	-50	-7	-17	-2	-17
Critical (15%)											
No Action Alternative	11,740,400	395,039	0	2,255,935	0	3,441,525	12,058	446,671	30,467	79,854	18,401,950
Alternative 3	7,449,300	428,029	0	3,507,175	0	3,723,000	9,030	452,064	26,148	82,864	15,677,609
Difference	-4,291,101	32,990	0	1,251,240	0	281,475	-3,028	5,392	-4,320	3,010	-2,724,340
Percent Difference	-37	8	0	55	0	8	-25	1	-14	4	-15

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-11. Annual Potential Production for Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	16,838,069
Alternative 5	16,908,477
Difference	70,408
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	16,537,313
Alternative 5	16,493,092
Difference	-44,221
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	15,696,855
Alternative 5	15,891,098
Difference	194,243
Percent Difference	1
Below Normal (17.5%)	
No Action Alternative	17,922,930
Alternative 5	17,951,192
Difference	28,262
Percent Difference	0
Dry (22.5%)	
No Action Alternative	17,754,135
Alternative 5	18,003,040
Difference	248,905
Percent Difference	1
Critical (15%)	
No Action Alternative	15,800,949
Alternative 5	15,797,949
Difference	-3,000
Percent Difference	0
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-1-12. Annual Mortality by Life Stage for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	7,894,954	4,684,028	272,676	47,521	320,197
Alternative 5	7,723,389	4,663,905	266,371	49,003	315,374
Difference	-171,565	-20,123	-6,305	1,482	-4,823
Percent Difference ³	-2	0	-2	3	-2
Water Year Types²					
Wet (32.5%)					
No Action Alternative	6,019,065	5,201,105	74,435	15,865	90,301
Alternative 5	6,169,444	5,177,967	78,031	16,578	94,608
Difference	150,379	-23,138	3,595	712	4,308
Percent Difference	2	0	5	4	5
Above Normal (12.5%)					
No Action Alternative	11,831,604	5,007,353	161,828	32,005	193,834
Alternative 5	11,229,256	4,990,191	153,381	34,302	187,683
Difference	-602,348	-17,162	-8,448	2,296	-6,151
Percent Difference	-5	0	-5	7	-3
Below Normal (17.5%)					
No Action Alternative	4,975,839	4,911,742	266,079	45,556	311,635
Alternative 5	4,934,725	4,906,604	268,136	45,725	313,861
Difference	-41,114	-5,138	2,056	169	2,226
Percent Difference	-1	0	1	0	1
Dry (22.5%)					
No Action Alternative	6,357,019	4,408,740	501,702	61,525	563,227
Alternative 5	5,727,952	4,357,900	490,190	66,478	556,668
Difference	-629,067	-50,840	-11,512	4,953	-6,559
Percent Difference	-10	-1	-2	8	-1
Critical (15%)					
No Action Alternative	14,391,374	3,441,525	458,729	110,322	569,051
Alternative 5	14,415,310	3,454,056	430,811	109,120	539,931
Difference	23,936	12,531	-27,918	-1,202	-29,120
Percent Difference	0	0	-6	-1	-5

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-1-13. Annual Mortality by Cause for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	5,949,693	6,949,486	12,899,179
Alternative 5	5,781,882	6,920,785	12,702,667
Difference	-167,811	-28,701	-196,511
Percent Difference ³	-3	0	-2
Water Year Types²			
Wet (32.5%)			
No Action Alternative	927,546	10,382,925	11,310,471
Alternative 5	1,088,909	10,353,111	11,442,020
Difference	161,363	-29,814	131,549
Percent Difference	17	0	1
Above Normal (12.5%)			
No Action Alternative	11,689,545	5,343,245	17,032,790
Alternative 5	11,083,720	5,323,409	16,407,129
Difference	-605,825	-19,836	-625,661
Percent Difference	-5	0	-4
Below Normal (17.5%)			
No Action Alternative	4,200,054	5,999,162	10,199,216
Alternative 5	4,169,106	5,986,084	10,155,190
Difference	-30,948	-13,078	-44,026
Percent Difference	-1	0	0
Dry (22.5%)			
No Action Alternative	5,983,150	5,345,836	11,328,986
Alternative 5	5,349,191	5,293,329	10,642,520
Difference	-633,958	-52,507	-686,466
Percent Difference	-11	-1	-6
Critical (15%)			
No Action Alternative	14,038,861	4,363,089	18,401,950
Alternative 5	14,062,400	4,346,896	18,409,296
Difference	23,539	-16,193	7,347
Percent Difference	0	0	0

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-14. Annual Mortality by Cause and Life Stage for Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	5,139,812	1,955,690	799,452	154	4,683,874	10,275	309,922	12,899,179
Alternative 5	4,786,653	1,951,663	985,073	154	4,663,751	10,003	305,371	12,702,667
Difference	-353,159	-4,026	185,621	0	-20,123	-272	-4,551	-196,511
Percent Difference ³	-7	0	23	0	0	-3	-1	-2
Water Year Types²								
Wet (32.5%)								
No Action Alternative	213,200	5,097,346	708,520	428	5,200,677	5,398	84,903	11,310,471
Alternative 5	348,257	5,086,105	735,082	436	5,177,531	5,134	89,475	11,442,020
Difference	135,058	-11,241	26,562	8	-23,146	-265	4,572	131,549
Percent Difference	63	0	4	2	0	-5	5	1
Above Normal (12.5%)								
No Action Alternative	11,397,132	146,831	287,640	34	5,007,318	4,738	189,095	17,032,790
Alternative 5	10,385,418	149,961	693,877	9	4,990,182	4,417	183,266	16,407,129
Difference	-1,011,714	3,130	406,236	-26	-17,136	-321	-5,830	-625,661
Percent Difference	-9	2	141	-75	0	-7	-3	-4
Below Normal (17.5%)								
No Action Alternative	4,050,002	780,040	145,797	60	4,911,682	4,196	307,440	10,199,216
Alternative 5	4,052,333	769,810	112,581	59	4,906,545	4,133	309,728	10,155,190
Difference	2,331	-10,229	-33,215	0	-5,137	-63	2,289	-44,026
Percent Difference	0	-1	-23	-1	0	-1	1	0
Dry (22.5%)								
No Action Alternative	5,226,978	377,492	752,548	0	4,408,740	3,623	559,604	11,328,986
Alternative 5	4,376,903	382,888	968,162	1	4,357,898	4,125	552,543	10,642,520
Difference	-850,076	5,395	215,614	1	-50,841	502	-7,061	-686,466
Percent Difference	-16	1	29	0	-1	14	-1	-6
Critical (15%)								
No Action Alternative	11,740,400	395,039	2,255,935	0	3,441,525	42,525	526,526	18,401,950
Alternative 5	11,208,869	393,784	2,812,657	0	3,454,056	40,874	499,057	18,409,296
Difference	-531,531	-1,255	556,722	0	12,531	-1,651	-27,469	7,347
Percent Difference	-5	0	25	0	0	-4	-5	0

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-1-15. Annual Mortality by All Factors for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
No Action Alternative	5,139,812	1,449,851	505,839	799,452	154	4,683,874	4,419	268,257	5,856	41,665	12,899,179
Alternative 5	4,786,653	1,450,386	501,277	985,073	154	4,663,751	4,489	261,882	5,514	43,488	12,702,667
Difference	-353,159	535	-4,561	185,621	0	-20,123	70	-6,375	-342	1,824	-196,511
Percent Difference ³	-7	0	-1	23	0	0	2	-2	-6	4	-2
Water Year Types²											
Wet (32.5%)											
No Action Alternative	213,200	3,859,065	1,238,281	708,520	428	5,200,677	4,236	70,199	1,162	14,703	11,310,471
Alternative 5	348,257	3,861,662	1,224,443	735,082	436	5,177,531	4,005	74,026	1,129	15,449	11,442,020
Difference	135,058	2,597	-13,838	26,562	8	-23,146	-231	3,827	-33	746	131,549
Percent Difference	63	0	-1	4	2	0	-5	5	-3	5	1
Above Normal (12.5%)											
No Action Alternative	11,397,132	67,263	79,569	287,640	34	5,007,318	3,300	158,529	1,438	30,567	17,032,790
Alternative 5	10,385,418	69,983	79,978	693,877	9	4,990,182	3,244	150,137	1,173	33,128	16,407,129
Difference	-1,011,714	2,721	409	406,236	-26	-17,136	-56	-8,391	-265	2,561	-625,661
Percent Difference	-9	4	1	141	-75	0	-2	-5	-18	8	-4
Below Normal (17.5%)											
No Action Alternative	4,050,002	246,033	534,007	145,797	60	4,911,682	2,887	263,192	1,308	44,248	10,199,216
Alternative 5	4,052,333	236,463	533,348	112,581	59	4,906,545	2,782	265,353	1,350	44,375	10,155,190
Difference	2,331	-9,570	-659	-33,215	0	-5,137	-105	2,161	42	128	-44,026
Percent Difference	0	-4	0	-23	-1	0	-4	1	3	0	0
Dry (22.5%)											
No Action Alternative	5,226,978	377,492	0	752,548	0	4,408,740	1,403	500,298	2,220	59,306	11,328,986
Alternative 5	4,376,903	382,888	0	968,162	1	4,357,898	1,827	488,363	2,298	64,180	10,642,520
Difference	-850,076	5,395	0	215,614	1	-50,841	424	-11,936	79	4,874	-686,466
Percent Difference	-16	1	0	29	0	-1	30	-2	4	8	-6
Critical (15%)											
No Action Alternative	11,740,400	395,039	0	2,255,935	0	3,441,525	12,058	446,671	30,467	79,854	18,401,950
Alternative 5	11,208,869	393,784	0	2,812,657	0	3,454,056	12,558	418,253	28,316	80,804	18,409,296
Difference	-531,531	-1,255	0	556,722	0	12,531	500	-28,418	-2,151	949	7,347
Percent Difference	-5	0	0	25	0	0	4	-6	-7	1	0

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-16. Annual Potential Production for Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	17,037,309
No Action Alternative	16,838,069
Difference	-199,240
Percent Difference ³	-1
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	16,525,365
No Action Alternative	16,537,313
Difference	11,948
Percent Difference	0
Above Normal (12.5%)	
Second Basis of Comparison	15,746,827
No Action Alternative	15,696,855
Difference	-49,972
Percent Difference	0
Below Normal (17.5%)	
Second Basis of Comparison	17,847,310
No Action Alternative	17,922,930
Difference	75,620
Percent Difference	0
Dry (22.5%)	
Second Basis of Comparison	17,934,726
No Action Alternative	17,754,135
Difference	-180,590
Percent Difference	-1
Critical (15%)	
Second Basis of Comparison	16,930,799
No Action Alternative	15,800,949
Difference	-1,129,850
Percent Difference	-7
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-1-17. Annual Mortality by Life Stage for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	7,110,950	4,709,109	269,215	49,405	318,621
No Action Alternative	7,894,954	4,684,028	272,676	47,521	320,197
Difference	784,003	-25,081	3,461	-1,885	1,576
Percent Difference ³	11	-1	1	-4	0
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	6,023,551	5,129,591	71,744	16,838	88,581
No Action Alternative	6,019,065	5,201,105	74,435	15,865	90,301
Difference	-4,486	71,514	2,692	-973	1,719
Percent Difference	0	1	4	-6	2
Above Normal (12.5%)					
Second Basis of Comparison	11,326,553	5,120,441	96,157	31,173	127,329
No Action Alternative	11,831,604	5,007,353	161,828	32,005	193,834
Difference	505,051	-113,088	65,672	833	66,505
Percent Difference	4	-2	68	3	52
Below Normal (17.5%)					
Second Basis of Comparison	4,943,736	4,895,243	284,538	50,880	335,418
No Action Alternative	4,975,839	4,911,742	266,079	45,556	311,635
Difference	32,103	16,499	-18,459	-5,324	-23,783
Percent Difference	1	0	-6	-10	-7
Dry (22.5%)					
Second Basis of Comparison	5,846,335	4,371,799	440,615	59,727	500,342
No Action Alternative	6,357,019	4,408,740	501,702	61,525	563,227
Difference	510,683	36,940	61,087	1,798	62,885
Percent Difference	9	1	14	3	13
Critical (15%)					
Second Basis of Comparison	10,379,320	3,744,097	566,311	117,959	684,270
No Action Alternative	14,391,374	3,441,525	458,729	110,322	569,051
Difference	4,012,054	-302,572	-107,582	-7,638	-115,220
Percent Difference	39	-8	-19	-6	-17

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-1-18. Annual Mortality by Cause for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	5,010,581	7,128,100	12,138,680
No Action Alternative	5,949,693	6,949,486	12,899,179
Difference	939,112	-178,614	760,499
Percent Difference ³	19	-3	6
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	485,103	10,756,621	11,241,723
No Action Alternative	927,546	10,382,925	11,310,471
Difference	442,443	-373,695	68,747
Percent Difference	91	-3	1
Above Normal (12.5%)			
Second Basis of Comparison	11,136,551	5,437,771	16,574,323
No Action Alternative	11,689,545	5,343,245	17,032,790
Difference	552,994	-94,526	458,468
Percent Difference	5	-2	3
Below Normal (17.5%)			
Second Basis of Comparison	4,155,751	6,018,646	10,174,397
No Action Alternative	4,200,054	5,999,162	10,199,216
Difference	44,304	-19,484	24,819
Percent Difference	1	0	0
Dry (22.5%)			
Second Basis of Comparison	5,469,925	5,248,551	10,718,477
No Action Alternative	5,983,150	5,345,836	11,328,986
Difference	513,224	97,285	610,509
Percent Difference	9	2	6
Critical (15%)			
Second Basis of Comparison	10,019,091	4,788,596	14,807,687
No Action Alternative	14,038,861	4,363,089	18,401,950
Difference	4,019,770	-425,507	3,594,263
Percent Difference	40	-9	24

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-19. Annual Mortality by Cause and Life Stage for Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	4,292,224	2,108,590	710,136	151	4,708,958	8,069	310,552	12,138,680
No Action Alternative	5,139,812	1,955,690	799,452	154	4,683,874	10,275	309,922	12,899,179
Difference	847,588	-152,900	89,315	3	-25,084	2,206	-630	760,499
Percent Difference ³	20	-7	13	2	-1	27	0	6
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	76,487	5,544,710	402,355	446	5,129,145	5,816	82,766	11,241,723
No Action Alternative	213,200	5,097,346	708,520	428	5,200,677	5,398	84,903	11,310,471
Difference	136,713	-447,364	306,165	-18	71,532	-417	2,137	68,747
Percent Difference	179	-8	76	-4	1	-7	3	1
Above Normal (12.5%)								
Second Basis of Comparison	10,875,176	194,605	256,772	9	5,120,432	4,595	122,734	16,574,323
No Action Alternative	11,397,132	146,831	287,640	34	5,007,318	4,738	189,095	17,032,790
Difference	521,956	-47,774	30,868	26	-113,113	144	66,361	458,468
Percent Difference	5	-25	12	287	-2	3	54	3
Below Normal (17.5%)								
Second Basis of Comparison	4,055,314	789,925	98,496	25	4,895,218	1,915	333,503	10,174,397
No Action Alternative	4,050,002	780,040	145,797	60	4,911,682	4,196	307,440	10,199,216
Difference	-5,312	-9,886	47,300	35	16,465	2,280	-26,064	24,819
Percent Difference	0	-1	48	138	0	119	-8	0
Dry (22.5%)								
Second Basis of Comparison	4,603,020	378,293	865,023	0	4,371,799	1,883	498,459	10,718,477
No Action Alternative	5,226,978	377,492	752,548	0	4,408,740	3,623	559,604	11,328,986
Difference	623,959	-801	-112,475	0	36,940	1,740	61,145	610,509
Percent Difference	14	0	-13	0	1	92	12	6
Critical (15%)								
Second Basis of Comparison	7,750,732	392,537	2,236,052	0	3,744,097	32,307	651,963	14,807,687
No Action Alternative	11,740,400	395,039	2,255,935	0	3,441,525	42,525	526,526	18,401,950
Difference	3,989,668	2,502	19,884	0	-302,572	10,218	-125,438	3,594,263
Percent Difference	51	1	1	0	-8	32	-19	24

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-1-20. Annual Mortality by All Factors for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	4,292,224	1,473,372	635,217	710,136	151	4,708,958	3,312	265,903	4,757	44,648	12,138,680
No Action Alternative	5,139,812	1,449,851	505,839	799,452	154	4,683,874	4,419	268,257	5,856	41,665	12,899,179
Difference	847,588	-23,521	-129,379	89,315	3	-25,084	1,106	2,354	1,099	-2,984	760,499
Percent Difference ³	20	-2	-20	13	2	-1	33	1	23	-7	6
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	76,487	3,907,496	1,637,214	402,355	446	5,129,145	4,203	67,541	1,613	15,225	11,241,723
No Action Alternative	213,200	3,859,065	1,238,281	708,520	428	5,200,677	4,236	70,199	1,162	14,703	11,310,471
Difference	136,713	-48,431	-398,933	306,165	-18	71,532	33	2,659	-451	-522	68,747
Percent Difference	179	-1	-24	76	-4	1	1	4	-28	-3	1
Above Normal (12.5%)											
Second Basis of Comparison	10,875,176	114,650	79,955	256,772	9	5,120,432	3,015	93,141	1,579	29,593	16,574,323
No Action Alternative	11,397,132	67,263	79,569	287,640	34	5,007,318	3,300	158,529	1,438	30,567	17,032,790
Difference	521,956	-47,387	-386	30,868	26	-113,113	285	65,387	-141	974	458,468
Percent Difference	5	-41	0	12	287	-2	9	70	-9	3	3
Below Normal (17.5%)											
Second Basis of Comparison	4,055,314	257,762	532,163	98,496	25	4,895,218	1,115	283,424	801	50,079	10,174,397
No Action Alternative	4,050,002	246,033	534,007	145,797	60	4,911,682	2,887	263,192	1,308	44,248	10,199,216
Difference	-5,312	-11,729	1,844	47,300	35	16,465	1,773	-20,232	508	-5,832	24,819
Percent Difference	0	-5	0	48	138	0	159	-7	63	-12	0
Dry (22.5%)											
Second Basis of Comparison	4,603,020	378,293	0	865,023	0	4,371,799	423	440,192	1,460	58,267	10,718,477
No Action Alternative	5,226,978	377,492	0	752,548	0	4,408,740	1,403	500,298	2,220	59,306	11,328,986
Difference	623,959	-801	0	-112,475	0	36,940	980	60,107	760	1,038	610,509
Percent Difference	14	0	0	-13	0	1	232	14	52	2	6
Critical (15%)											
Second Basis of Comparison	7,750,732	392,537	0	2,236,052	0	3,744,097	8,529	557,782	23,779	94,181	14,807,687
No Action Alternative	11,740,400	395,039	0	2,255,935	0	3,441,525	12,058	446,671	30,467	79,854	18,401,950
Difference	3,989,668	2,502	0	19,884	0	-302,572	3,529	-111,111	6,689	-14,327	3,594,263
Percent Difference	51	1	0	1	0	-8	41	-20	28	-15	24

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-1-21. Annual Potential Production for Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	17,037,309
Alternative 3	17,129,024
Difference	91,715
Percent Difference ³	1
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	16,525,365
Alternative 3	16,544,696
Difference	19,331
Percent Difference	0
Above Normal (12.5%)	
Second Basis of Comparison	15,746,827
Alternative 3	15,897,563
Difference	150,736
Percent Difference	1
Below Normal (17.5%)	
Second Basis of Comparison	17,847,310
Alternative 3	17,877,415
Difference	30,105
Percent Difference	0
Dry (22.5%)	
Second Basis of Comparison	17,934,726
Alternative 3	18,382,793
Difference	448,067
Percent Difference	2
Critical (15%)	
Second Basis of Comparison	16,930,799
Alternative 3	16,667,512
Difference	-263,288
Percent Difference	-2
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-1-22. Annual Mortality by Life Stage for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	7,110,950	4,709,109	269,215	49,405	318,621
Alternative 3	6,873,719	4,709,136	258,786	47,224	306,009
Difference	-237,232	27	-10,430	-2,182	-12,611
Percent Difference ³	-3	0	-4	-4	-4
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	6,023,551	5,129,591	71,744	16,838	88,581
Alternative 3	5,981,293	5,099,805	75,392	16,365	91,757
Difference	-42,258	-29,786	3,648	-473	3,176
Percent Difference	-1	-1	5	-3	4
Above Normal (12.5%)					
Second Basis of Comparison	11,326,553	5,120,441	96,157	31,173	127,329
Alternative 3	10,983,177	5,061,047	110,803	26,403	137,207
Difference	-343,376	-59,394	14,647	-4,769	9,878
Percent Difference	-3	-1	15	-15	8
Below Normal (17.5%)					
Second Basis of Comparison	4,943,736	4,895,243	284,538	50,880	335,418
Alternative 3	4,905,579	4,909,824	267,778	50,091	317,869
Difference	-38,157	14,582	-16,760	-789	-17,549
Percent Difference	-1	0	-6	-2	-5
Dry (22.5%)					
Second Basis of Comparison	5,846,335	4,371,799	440,615	59,727	500,342
Alternative 3	4,403,331	4,450,665	464,033	59,943	523,976
Difference	-1,443,004	78,865	23,419	215	23,634
Percent Difference	-25	2	5	0	5
Critical (15%)					
Second Basis of Comparison	10,379,320	3,744,097	566,311	117,959	684,270
Alternative 3	11,384,504	3,723,000	461,093	109,012	570,105
Difference	1,005,183	-21,096	-105,218	-8,947	-114,165
Percent Difference	10	-1	-19	-8	-17

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-1-23. Annual Mortality by Cause for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	5,010,581	7,128,100	12,138,680
Alternative 3	4,751,566	7,137,299	11,888,865
Difference	-259,015	9,199	-249,816
Percent Difference ³	-5	0	-2
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	485,103	10,756,621	11,241,723
Alternative 3	389,939	10,782,916	11,172,855
Difference	-95,164	26,295	-68,868
Percent Difference	-20	0	-1
Above Normal (12.5%)			
Second Basis of Comparison	11,136,551	5,437,771	16,574,323
Alternative 3	10,788,099	5,393,332	16,181,431
Difference	-348,452	-44,440	-392,892
Percent Difference	-3	-1	-2
Below Normal (17.5%)			
Second Basis of Comparison	4,155,751	6,018,646	10,174,397
Alternative 3	4,135,609	5,997,663	10,133,272
Difference	-20,141	-20,983	-41,125
Percent Difference	0	0	0
Dry (22.5%)			
Second Basis of Comparison	5,469,925	5,248,551	10,718,477
Alternative 3	4,017,083	5,360,888	9,377,972
Difference	-1,452,842	112,337	-1,340,505
Percent Difference	-27	2	-13
Critical (15%)			
Second Basis of Comparison	10,019,091	4,788,596	14,807,687
Alternative 3	10,991,653	4,685,957	15,677,609
Difference	972,562	-102,640	869,922
Percent Difference	10	-2	6

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-24. Annual Mortality by Cause and Life Stage for Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Temperature	Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat				
Long-term									
Full Simulation Period¹									
Second Basis of Comparison	4,292,224	2,108,590	710,136	151	4,708,958	8,069	310,552	12,138,680	
Alternative 3	3,882,019	2,130,887	860,812	146	4,708,991	8,589	297,421	11,888,865	
Difference	-410,205	22,298	150,676	-5	32	520	-13,131	-249,816	
Percent Difference ³	-10	1	21	-3	0	6	-4	-2	
Water Year Types²									
Wet (32.5%)									
Second Basis of Comparison	76,487	5,544,710	402,355	446	5,129,145	5,816	82,766	11,241,723	
Alternative 3	37,613	5,597,671	346,009	441	5,099,364	5,877	85,881	11,172,855	
Difference	-38,874	52,961	-56,345	-5	-29,781	61	3,115	-68,868	
Percent Difference	-51	1	-14	-1	-1	1	4	-1	
Above Normal (12.5%)									
Second Basis of Comparison	10,875,176	194,605	256,772	9	5,120,432	4,595	122,734	16,574,323	
Alternative 3	10,309,394	196,462	477,321	0	5,061,047	1,384	135,823	16,181,431	
Difference	-565,781	1,857	220,549	-9	-59,385	-3,210	13,088	-392,892	
Percent Difference	-5	1	86	-100	-1	-70	11	-2	
Below Normal (17.5%)									
Second Basis of Comparison	4,055,314	789,925	98,496	25	4,895,218	1,915	333,503	10,174,397	
Alternative 3	4,049,375	773,748	82,456	14	4,909,811	3,764	314,105	10,133,272	
Difference	-5,939	-16,178	-16,041	-12	14,593	1,849	-19,399	-41,125	
Percent Difference	0	-2	-16	-46	0	97	-6	0	
Dry (22.5%)									
Second Basis of Comparison	4,603,020	378,293	865,023	0	4,371,799	1,883	498,459	10,718,477	
Alternative 3	3,355,934	388,784	658,614	0	4,450,665	2,536	521,440	9,377,972	
Difference	-1,247,086	10,491	-206,409	0	78,865	653	22,981	-1,340,505	
Percent Difference	-27	3	-24	0	2	35	5	-13	
Critical (15%)									
Second Basis of Comparison	7,750,732	392,537	2,236,052	0	3,744,097	32,307	651,963	14,807,687	
Alternative 3	7,449,300	428,029	3,507,175	0	3,723,000	35,178	534,928	15,677,609	
Difference	-301,433	35,492	1,271,124	0	-21,096	2,870	-117,035	869,922	
Percent Difference	-4	9	57	0	-1	9	-18	6	

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-1-25. Annual Mortality by All Factors for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	4,292,224	1,473,372	635,217	710,136	151	4,708,958	3,312	265,903	4,757	44,648	12,138,680
Alternative 3	3,882,019	1,491,155	639,732	860,812	146	4,708,991	3,342	255,443	5,247	41,977	11,888,865
Difference	-410,205	17,783	4,515	150,676	-5	32	30	-10,460	490	-2,671	-249,816
Percent Difference ³	-10	1	1	21	-3	0	1	-4	10	-6	-2
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	76,487	3,907,496	1,637,214	402,355	446	5,129,145	4,203	67,541	1,613	15,225	11,241,723
Alternative 3	37,613	3,945,868	1,651,803	346,009	441	5,099,364	4,272	71,120	1,605	14,761	11,172,855
Difference	-38,874	38,372	14,589	-56,345	-5	-29,781	69	3,579	-8	-465	-68,868
Percent Difference	-51	1	1	-14	-1	-1	2	5	-1	-3	-1
Above Normal (12.5%)											
Second Basis of Comparison	10,875,176	114,650	79,955	256,772	9	5,120,432	3,015	93,141	1,579	29,593	16,574,323
Alternative 3	10,309,394	116,493	79,969	477,321	0	5,061,047	576	110,227	808	25,595	16,181,431
Difference	-565,781	1,843	14	220,549	-9	-59,385	-2,439	17,086	-771	-3,998	-392,892
Percent Difference	-5	2	0	86	-100	-1	-81	18	-49	-14	-2
Below Normal (17.5%)											
Second Basis of Comparison	4,055,314	257,762	532,163	98,496	25	4,895,218	1,115	283,424	801	50,079	10,174,397
Alternative 3	4,049,375	242,891	530,857	82,456	14	4,909,811	2,116	265,663	1,649	48,442	10,133,272
Difference	-5,939	-14,871	-1,307	-16,041	-12	14,593	1,001	-17,761	848	-1,637	-41,125
Percent Difference	0	-6	0	-16	-46	0	90	-6	106	-3	0
Dry (22.5%)											
Second Basis of Comparison	4,603,020	378,293	0	865,023	0	4,371,799	423	440,192	1,460	58,267	10,718,477
Alternative 3	3,355,934	388,784	0	658,614	0	4,450,665	698	463,335	1,837	58,105	9,377,972
Difference	-1,247,086	10,491	0	-206,409	0	78,865	275	23,144	378	-162	-1,340,505
Percent Difference	-27	3	0	-24	0	2	65	5	26	0	-13
Critical (15%)											
Second Basis of Comparison	7,750,732	392,537	0	2,236,052	0	3,744,097	8,529	557,782	23,779	94,181	14,807,687
Alternative 3	7,449,300	428,029	0	3,507,175	0	3,723,000	9,030	452,064	26,148	82,864	15,677,609
Difference	-301,433	35,492	0	1,271,124	0	-21,096	501	-105,719	2,369	-11,317	869,922
Percent Difference	-4	9	0	57	0	-1	6	-19	10	-12	6

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-26. Annual Potential Production for Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	17,037,309
Alternative 5	16,908,477
Difference	-128,832
Percent Difference ³	-1
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	16,525,365
Alternative 5	16,493,092
Difference	-32,272
Percent Difference	0
Above Normal (12.5%)	
Second Basis of Comparison	15,746,827
Alternative 5	15,891,098
Difference	144,271
Percent Difference	1
Below Normal (17.5%)	
Second Basis of Comparison	17,847,310
Alternative 5	17,951,192
Difference	103,882
Percent Difference	1
Dry (22.5%)	
Second Basis of Comparison	17,934,726
Alternative 5	18,003,040
Difference	68,315
Percent Difference	0
Critical (15%)	
Second Basis of Comparison	16,930,799
Alternative 5	15,797,949
Difference	-1,132,850
Percent Difference	-7
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-1-27. Annual Mortality by Life Stage for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	7,110,950	4,709,109	269,215	49,405	318,621
Alternative 5	7,723,389	4,663,905	266,371	49,003	315,374
Difference	612,438	-45,204	-2,845	-402	-3,247
Percent Difference ³	9	-1	-1	-1	-1
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	6,023,551	5,129,591	71,744	16,838	88,581
Alternative 5	6,169,444	5,177,967	78,031	16,578	94,608
Difference	145,893	48,376	6,287	-260	6,027
Percent Difference	2	1	9	-2	7
Above Normal (12.5%)					
Second Basis of Comparison	11,326,553	5,120,441	96,157	31,173	127,329
Alternative 5	11,229,256	4,990,191	153,381	34,302	187,683
Difference	-97,297	-130,250	57,224	3,129	60,354
Percent Difference	-1	-3	60	10	47
Below Normal (17.5%)					
Second Basis of Comparison	4,943,736	4,895,243	284,538	50,880	335,418
Alternative 5	4,934,725	4,906,604	268,136	45,725	313,861
Difference	-9,011	11,362	-16,403	-5,155	-21,557
Percent Difference	0	0	-6	-10	-6
Dry (22.5%)					
Second Basis of Comparison	5,846,335	4,371,799	440,615	59,727	500,342
Alternative 5	5,727,952	4,357,900	490,190	66,478	556,668
Difference	-118,383	-13,900	49,576	6,751	56,326
Percent Difference	-2	0	11	11	11
Critical (15%)					
Second Basis of Comparison	10,379,320	3,744,097	566,311	117,959	684,270
Alternative 5	14,415,310	3,454,056	430,811	109,120	539,931
Difference	4,035,990	-290,041	-135,500	-8,839	-144,340
Percent Difference	39	-8	-24	-7	-21

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-1-28. Annual Mortality by Cause for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	5,010,581	7,128,100	12,138,680
Alternative 5	5,781,882	6,920,785	12,702,667
Difference	771,302	-207,314	563,987
Percent Difference ³	15	-3	5
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	485,103	10,756,621	11,241,723
Alternative 5	1,088,909	10,353,111	11,442,020
Difference	603,806	-403,510	200,296
Percent Difference	124	-4	2
Above Normal (12.5%)			
Second Basis of Comparison	11,136,551	5,437,771	16,574,323
Alternative 5	11,083,720	5,323,409	16,407,129
Difference	-52,831	-114,362	-167,193
Percent Difference	0	-2	-1
Below Normal (17.5%)			
Second Basis of Comparison	4,155,751	6,018,646	10,174,397
Alternative 5	4,169,106	5,986,084	10,155,190
Difference	13,356	-32,563	-19,207
Percent Difference	0	-1	0
Dry (22.5%)			
Second Basis of Comparison	5,469,925	5,248,551	10,718,477
Alternative 5	5,349,191	5,293,329	10,642,520
Difference	-120,734	44,777	-75,957
Percent Difference	-2	1	-1
Critical (15%)			
Second Basis of Comparison	10,019,091	4,788,596	14,807,687
Alternative 5	14,062,400	4,346,896	18,409,296
Difference	4,043,309	-441,700	3,601,609
Percent Difference	40	-9	24

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-29. Annual Mortality by Cause and Life Stage for Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Temperature	Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat				
Long-term									
Full Simulation Period¹									
Second Basis of Comparison	4,292,224	2,108,590	710,136	151	4,708,958	8,069	310,552	12,138,680	
Alternative 5	4,786,653	1,951,663	985,073	154	4,663,751	10,003	305,371	12,702,667	
Difference	494,428	-156,926	274,936	3	-45,207	1,934	-5,181	563,987	
Percent Difference ³	12	-7	39	2	-1	24	-2	5	
Water Year Types²									
Wet (32.5%)									
Second Basis of Comparison	76,487	5,544,710	402,355	446	5,129,145	5,816	82,766	11,241,723	
Alternative 5	348,257	5,086,105	735,082	436	5,177,531	5,134	89,475	11,442,020	
Difference	271,771	-458,605	332,727	-10	48,386	-682	6,709	200,296	
Percent Difference	355	-8	83	-2	1	-12	8	2	
Above Normal (12.5%)									
Second Basis of Comparison	10,875,176	194,605	256,772	9	5,120,432	4,595	122,734	16,574,323	
Alternative 5	10,385,418	149,961	693,877	9	4,990,182	4,417	183,266	16,407,129	
Difference	-489,758	-44,644	437,104	0	-130,249	-178	60,531	-167,193	
Percent Difference	-5	-23	170	-4	-3	-4	49	-1	
Below Normal (17.5%)									
Second Basis of Comparison	4,055,314	789,925	98,496	25	4,895,218	1,915	333,503	10,174,397	
Alternative 5	4,052,333	769,810	112,581	59	4,906,545	4,133	309,728	10,155,190	
Difference	-2,981	-20,115	14,085	34	11,327	2,218	-23,775	-19,207	
Percent Difference	0	-3	14	137	0	116	-7	0	
Dry (22.5%)									
Second Basis of Comparison	4,603,020	378,293	865,023	0	4,371,799	1,883	498,459	10,718,477	
Alternative 5	4,376,903	382,888	968,162	1	4,357,898	4,125	552,543	10,642,520	
Difference	-226,117	4,595	103,139	1	-13,901	2,243	54,084	-75,957	
Percent Difference	-5	1	12	0	0	119	11	-1	
Critical (15%)									
Second Basis of Comparison	7,750,732	392,537	2,236,052	0	3,744,097	32,307	651,963	14,807,687	
Alternative 5	11,208,869	393,784	2,812,657	0	3,454,056	40,874	499,057	18,409,296	
Difference	3,458,137	1,247	576,606	0	-290,041	8,567	-152,907	3,601,609	
Percent Difference	45	0	26	0	-8	27	-23	24	

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-1-30. Annual Mortality by All Factors for Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	4,292,224	1,473,372	635,217	710,136	151	4,708,958	3,312	265,903	4,757	44,648	12,138,680
Alternative 5	4,786,653	1,450,386	501,277	985,073	154	4,663,751	4,489	261,882	5,514	43,488	12,702,667
Difference	494,428	-22,986	-133,940	274,936	3	-45,207	1,176	-4,021	758	-1,160	563,987
Percent Difference ³	12	-2	-21	39	2	-1	36	-2	16	-3	5
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	76,487	3,907,496	1,637,214	402,355	446	5,129,145	4,203	67,541	1,613	15,225	11,241,723
Alternative 5	348,257	3,861,662	1,224,443	735,082	436	5,177,531	4,005	74,026	1,129	15,449	11,442,020
Difference	271,771	-45,835	-412,770	332,727	-10	48,386	-198	6,485	-484	224	200,296
Percent Difference	355	-1	-25	83	-2	1	-5	10	-30	1	2
Above Normal (12.5%)											
Second Basis of Comparison	10,875,176	114,650	79,955	256,772	9	5,120,432	3,015	93,141	1,579	29,593	16,574,323
Alternative 5	10,385,418	69,983	79,978	693,877	9	4,990,182	3,244	150,137	1,173	33,128	16,407,129
Difference	-489,758	-44,667	23	437,104	0	-130,249	228	56,996	-406	3,535	-167,193
Percent Difference	-5	-39	0	170	-4	-3	8	61	-26	12	-1
Below Normal (17.5%)											
Second Basis of Comparison	4,055,314	257,762	532,163	98,496	25	4,895,218	1,115	283,424	801	50,079	10,174,397
Alternative 5	4,052,333	236,463	533,348	112,581	59	4,906,545	2,782	265,353	1,350	44,375	10,155,190
Difference	-2,981	-21,299	1,184	14,085	34	11,327	1,668	-18,071	550	-5,704	-19,207
Percent Difference	0	-8	0	14	137	0	150	-6	69	-11	0
Dry (22.5%)											
Second Basis of Comparison	4,603,020	378,293	0	865,023	0	4,371,799	423	440,192	1,460	58,267	10,718,477
Alternative 5	4,376,903	382,888	0	968,162	1	4,357,898	1,827	488,363	2,298	64,180	10,642,520
Difference	-226,117	4,595	0	103,139	1	-13,901	1,404	48,171	838	5,912	-75,957
Percent Difference	-5	1	0	12	0	0	332	11	57	10	-1
Critical (15%)											
Second Basis of Comparison	7,750,732	392,537	0	2,236,052	0	3,744,097	8,529	557,782	23,779	94,181	14,807,687
Alternative 5	11,208,869	393,784	0	2,812,657	0	3,454,056	12,558	418,253	28,316	80,804	18,409,296
Difference	3,458,137	1,247	0	576,606	0	-290,041	4,029	-139,529	4,538	-13,377	3,601,609
Percent Difference	45	0	0	26	0	-8	47	-25	19	-14	24

¹ Based on the 80-year simulation period


² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality


1 **B.2. Late Fall-Run Chinook Salmon**
2

Figure B-2-1. Annual Potential Production for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-2. Annual Mortality for Late Fall-Run Chinook Salmon - Eggs




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-3. Annual Mortality for Late Fall-Run Chinook Salmon - Fry




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-4. Annual Mortality for Late Fall-Run Chinook Salmon - Pre-Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-5. Annual Mortality for Late Fall-Run Chinook Salmon - Immature Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-6. Annual Mortality for Late Fall-Run Chinook Salmon - Pre- & Immature Smolts




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-7. Annual Mortality for Late Fall-Run Chinook Salmon - All Lifestages




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-8. Incubation - Habitat based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-9. Super-imposition - Habitat based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-10. Fry - Habitat based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-11. Pre-smolt - Habitat based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-12. Immature Smolt - Habitat based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-13. Total Habitat based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-14. Pre-Spawn Mortality - Temperature based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-15. Eggs - Temperature based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-16. Fry - Temperature based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-17. Pre-smolt - Temperature based Annual Mortality for Late Fall-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-18. Immature Smolt - Temperature based Annual Mortality for Late Fall-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-2-19. Total Temperature based Annual Mortality for Late Fall-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Table B-2-1. Annual Potential Production for Late Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	2,813,219
Alternative 1	2,800,061
Difference	-13,158
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	2,692,145
Alternative 1	2,691,035
Difference	-1,111
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	2,860,264
Alternative 1	2,802,912
Difference	-57,352
Percent Difference	-2
Below Normal (17.5%)	
No Action Alternative	2,982,412
Alternative 1	2,930,472
Difference	-51,940
Percent Difference	-2
Dry (22.5%)	
No Action Alternative	3,023,892
Alternative 1	2,976,338
Difference	-47,554
Percent Difference	-2
Critical (15%)	
No Action Alternative	2,522,939
Alternative 1	2,617,343
Difference	94,404
Percent Difference	4
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-2-2. Annual Mortality by Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	492,142	1,757,035	82,787	37,844	120,631
Alternative 1	513,890	1,802,954	68,169	30,510	98,679
Difference	21,748	45,920	-14,618	-7,334	-21,952
Percent Difference ³	4	3	-18	-19	-18
Water Year Types²					
Wet (32.5%)					
No Action Alternative	1,305,939	1,487,095	6,012	78	6,089
Alternative 1	1,331,500	1,479,904	4,935	609	5,544
Difference	25,561	-7,191	-1,076	531	-545
Percent Difference	2	0	-18	684	-9
Above Normal (12.5%)					
No Action Alternative	371,926	1,810,494	1,361	103	1,464
Alternative 1	482,073	1,869,446	2,387	187	2,573
Difference	110,146	58,952	1,025	84	1,109
Percent Difference	30	3	75	82	76
Below Normal (17.5%)					
No Action Alternative	38,722	1,885,067	14,022	4,588	18,610
Alternative 1	41,496	1,985,382	9,337	3,123	12,460
Difference	2,774	100,315	-4,685	-1,465	-6,150
Percent Difference	7	5	-33	-32	-33
Dry (22.5%)					
No Action Alternative	34,945	1,894,612	38,990	16,946	55,936
Alternative 1	34,962	1,979,833	29,461	15,809	45,270
Difference	17	85,221	-9,529	-1,137	-10,666
Percent Difference	0	4	-24	-7	-19
Critical (15%)					
No Action Alternative	43,879	1,941,615	462,907	221,268	684,174
Alternative 1	38,435	1,969,335	386,693	174,569	561,262
Difference	-5,445	27,720	-76,214	-46,699	-122,912
Percent Difference	-12	1	-16	-21	-18

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-2-3. Annual Mortality by Cause for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	117,312	2,252,495	2,369,807
Alternative 1	100,569	2,314,954	2,415,523
Difference	-16,743	62,459	45,716
Percent Difference ³	-14	3	2
Water Year Types²			
Wet (32.5%)			
No Action Alternative	11,538	2,787,586	2,799,124
Alternative 1	13,087	2,803,861	2,816,949
Difference	1,549	16,276	17,825
Percent Difference	13	1	1
Above Normal (12.5%)			
No Action Alternative	9,419	2,174,466	2,183,885
Alternative 1	9,812	2,344,280	2,354,092
Difference	393	169,814	170,208
Percent Difference	4	8	8
Below Normal (17.5%)			
No Action Alternative	16,631	1,925,768	1,942,399
Alternative 1	15,158	2,024,180	2,039,338
Difference	-1,474	98,412	96,938
Percent Difference	-9	5	5
Dry (22.5%)			
No Action Alternative	44,530	1,940,964	1,985,493
Alternative 1	40,463	2,019,602	2,060,065
Difference	-4,067	78,638	74,572
Percent Difference	-9	4	4
Critical (15%)			
No Action Alternative	663,032	2,006,637	2,669,669
Alternative 1	555,549	2,013,483	2,569,032
Difference	-107,483	6,846	-100,637
Percent Difference	-16	0	-4

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-4. Annual Mortality by Cause and Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)					Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature	Juvenile Habitat	
Long-term								
Full Simulation Period¹								
No Action Alternative	0	482,477	9,665	3,749	1,753,285	103,897	16,733	2,369,807
Alternative 1	0	504,586	9,304	3,662	1,799,292	87,603	11,076	2,415,523
Difference	0	22,110	-361	-87	46,006	-16,294	-5,657	45,716
Percent Difference ³	0	5	-4	-2	3	-16	-34	2
Water Year Types²								
Wet (32.5%)								
No Action Alternative	0	1,294,487	11,452	61	1,487,035	26	6,063	2,799,124
Alternative 1	0	1,319,517	11,983	61	1,479,843	1,043	4,501	2,816,949
Difference	0	25,030	531	0	-7,192	1,018	-1,563	17,825
Percent Difference	0	2	5	1	0	3,925	-26	1
Above Normal (12.5%)								
No Action Alternative	0	362,747	9,179	167	1,810,328	73	1,392	2,183,885
Alternative 1	0	472,813	9,259	147	1,869,299	405	2,168	2,354,092
Difference	0	110,066	80	-19	58,971	333	776	170,208
Percent Difference	0	30	1	-12	3	459	56	8
Below Normal (17.5%)								
No Action Alternative	0	28,022	10,701	143	1,884,924	5,787	12,822	1,942,399
Alternative 1	0	30,282	11,214	62	1,985,320	3,882	8,578	2,039,338
Difference	0	2,261	513	-81	100,396	-1,906	-4,244	96,938
Percent Difference	0	8	5	-57	5	-33	-33	5
Dry (22.5%)								
No Action Alternative	0	28,946	5,999	570	1,894,042	37,961	17,975	1,985,493
Alternative 1	0	30,519	4,444	1,218	1,978,615	34,802	10,468	2,060,065
Difference	0	1,573	-1,556	648	84,573	-3,159	-7,508	74,572
Percent Difference	0	5	-26	114	4	-8	-42	4
Critical (15%)								
No Action Alternative	0	33,389	10,490	23,702	1,917,913	628,839	55,335	2,669,669
Alternative 1	0	29,837	8,597	22,262	1,947,073	524,689	36,573	2,569,032
Difference	0	-3,552	-1,893	-1,440	29,160	-104,150	-18,762	-100,637
Percent Difference	0	-11	-18	-6	2	-17	-34	-4

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-5. Annual Mortality by All Factors for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super-imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	0	170,688	311,789	9,665	3,749	1,753,285	66,626	16,161	37,272	572	2,369,807
Alternative 1	0	171,160	333,426	9,304	3,662	1,799,292	57,690	10,479	29,913	597	2,415,523
Difference	0	472	21,637	-361	-87	46,006	-8,936	-5,682	-7,359	25	45,716
Percent Difference ³	0	0	7	-4	-2	3	-13	-35	-20	4	2
Water Year Types²											
Wet (32.5%)											
No Action Alternative	0	465,305	829,182	11,452	61	1,487,035	19	5,993	7	71	2,799,124
Alternative 1	0	464,856	854,662	11,983	61	1,479,843	549	4,386	494	114	2,816,949
Difference	0	-449	25,479	531	0	-7,192	530	-1,606	488	43	17,825
Percent Difference	0	0	3	5	1	0	2,784	-27	7,082	61	1
Above Normal (12.5%)											
No Action Alternative	0	24,311	338,436	9,179	167	1,810,328	54	1,307	18	84	2,183,885
Alternative 1	0	27,524	445,289	9,259	147	1,869,299	297	2,089	108	79	2,354,092
Difference	0	3,213	106,853	80	-19	58,971	243	782	90	-6	170,208
Percent Difference	0	13	32	1	-12	3	448	60	491	-7	8
Below Normal (17.5%)											
No Action Alternative	0	28,022	0	10,701	143	1,884,924	1,766	12,256	4,022	566	1,942,399
Alternative 1	0	30,282	0	11,214	62	1,985,320	1,247	8,090	2,635	488	2,039,338
Difference	0	2,261	0	513	-81	100,396	-519	-4,166	-1,386	-79	96,938
Percent Difference	0	8	0	5	-57	5	-29	-34	-34	-14	5
Dry (22.5%)											
No Action Alternative	0	28,946	0	5,999	570	1,894,042	21,850	17,140	16,111	835	1,985,493
Alternative 1	0	30,519	0	4,444	1,218	1,978,615	19,975	9,486	14,827	982	2,060,065
Difference	0	1,573	0	-1,556	648	84,573	-1,875	-7,654	-1,284	147	74,572
Percent Difference	0	5	0	-26	114	4	-9	-45	-8	18	4
Critical (15%)											
No Action Alternative	0	33,389	0	10,490	23,702	1,917,913	409,251	53,656	219,588	1,679	2,669,669
Alternative 1	0	29,837	0	8,597	22,262	1,947,073	351,747	34,946	172,942	1,627	2,569,032
Difference	0	-3,552	0	-1,893	-1,440	29,160	-57,504	-18,710	-46,646	-52	-100,637
Percent Difference	0	-11	0	-18	-6	2	-14	-35	-21	-3	-4

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-2-6. Annual Potential Production for Late Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	2,813,219
Alternative 3	2,812,234
Difference	-985
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	2,692,145
Alternative 3	2,691,402
Difference	-743
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	2,860,264
Alternative 3	2,810,515
Difference	-49,749
Percent Difference	-2
Below Normal (17.5%)	
No Action Alternative	2,982,412
Alternative 3	2,961,353
Difference	-21,059
Percent Difference	-1
Dry (22.5%)	
No Action Alternative	3,023,892
Alternative 3	3,012,660
Difference	-11,233
Percent Difference	0
Critical (15%)	
No Action Alternative	2,522,939
Alternative 3	2,600,856
Difference	77,917
Percent Difference	3
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-2-7. Annual Mortality by Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	492,142	1,757,035	82,787	37,844	120,631
Alternative 3	517,818	1,792,455	66,941	28,700	95,641
Difference	25,677	35,421	-15,845	-9,144	-24,990
Percent Difference ³	5	2	-19	-24	-21
Water Year Types²					
Wet (32.5%)					
No Action Alternative	1,305,939	1,487,095	6,012	78	6,089
Alternative 3	1,334,935	1,484,912	3,275	536	3,812
Difference	28,996	-2,184	-2,736	459	-2,278
Percent Difference	2	0	-46	590	-37
Above Normal (12.5%)					
No Action Alternative	371,926	1,810,494	1,361	103	1,464
Alternative 3	504,894	1,838,570	2,383	216	2,598
Difference	132,968	28,076	1,021	113	1,134
Percent Difference	36	2	75	110	77
Below Normal (17.5%)					
No Action Alternative	38,722	1,885,067	14,022	4,588	18,610
Alternative 3	39,609	1,946,219	10,333	2,164	12,497
Difference	887	61,152	-3,689	-2,424	-6,113
Percent Difference	2	3	-26	-53	-33
Dry (22.5%)					
No Action Alternative	34,945	1,894,612	38,990	16,946	55,936
Alternative 3	34,674	1,958,252	19,261	12,124	31,385
Difference	-271	63,640	-19,729	-4,822	-24,551
Percent Difference	-1	3	-51	-28	-44
Critical (15%)					
No Action Alternative	43,879	1,941,615	462,907	221,268	684,174
Alternative 3	40,798	1,992,284	396,247	169,277	565,524
Difference	-3,082	50,669	-66,660	-51,990	-118,650
Percent Difference	-7	3	-14	-23	-17

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-2-8. Annual Mortality by Cause for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	117,312	2,252,495	2,369,807
Alternative 3	96,645	2,309,269	2,405,915
Difference	-20,666	56,774	36,108
Percent Difference ³	-18	3	2
Water Year Types²			
Wet (32.5%)			
No Action Alternative	11,538	2,787,586	2,799,124
Alternative 3	13,133	2,810,525	2,823,658
Difference	1,595	22,940	24,535
Percent Difference	14	1	1
Above Normal (12.5%)			
No Action Alternative	9,419	2,174,466	2,183,885
Alternative 3	6,036	2,340,026	2,346,062
Difference	-3,382	165,560	162,178
Percent Difference	-36	8	7
Below Normal (17.5%)			
No Action Alternative	16,631	1,925,768	1,942,399
Alternative 3	13,519	1,984,806	1,998,326
Difference	-3,112	59,038	55,926
Percent Difference	-19	3	3
Dry (22.5%)			
No Action Alternative	44,530	1,940,964	1,985,493
Alternative 3	27,396	1,996,915	2,024,311
Difference	-17,134	55,952	38,818
Percent Difference	-38	3	2
Critical (15%)			
No Action Alternative	663,032	2,006,637	2,669,669
Alternative 3	553,950	2,044,656	2,598,606
Difference	-109,082	38,019	-71,063
Percent Difference	-16	2	-3
¹ Based on the 90-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the Annual average ⁴ Mortality values do not include base mortality			

Table B-2-9. Annual Mortality by Cause and Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)					Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature	Juvenile Habitat	
Long-term								
Full Simulation Period¹								
No Action Alternative	0	482,477	9,665	3,749	1,753,285	103,897	16,733	2,369,807
Alternative 3	0	509,000	8,818	3,126	1,789,329	84,700	10,941	2,405,915
Difference	0	26,523	-847	-623	36,043	-19,197	-5,793	36,108
Percent Difference ³	0	5	-9	-17	2	-18	-35	2
Water Year Types²								
Wet (32.5%)								
No Action Alternative	0	1,294,487	11,452	61	1,487,035	26	6,063	2,799,124
Alternative 3	0	1,322,789	12,146	61	1,484,851	927	2,885	2,823,658
Difference	0	28,302	694	0	-2,184	901	-3,178	24,535
Percent Difference	0	2	6	0	0	3,475	-52	1
Above Normal (12.5%)								
No Action Alternative	0	362,747	9,179	167	1,810,328	73	1,392	2,183,885
Alternative 3	0	499,275	5,619	31	1,838,539	386	2,212	2,346,062
Difference	0	136,528	-3,560	-136	28,212	314	821	162,178
Percent Difference	0	38	-39	-82	2	433	59	7
Below Normal (17.5%)								
No Action Alternative	0	28,022	10,701	143	1,884,924	5,787	12,822	1,942,399
Alternative 3	0	28,753	10,857	75	1,946,144	2,588	9,910	1,998,326
Difference	0	731	156	-68	61,220	-3,200	-2,913	55,926
Percent Difference	0	3	1	-47	3	-55	-23	3
Dry (22.5%)								
No Action Alternative	0	28,946	5,999	570	1,894,042	37,961	17,975	1,985,493
Alternative 3	0	30,082	4,592	188	1,958,065	22,616	8,769	2,024,311
Difference	0	1,136	-1,407	-382	64,022	-15,345	-9,206	38,818
Percent Difference	0	4	-23	-67	3	-40	-51	2
Critical (15%)								
No Action Alternative	0	33,389	10,490	23,702	1,917,913	628,839	55,335	2,669,669
Alternative 3	0	32,561	8,237	20,317	1,971,967	525,396	40,128	2,598,606
Difference	0	-829	-2,253	-3,386	54,055	-103,443	-15,207	-71,063
Percent Difference	0	-2	-21	-14	3	-16	-27	-3

1 Based on the 80-year simulation period
2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
3 Relative difference of the Annual average
4 Mortality values do not include base mortality

Table B-2-10. Annual Mortality by All Factors for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	0	170,688	311,789	9,665	3,749	1,753,285	66,626	16,161	37,272	572	2,369,807
Alternative 3	0	171,685	337,315	8,818	3,126	1,789,329	56,543	10,398	28,158	542	2,405,915
Difference	0	997	25,526	-847	-623	36,043	-10,083	-5,762	-9,114	-30	36,108
Percent Difference ³	0	1	8	-9	-17	2	-15	-36	-24	-5	2
Water Year Types²											
Wet (32.5%)											
No Action Alternative	0	465,305	829,182	11,452	61	1,487,035	19	5,993	7	71	2,799,124
Alternative 3	0	466,004	856,785	12,146	61	1,484,851	516	2,759	411	126	2,823,658
Difference	0	699	27,603	694	0	-2,184	497	-3,233	404	55	24,535
Percent Difference	0	0	3	6	0	0	2,610	-54	5,866	77	1
Above Normal (12.5%)											
No Action Alternative	0	24,311	338,436	9,179	167	1,810,328	54	1,307	18	84	2,183,885
Alternative 3	0	28,397	470,878	5,619	31	1,838,539	296	2,087	90	125	2,346,062
Difference	0	4,086	132,442	-3,560	-136	28,212	242	779	72	41	162,178
Percent Difference	0	17	39	-39	-82	2	446	60	392	49	7
Below Normal (17.5%)											
No Action Alternative	0	28,022	0	10,701	143	1,884,924	1,766	12,256	4,022	566	1,942,399
Alternative 3	0	28,753	0	10,857	75	1,946,144	823	9,510	1,765	400	1,998,326
Difference	0	731	0	156	-68	61,220	-943	-2,746	-2,257	-167	55,926
Percent Difference	0	3	0	1	-47	3	-53	-22	-56	-29	3
Dry (22.5%)											
No Action Alternative	0	28,946	0	5,999	570	1,894,042	21,850	17,140	16,111	835	1,985,493
Alternative 3	0	30,082	0	4,592	188	1,958,065	11,401	7,860	11,215	909	2,024,311
Difference	0	1,136	0	-1,407	-382	64,022	-10,449	-9,280	-4,896	74	38,818
Percent Difference	0	4	0	-23	-67	3	-48	-54	-30	9	2
Critical (15%)											
No Action Alternative	0	33,389	0	10,490	23,702	1,917,913	409,251	53,656	219,588	1,679	2,669,669
Alternative 3	0	32,561	0	8,237	20,317	1,971,967	357,527	38,720	167,870	1,408	2,598,606
Difference	0	-829	0	-2,253	-3,386	54,055	-51,725	-14,935	-51,719	-272	-71,063
Percent Difference	0	-2	0	-21	-14	3	-13	-28	-24	-16	-3

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-2-11. Annual Potential Production for Late Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	2,813,219
Alternative 5	2,805,566
Difference	-7,653
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	2,692,145
Alternative 5	2,700,194
Difference	8,049
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	2,860,264
Alternative 5	2,829,088
Difference	-31,176
Percent Difference	-1
Below Normal (17.5%)	
No Action Alternative	2,982,412
Alternative 5	2,951,992
Difference	-30,420
Percent Difference	-1
Dry (22.5%)	
No Action Alternative	3,023,892
Alternative 5	3,004,835
Difference	-19,057
Percent Difference	-1
Critical (15%)	
No Action Alternative	2,522,939
Alternative 5	2,544,537
Difference	21,598
Percent Difference	1
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-2-12. Annual Mortality by Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	492,142	1,757,035	82,787	37,844	120,631
Alternative 5	486,679	1,779,342	78,549	38,177	116,726
Difference	-5,463	22,307	-4,237	333	-3,904
Percent Difference ³	-1	1	-5	1	-3
Water Year Types²					
Wet (32.5%)					
No Action Alternative	1,305,939	1,487,095	6,012	78	6,089
Alternative 5	1,284,631	1,490,907	4,027	74	4,101
Difference	-21,308	3,812	-1,985	-4	-1,989
Percent Difference	-2	0	-33	-5	-33
Above Normal (12.5%)					
No Action Alternative	371,926	1,810,494	1,361	103	1,464
Alternative 5	385,985	1,859,656	1,357	82	1,439
Difference	14,059	49,162	-5	-21	-25
Percent Difference	4	3	0	-20	-2
Below Normal (17.5%)					
No Action Alternative	38,722	1,885,067	14,022	4,588	18,610
Alternative 5	39,141	1,943,539	13,998	4,481	18,480
Difference	419	58,471	-23	-107	-130
Percent Difference	1	3	0	-2	-1
Dry (22.5%)					
No Action Alternative	34,945	1,894,612	38,990	16,946	55,936
Alternative 5	34,298	1,930,739	31,905	14,697	46,602
Difference	-647	36,127	-7,085	-2,249	-9,334
Percent Difference	-2	2	-18	-13	-17
Critical (15%)					
No Action Alternative	43,879	1,941,615	462,907	221,268	684,174
Alternative 5	42,394	1,918,694	449,617	227,011	676,628
Difference	-1,485	-22,921	-13,290	5,743	-7,547
Percent Difference	-3	-1	-3	3	-1

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

⁵ Eggs mortality includes pre-spawn mortality

Table B-2-13. Annual Mortality by Cause for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	117,312	2,252,495	2,369,807
Alternative 5	115,323	2,267,424	2,382,747
Difference	-1,989	14,929	12,940
Percent Difference ³	-2	1	1
Water Year Types²			
Wet (32.5%)			
No Action Alternative	11,538	2,787,586	2,799,124
Alternative 5	11,470	2,768,169	2,779,639
Difference	-68	-19,417	-19,485
Percent Difference	-1	-1	-1
Above Normal (12.5%)			
No Action Alternative	9,419	2,174,466	2,183,885
Alternative 5	9,777	2,237,304	2,247,081
Difference	359	62,838	63,196
Percent Difference	4	3	3
Below Normal (17.5%)			
No Action Alternative	16,631	1,925,768	1,942,399
Alternative 5	16,938	1,984,222	2,001,160
Difference	307	58,454	58,760
Percent Difference	2	3	3
Dry (22.5%)			
No Action Alternative	44,530	1,940,964	1,985,493
Alternative 5	40,257	1,971,382	2,011,639
Difference	-4,273	30,419	26,146
Percent Difference	-10	2	1
Critical (15%)			
No Action Alternative	663,032	2,006,637	2,669,669
Alternative 5	655,672	1,982,044	2,637,716
Difference	-7,360	-24,593	-31,953
Percent Difference	-1	-1	-1

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-14. Annual Mortality by Cause and Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)					Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature	Juvenile Habitat	
Long-term								
Full Simulation Period¹								
No Action Alternative	0	482,477	9,665	3,749	1,753,285	103,897	16,733	2,369,807
Alternative 5	0	476,778	9,902	2,705	1,776,637	102,717	14,010	2,382,747
Difference	0	-5,699	236	-1,044	23,351	-1,181	-2,724	12,940
Percent Difference ³	0	-1	2	-28	1	-1	-16	1
Water Year Types²								
Wet (32.5%)								
No Action Alternative	0	1,294,487	11,452	61	1,487,035	26	6,063	2,799,124
Alternative 5	0	1,273,245	11,386	61	1,490,847	24	4,077	2,779,639
Difference	0	-21,242	-66	0	3,812	-2	-1,987	-19,485
Percent Difference	0	-2	-1	0	0	-8	-33	-1
Above Normal (12.5%)								
No Action Alternative	0	362,747	9,179	167	1,810,328	73	1,392	2,183,885
Alternative 5	0	376,400	9,586	142	1,859,515	50	1,389	2,247,081
Difference	0	13,653	406	-25	49,187	-23	-2	63,196
Percent Difference	0	4	4	-15	3	-31	0	3
Below Normal (17.5%)								
No Action Alternative	0	28,022	10,701	143	1,884,924	5,787	12,822	1,942,399
Alternative 5	0	28,128	11,014	147	1,943,392	5,777	12,702	2,001,160
Difference	0	106	313	4	58,468	-10	-120	58,760
Percent Difference	0	0	3	3	3	0	-1	3
Dry (22.5%)								
No Action Alternative	0	28,946	5,999	570	1,894,042	37,961	17,975	1,985,493
Alternative 5	0	28,043	6,255	761	1,929,979	33,241	13,361	2,011,639
Difference	0	-903	256	191	35,936	-4,720	-4,614	26,146
Percent Difference	0	-3	4	34	2	-12	-26	1
Critical (15%)								
No Action Alternative	0	33,389	10,490	23,702	1,917,913	628,839	55,335	2,669,669
Alternative 5	0	31,273	11,121	16,469	1,902,225	628,081	48,546	2,637,716
Difference	0	-2,116	631	-7,233	-15,688	-758	-6,789	-31,953
Percent Difference	0	-6	6	-31	-1	0	-12	-1

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-2-15. Annual Mortality by All Factors for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	0	170,688	311,789	9,665	3,749	1,753,285	66,626	16,161	37,272	572	2,369,807
Alternative 5	0	170,227	306,551	9,902	2,705	1,776,637	65,089	13,460	37,628	549	2,382,747
Difference	0	-461	-5,238	236	-1,044	23,351	-1,537	-2,700	356	-23	12,940
Percent Difference ³	0	0	-2	2	-28	1	-2	-17	1	-4	1
Water Year Types²											
Wet (32.5%)											
No Action Alternative	0	465,305	829,182	11,452	61	1,487,035	19	5,993	7	71	2,799,124
Alternative 5	0	465,569	807,677	11,386	61	1,490,847	18	4,009	6	68	2,779,639
Difference	0	264	-21,506	-66	0	3,812	-1	-1,984	-1	-3	-19,485
Percent Difference	0	0	-3	-1	0	0	-3	-33	-20	-4	-1
Above Normal (12.5%)											
No Action Alternative	0	24,311	338,436	9,179	167	1,810,328	54	1,307	18	84	2,183,885
Alternative 5	0	23,955	352,445	9,586	142	1,859,515	32	1,325	18	64	2,247,081
Difference	0	-356	14,009	406	-25	49,187	-22	18	-1	-20	63,196
Percent Difference	0	-1	4	4	-15	3	-41	1	-3	-24	3
Below Normal (17.5%)											
No Action Alternative	0	28,022	0	10,701	143	1,884,924	1,766	12,256	4,022	566	1,942,399
Alternative 5	0	28,128	0	11,014	147	1,943,392	1,852	12,147	3,925	556	2,001,160
Difference	0	106	0	313	4	58,468	86	-110	-96	-11	58,760
Percent Difference	0	0	0	3	3	3	5	-1	-2	-2	3
Dry (22.5%)											
No Action Alternative	0	28,946	0	5,999	570	1,894,042	21,850	17,140	16,111	835	1,985,493
Alternative 5	0	28,043	0	6,255	761	1,929,979	19,310	12,595	13,932	766	2,011,639
Difference	0	-903	0	256	191	35,936	-2,540	-4,545	-2,179	-70	26,146
Percent Difference	0	-3	0	4	34	2	-12	-27	-14	-8	1
Critical (15%)											
No Action Alternative	0	33,389	0	10,490	23,702	1,917,913	409,251	53,656	219,588	1,679	2,669,669
Alternative 5	0	31,273	0	11,121	16,469	1,902,225	402,734	46,883	225,348	1,663	2,637,716
Difference	0	-2,116	0	631	-7,233	-15,688	-6,517	-6,773	5,759	-16	-31,953
Percent Difference	0	-6	0	6	-31	-1	-2	-13	3	-1	-1

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table C-2-16. Annual Potential Production for Late Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	2,800,061
No Action Alternative	2,813,219
Difference	13,158
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	2,691,035
No Action Alternative	2,692,145
Difference	1,111
Percent Difference	0
Above Normal (12.5%)	
Second Basis of Comparison	2,802,912
No Action Alternative	2,860,264
Difference	57,352
Percent Difference	2
Below Normal (17.5%)	
Second Basis of Comparison	2,930,472
No Action Alternative	2,982,412
Difference	51,940
Percent Difference	2
Dry (22.5%)	
Second Basis of Comparison	2,976,338
No Action Alternative	3,023,892
Difference	47,554
Percent Difference	2
Critical (15%)	
Second Basis of Comparison	2,617,343
No Action Alternative	2,522,939
Difference	-94,404
Percent Difference	-4
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table C-2-17. Annual Mortality by Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	513,890	1,802,954	68,169	30,510	98,679
No Action Alternative	492,142	1,757,035	82,787	37,844	120,631
Difference	-21,748	-45,920	14,618	7,334	21,952
Percent Difference ³	-4	-3	21	24	22
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	1,331,500	1,479,904	4,935	609	5,544
No Action Alternative	1,305,939	1,487,095	6,012	78	6,089
Difference	-25,561	7,191	1,076	-531	545
Percent Difference	-2	0	22	-87	10
Above Normal (12.5%)					
Second Basis of Comparison	482,073	1,869,446	2,387	187	2,573
No Action Alternative	371,926	1,810,494	1,361	103	1,464
Difference	-110,146	-58,952	-1,025	-84	-1,109
Percent Difference	-23	-3	-43	-45	-43
Below Normal (17.5%)					
Second Basis of Comparison	41,496	1,985,382	9,337	3,123	12,460
No Action Alternative	38,722	1,885,067	14,022	4,588	18,610
Difference	-2,774	-100,315	4,685	1,465	6,150
Percent Difference	-7	-5	50	47	49
Dry (22.5%)					
Second Basis of Comparison	34,962	1,979,833	29,461	15,809	45,270
No Action Alternative	34,945	1,894,612	38,990	16,946	55,936
Difference	-17	-85,221	9,529	1,137	10,666
Percent Difference	0	-4	32	7	24
Critical (15%)					
Second Basis of Comparison	38,435	1,969,335	386,693	174,569	561,262
No Action Alternative	43,879	1,941,615	462,907	221,268	684,174
Difference	5,445	-27,720	76,214	46,699	122,912
Percent Difference	14	-1	20	27	22

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table C-2-18. Annual Mortality by Cause for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	100,569	2,314,954	2,415,523
No Action Alternative	117,312	2,252,495	2,369,807
Difference	16,743	-62,459	-45,716
Percent Difference ³	17	-3	-2
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	13,087	2,803,861	2,816,949
No Action Alternative	11,538	2,787,586	2,799,124
Difference	-1,549	-16,276	-17,825
Percent Difference	-12	-1	-1
Above Normal (12.5%)			
Second Basis of Comparison	9,812	2,344,280	2,354,092
No Action Alternative	9,419	2,174,466	2,183,885
Difference	-393	-169,814	-170,208
Percent Difference	-4	-7	-7
Below Normal (17.5%)			
Second Basis of Comparison	15,158	2,024,180	2,039,338
No Action Alternative	16,631	1,925,768	1,942,399
Difference	1,474	-98,412	-96,938
Percent Difference	10	-5	-5
Dry (22.5%)			
Second Basis of Comparison	40,463	2,019,602	2,060,065
No Action Alternative	44,530	1,940,964	1,985,493
Difference	4,067	-78,638	-74,572
Percent Difference	10	-4	-4
Critical (15%)			
Second Basis of Comparison	555,549	2,013,483	2,569,032
No Action Alternative	663,032	2,006,637	2,669,669
Difference	107,483	-6,846	100,637
Percent Difference	19	0	4

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table C-2-19. Annual Mortality by Cause and Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	0	504,586	9,304	3,662	1,799,292	87,603	11,076	2,415,523
No Action Alternative	0	482,477	9,665	3,749	1,753,285	103,897	16,733	2,369,807
Difference	0	-22,110	361	87	-46,006	16,294	5,657	-45,716
Percent Difference ³	0	-4	4	2	-3	19	51	-2
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	0	1,319,517	11,983	61	1,479,843	1,043	4,501	2,816,949
No Action Alternative	0	1,294,487	11,452	61	1,487,035	26	6,063	2,799,124
Difference	0	-25,030	-531	0	7,192	-1,018	1,563	-17,825
Percent Difference	0	-2	-4	-1	0	-98	35	-1
Above Normal (12.5%)								
Second Basis of Comparison	0	472,813	9,259	147	1,869,299	405	2,168	2,354,092
No Action Alternative	0	362,747	9,179	167	1,810,328	73	1,392	2,183,885
Difference	0	-110,066	-80	19	-58,971	-333	-776	-170,208
Percent Difference	0	-23	-1	13	-3	-82	-36	-7
Below Normal (17.5%)								
Second Basis of Comparison	0	30,282	11,214	62	1,985,320	3,882	8,578	2,039,338
No Action Alternative	0	28,022	10,701	143	1,884,924	5,787	12,822	1,942,399
Difference	0	-2,261	-513	81	-100,396	1,906	4,244	-96,938
Percent Difference	0	-7	-5	131	-5	49	49	-5
Dry (22.5%)								
Second Basis of Comparison	0	30,519	4,444	1,218	1,978,615	34,802	10,468	2,060,065
No Action Alternative	0	28,946	5,999	570	1,894,042	37,961	17,975	1,985,493
Difference	0	-1,573	1,556	-648	-84,573	3,159	7,508	-74,572
Percent Difference	0	-5	35	-53	-4	9	72	-4
Critical (15%)								
Second Basis of Comparison	0	29,837	8,597	22,262	1,947,073	524,689	36,573	2,569,032
No Action Alternative	0	33,389	10,490	23,702	1,917,913	628,839	55,335	2,669,669
Difference	0	3,552	1,893	1,440	-29,160	104,150	18,762	100,637
Percent Difference	0	12	22	6	-1	20	51	4

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table C-2-20. Annual Mortality by All Factors for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super-imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	0	171,160	333,426	9,304	3,662	1,799,292	57,690	10,479	29,913	597	2,415,523
No Action Alternative	0	170,688	311,789	9,665	3,749	1,753,285	66,626	16,161	37,272	572	2,369,807
Difference	0	-472	-21,637	361	87	-46,006	8,936	5,682	7,359	-25	-45,716
Percent Difference ³	0	0	-6	4	2	-3	15	54	25	-4	-2
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	0	464,856	854,662	11,983	61	1,479,843	549	4,386	494	114	2,816,949
No Action Alternative	0	465,305	829,182	11,452	61	1,487,035	19	5,993	7	71	2,799,124
Difference	0	449	-25,479	-531	0	7,192	-530	1,606	-488	-43	-17,825
Percent Difference	0	0	-3	-4	-1	0	-97	37	-99	-38	-1
Above Normal (12.5%)											
Second Basis of Comparison	0	27,524	445,289	9,259	147	1,869,299	297	2,089	108	79	2,354,092
No Action Alternative	0	24,311	338,436	9,179	167	1,810,328	54	1,307	18	84	2,183,885
Difference	0	-3,213	-106,853	-80	19	-58,971	-243	-782	-90	6	-170,208
Percent Difference	0	-12	-24	-1	13	-3	-82	-37	-83	7	-7
Below Normal (17.5%)											
Second Basis of Comparison	0	30,282	0	11,214	62	1,985,320	1,247	8,090	2,635	488	2,039,338
No Action Alternative	0	28,022	0	10,701	143	1,884,924	1,766	12,256	4,022	566	1,942,399
Difference	0	-2,261	0	-513	81	-100,396	519	4,166	1,386	79	-96,938
Percent Difference	0	-7	0	-5	131	-5	42	51	53	16	-5
Dry (22.5%)											
Second Basis of Comparison	0	30,519	0	4,444	1,218	1,978,615	19,975	9,486	14,827	982	2,060,065
No Action Alternative	0	28,946	0	5,999	570	1,894,042	21,850	17,140	16,111	835	1,985,493
Difference	0	-1,573	0	1,556	-648	-84,573	1,875	7,654	1,284	-147	-74,572
Percent Difference	0	-5	0	35	-53	-4	9	81	9	-15	-4
Critical (15%)											
Second Basis of Comparison	0	29,837	0	8,597	22,262	1,947,073	351,747	34,946	172,942	1,627	2,569,032
No Action Alternative	0	33,389	0	10,490	23,702	1,917,913	409,251	53,656	219,588	1,679	2,669,669
Difference	0	3,552	0	1,893	1,440	-29,160	57,504	18,710	46,646	52	100,637
Percent Difference	0	12	0	22	6	-1	16	54	27	3	4

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-21. Annual Potential Production for Late Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	2,800,061
Alternative 3	2,812,234
Difference	12,173
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	2,691,035
Alternative 3	2,691,402
Difference	367
Percent Difference	0
Above Normal (12.5%)	
Second Basis of Comparison	2,802,912
Alternative 3	2,810,515
Difference	7,603
Percent Difference	0
Below Normal (17.5%)	
Second Basis of Comparison	2,930,472
Alternative 3	2,961,353
Difference	30,881
Percent Difference	1
Dry (22.5%)	
Second Basis of Comparison	2,976,338
Alternative 3	3,012,660
Difference	36,322
Percent Difference	1
Critical (15%)	
Second Basis of Comparison	2,617,343
Alternative 3	2,600,856
Difference	-16,487
Percent Difference	-1
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-2-22. Annual Mortality by Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	513,890	1,802,954	68,169	30,510	98,679
Alternative 3	517,818	1,792,455	66,941	28,700	95,641
Difference	3,928	-10,499	-1,228	-1,811	-3,038
Percent Difference ³	1	-1	-2	-6	-3
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	1,331,500	1,479,904	4,935	609	5,544
Alternative 3	1,334,935	1,484,912	3,275	536	3,812
Difference	3,434	5,008	-1,660	-72	-1,732
Percent Difference	0	0	-34	-12	-31
Above Normal (12.5%)					
Second Basis of Comparison	482,073	1,869,446	2,387	187	2,573
Alternative 3	504,894	1,838,570	2,383	216	2,598
Difference	22,822	-30,877	-4	29	25
Percent Difference	5	-2	0	15	1
Below Normal (17.5%)					
Second Basis of Comparison	41,496	1,985,382	9,337	3,123	12,460
Alternative 3	39,609	1,946,219	10,333	2,164	12,497
Difference	-1,887	-39,163	996	-959	37
Percent Difference	-5	-2	11	-31	0
Dry (22.5%)					
Second Basis of Comparison	34,962	1,979,833	29,461	15,809	45,270
Alternative 3	34,674	1,958,252	19,261	12,124	31,385
Difference	-288	-21,580	-10,200	-3,685	-13,885
Percent Difference	-1	-1	-35	-23	-31
Critical (15%)					
Second Basis of Comparison	38,435	1,969,335	386,693	174,569	561,262
Alternative 3	40,798	1,992,284	396,247	169,277	565,524
Difference	2,363	22,949	9,554	-5,292	4,262
Percent Difference	6	1	2	-3	1

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-2-23. Annual Mortality by Cause for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	100,569	2,314,954	2,415,523
Alternative 3	96,645	2,309,269	2,405,915
Difference	-3,924	-5,685	-9,609
Percent Difference ³	-4	0	0
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	13,087	2,803,861	2,816,949
Alternative 3	13,133	2,810,525	2,823,658
Difference	45	6,664	6,710
Percent Difference	0	0	0
Above Normal (12.5%)			
Second Basis of Comparison	9,812	2,344,280	2,354,092
Alternative 3	6,036	2,340,026	2,346,062
Difference	-3,776	-4,254	-8,030
Percent Difference	-38	0	0
Below Normal (17.5%)			
Second Basis of Comparison	15,158	2,024,180	2,039,338
Alternative 3	13,519	1,984,806	1,998,326
Difference	-1,638	-39,374	-41,012
Percent Difference	-11	-2	-2
Dry (22.5%)			
Second Basis of Comparison	40,463	2,019,602	2,060,065
Alternative 3	27,396	1,996,915	2,024,311
Difference	-13,067	-22,686	-35,754
Percent Difference	-32	-1	-2
Critical (15%)			
Second Basis of Comparison	555,549	2,013,483	2,569,032
Alternative 3	553,950	2,044,656	2,598,606
Difference	-1,599	31,172	29,574
Percent Difference	0	2	1

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-24. Annual Mortality by Cause and Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	0	504,586	9,304	3,662	1,799,292	87,603	11,076	2,415,523
Alternative 3	0	509,000	8,818	3,126	1,789,329	84,700	10,941	2,405,915
Difference	0	4,414	-485	-536	-9,963	-2,903	-136	-9,609
Percent Difference ³	0	1	-5	-15	-1	-3	-1	0
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	0	1,319,517	11,983	61	1,479,843	1,043	4,501	2,816,949
Alternative 3	0	1,322,789	12,146	61	1,484,851	927	2,885	2,823,658
Difference	0	3,272	162	0	5,008	-117	-1,616	6,710
Percent Difference	0	0	1	0	0	-11	-36	0
Above Normal (12.5%)								
Second Basis of Comparison	0	472,813	9,259	147	1,869,299	405	2,168	2,354,092
Alternative 3	0	499,275	5,619	31	1,838,539	386	2,212	2,346,062
Difference	0	26,462	-3,640	-117	-30,760	-19	44	-8,030
Percent Difference	0	6	-39	-79	-2	-5	2	0
Below Normal (17.5%)								
Second Basis of Comparison	0	30,282	11,214	62	1,985,320	3,882	8,578	2,039,338
Alternative 3	0	28,753	10,857	75	1,946,144	2,588	9,910	1,998,326
Difference	0	-1,530	-357	13	-39,176	-1,294	1,332	-41,012
Percent Difference	0	-5	-3	21	-2	-33	16	-2
Dry (22.5%)								
Second Basis of Comparison	0	30,519	4,444	1,218	1,978,615	34,802	10,468	2,060,065
Alternative 3	0	30,082	4,592	188	1,958,065	22,616	8,769	2,024,311
Difference	0	-437	149	-1,030	-20,551	-12,186	-1,699	-35,754
Percent Difference	0	-1	3	-85	-1	-35	-16	-2
Critical (15%)								
Second Basis of Comparison	0	29,837	8,597	22,262	1,947,073	524,689	36,573	2,569,032
Alternative 3	0	32,561	8,237	20,317	1,971,967	525,396	40,128	2,598,606
Difference	0	2,723	-360	-1,946	24,894	707	3,555	29,574
Percent Difference	0	9	-4	-9	1	0	10	1

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-25. Annual Mortality by All Factors for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	0	171,160	333,426	9,304	3,662	1,799,292	57,690	10,479	29,913	597	2,415,523
Alternative 3	0	171,685	337,315	8,818	3,126	1,789,329	56,543	10,398	28,158	542	2,405,915
Difference	0	525	3,889	-485	-536	-9,963	-1,147	-80	-1,755	-55	-9,609
Percent Difference ³	0	0	1	-5	-15	-1	-2	-1	-6	-9	0
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	0	464,856	854,662	11,983	61	1,479,843	549	4,386	494	114	2,816,949
Alternative 3	0	466,004	856,785	12,146	61	1,484,851	516	2,759	411	126	2,823,658
Difference	0	1,149	2,123	162	0	5,008	-33	-1,627	-84	11	6,710
Percent Difference	0	0	0	1	0	0	-6	-37	-17	10	0
Above Normal (12.5%)											
Second Basis of Comparison	0	27,524	445,289	9,259	147	1,869,299	297	2,089	108	79	2,354,092
Alternative 3	0	28,397	470,878	5,619	31	1,838,539	296	2,087	90	125	2,346,062
Difference	0	873	25,589	-3,640	-117	-30,760	-1	-3	-18	47	-8,030
Percent Difference	0	3	6	-39	-79	-2	0	0	-17	60	0
Below Normal (17.5%)											
Second Basis of Comparison	0	30,282	0	11,214	62	1,985,320	1,247	8,090	2,635	488	2,039,338
Alternative 3	0	28,753	0	10,857	75	1,946,144	823	9,510	1,765	400	1,998,326
Difference	0	-1,530	0	-357	13	-39,176	-424	1,420	-871	-88	-41,012
Percent Difference	0	-5	0	-3	21	-2	-34	18	-33	-18	-2
Dry (22.5%)											
Second Basis of Comparison	0	30,519	0	4,444	1,218	1,978,615	19,975	9,486	14,827	982	2,060,065
Alternative 3	0	30,082	0	4,592	188	1,958,065	11,401	7,860	11,215	909	2,024,311
Difference	0	-437	0	149	-1,030	-20,551	-8,574	-1,626	-3,612	-73	-35,754
Percent Difference	0	-1	0	3	-85	-1	-43	-17	-24	-7	-2
Critical (15%)											
Second Basis of Comparison	0	29,837	0	8,597	22,262	1,947,073	351,747	34,946	172,942	1,627	2,569,032
Alternative 3	0	32,561	0	8,237	20,317	1,971,967	357,527	38,720	167,870	1,408	2,598,606
Difference	0	2,723	0	-360	-1,946	24,894	5,780	3,774	-5,072	-219	29,574
Percent Difference	0	9	0	-4	-9	1	2	11	-3	-13	1

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-26. Annual Potential Production for Late Fall-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	2,800,061
Alternative 5	2,805,566
Difference	5,506
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	2,691,035
Alternative 5	2,700,194
Difference	9,159
Percent Difference	0
Above Normal (12.5%)	
Second Basis of Comparison	2,802,912
Alternative 5	2,829,088
Difference	26,176
Percent Difference	1
Below Normal (17.5%)	
Second Basis of Comparison	2,930,472
Alternative 5	2,951,992
Difference	21,520
Percent Difference	1
Dry (22.5%)	
Second Basis of Comparison	2,976,338
Alternative 5	3,004,835
Difference	28,497
Percent Difference	1
Critical (15%)	
Second Basis of Comparison	2,617,343
Alternative 5	2,544,537
Difference	-72,807
Percent Difference	-3
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-2-27. Annual Mortality by Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	513,890	1,802,954	68,169	30,510	98,679
Alternative 5	486,679	1,779,342	78,549	38,177	116,726
Difference	-27,211	-23,612	10,380	7,667	18,047
Percent Difference ³	-5	-1	15	25	18
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	1,331,500	1,479,904	4,935	609	5,544
Alternative 5	1,284,631	1,490,907	4,027	74	4,101
Difference	-46,869	11,003	-909	-535	-1,443
Percent Difference	-4	1	-18	-88	-26
Above Normal (12.5%)					
Second Basis of Comparison	482,073	1,869,446	2,387	187	2,573
Alternative 5	385,985	1,859,656	1,357	82	1,439
Difference	-96,087	-9,790	-1,030	-105	-1,134
Percent Difference	-20	-1	-43	-56	-44
Below Normal (17.5%)					
Second Basis of Comparison	41,496	1,985,382	9,337	3,123	12,460
Alternative 5	39,141	1,943,539	13,998	4,481	18,480
Difference	-2,355	-41,843	4,662	1,358	6,020
Percent Difference	-6	-2	50	43	48
Dry (22.5%)					
Second Basis of Comparison	34,962	1,979,833	29,461	15,809	45,270
Alternative 5	34,298	1,930,739	31,905	14,697	46,602
Difference	-664	-49,093	2,444	-1,112	1,332
Percent Difference	-2	-2	8	-7	3
Critical (15%)					
Second Basis of Comparison	38,435	1,969,335	386,693	174,569	561,262
Alternative 5	42,394	1,918,694	449,617	227,011	676,628
Difference	3,960	-50,641	62,924	52,442	115,365
Percent Difference	10	-3	16	30	21

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-2-28. Annual Mortality by Cause for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	100,569	2,314,954	2,415,523
Alternative 5	115,323	2,267,424	2,382,747
Difference	14,754	-47,530	-32,776
Percent Difference ³	15	-2	-1
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	13,087	2,803,861	2,816,949
Alternative 5	11,470	2,768,169	2,779,639
Difference	-1,617	-35,692	-37,310
Percent Difference	-12	-1	-1
Above Normal (12.5%)			
Second Basis of Comparison	9,812	2,344,280	2,354,092
Alternative 5	9,777	2,237,304	2,247,081
Difference	-35	-106,977	-107,012
Percent Difference	0	-5	-5
Below Normal (17.5%)			
Second Basis of Comparison	15,158	2,024,180	2,039,338
Alternative 5	16,938	1,984,222	2,001,160
Difference	1,780	-39,958	-38,178
Percent Difference	12	-2	-2
Dry (22.5%)			
Second Basis of Comparison	40,463	2,019,602	2,060,065
Alternative 5	40,257	1,971,382	2,011,639
Difference	-206	-48,219	-48,426
Percent Difference	-1	-2	-2
Critical (15%)			
Second Basis of Comparison	555,549	2,013,483	2,569,032
Alternative 5	655,672	1,982,044	2,637,716
Difference	100,123	-31,439	68,684
Percent Difference	18	-2	3

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-29. Annual Mortality by Cause and Life Stage for Late Fall-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	0	504,586	9,304	3,662	1,799,292	87,603	11,076	2,415,523
Alternative 5	0	476,778	9,902	2,705	1,776,637	102,717	14,010	2,382,747
Difference	0	-27,809	598	-958	-22,655	15,114	2,934	-32,776
Percent Difference ³	0	-6	6	-26	-1	17	26	-1
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	0	1,319,517	11,983	61	1,479,843	1,043	4,501	2,816,949
Alternative 5	0	1,273,245	11,386	61	1,490,847	24	4,077	2,779,639
Difference	0	-46,272	-597	0	11,003	-1,020	-424	-37,310
Percent Difference	0	-4	-5	-1	1	-98	-9	-1
Above Normal (12.5%)								
Second Basis of Comparison	0	472,813	9,259	147	1,869,299	405	2,168	2,354,092
Alternative 5	0	376,400	9,586	142	1,859,515	50	1,389	2,247,081
Difference	0	-96,413	326	-6	-9,784	-355	-779	-107,012
Percent Difference	0	-20	4	-4	-1	-88	-36	-5
Below Normal (17.5%)								
Second Basis of Comparison	0	30,282	11,214	62	1,985,320	3,882	8,578	2,039,338
Alternative 5	0	28,128	11,014	147	1,943,392	5,777	12,702	2,001,160
Difference	0	-2,155	-200	85	-41,928	1,896	4,124	-38,178
Percent Difference	0	-7	-2	137	-2	49	48	-2
Dry (22.5%)								
Second Basis of Comparison	0	30,519	4,444	1,218	1,978,615	34,802	10,468	2,060,065
Alternative 5	0	28,043	6,255	761	1,929,979	33,241	13,361	2,011,639
Difference	0	-2,476	1,812	-457	-48,637	-1,561	2,893	-48,426
Percent Difference	0	-8	41	-38	-2	-4	28	-2
Critical (15%)								
Second Basis of Comparison	0	29,837	8,597	22,262	1,947,073	524,689	36,573	2,569,032
Alternative 5	0	31,273	11,121	16,469	1,902,225	628,081	48,546	2,637,716
Difference	0	1,436	2,524	-5,793	-44,848	103,392	11,973	68,684
Percent Difference	0	5	29	-26	-2	20	33	3

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-2-30. Annual Mortality by All Factors for Late Fall-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	0	171,160	333,426	9,304	3,662	1,799,292	57,690	10,479	29,913	597	2,415,523
Alternative 5	0	170,227	306,551	9,902	2,705	1,776,637	65,089	13,460	37,628	549	2,382,747
Difference	0	-933	-26,876	598	-958	-22,655	7,399	2,982	7,715	-48	-32,776
Percent Difference ³	0	-1	-8	6	-26	-1	13	28	26	-8	-1
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	0	464,856	854,662	11,983	61	1,479,843	549	4,386	494	114	2,816,949
Alternative 5	0	465,569	807,677	11,386	61	1,490,847	18	4,009	6	68	2,779,639
Difference	0	713	-46,985	-597	0	11,003	-531	-378	-489	-46	-37,310
Percent Difference	0	0	-5	-5	-1	1	-97	-9	-99	-40	-1
Above Normal (12.5%)											
Second Basis of Comparison	0	27,524	445,289	9,259	147	1,869,299	297	2,089	108	79	2,354,092
Alternative 5	0	23,955	352,445	9,586	142	1,859,515	32	1,325	18	64	2,247,081
Difference	0	-3,569	-92,844	326	-6	-9,784	-265	-765	-90	-14	-107,012
Percent Difference	0	-13	-21	4	-4	-1	-89	-37	-84	-18	-5
Below Normal (17.5%)											
Second Basis of Comparison	0	30,282	0	11,214	62	1,985,320	1,247	8,090	2,635	488	2,039,338
Alternative 5	0	28,128	0	11,014	147	1,943,392	1,852	12,147	3,925	556	2,001,160
Difference	0	-2,155	0	-200	85	-41,928	605	4,056	1,290	68	-38,178
Percent Difference	0	-7	0	-2	137	-2	49	50	49	14	-2
Dry (22.5%)											
Second Basis of Comparison	0	30,519	0	4,444	1,218	1,978,615	19,975	9,486	14,827	982	2,060,065
Alternative 5	0	28,043	0	6,255	761	1,929,979	19,310	12,595	13,932	766	2,011,639
Difference	0	-2,476	0	1,812	-457	-48,637	-665	3,109	-896	-216	-48,426
Percent Difference	0	-8	0	41	-38	-2	-3	33	-6	-22	-2
Critical (15%)											
Second Basis of Comparison	0	29,837	0	8,597	22,262	1,947,073	351,747	34,946	172,942	1,627	2,569,032
Alternative 5	0	31,273	0	11,121	16,469	1,902,225	402,734	46,883	225,348	1,663	2,637,716
Difference	0	1,436	0	2,524	-5,793	-44,848	50,987	11,937	52,405	36	68,684
Percent Difference	0	5	0	29	-26	-2	14	34	30	2	3

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

1 **B.3. Spring-Run Chinook Salmon**


2

Figure B-3-1. Annual Potential Production for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-2. Annual Mortality for Spring-Run Chinook Salmon - Eggs




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-3. Annual Mortality for Spring-Run Chinook Salmon - Fry




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-4. Annual Mortality for Spring-Run Chinook Salmon - Pre-Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-5. Annual Mortality for Spring-Run Chinook Salmon - Immature Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-6. Annual Mortality for Spring-Run Chinook Salmon - Pre- & Immature Smolts




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-7. Annual Mortality for Spring-Run Chinook Salmon - All Lifestages




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-8. Incubation - Habitat based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-9. Super-imposition - Habitat based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-10. Fry - Habitat based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-11. Pre-smolt - Habitat based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-12. Immature Smolt - Habitat based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-13. Total Habitat based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-14. Pre-Spawn Mortality - Temperature based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-15. Eggs - Temperature based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-16. Fry - Temperature based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-17. Pre-smolt - Temperature based Annual Mortality for Spring-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-18. Immature Smolt - Temperature based Annual Mortality for Spring-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-3-19. Total Temperature based Annual Mortality for Spring-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Table B-3-1. Annual Potential Production for Spring-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	402,980
Alternative 1	410,722
Difference	7,742
Percent Difference ³	2
Water Year Types²	
Wet (32.5%)	
No Action Alternative	442,676
Alternative 1	449,832
Difference	7,156
Percent Difference	2
Above Normal (12.5%)	
No Action Alternative	362,537
Alternative 1	367,591
Difference	5,054
Percent Difference	1
Below Normal (17.5%)	
No Action Alternative	428,569
Alternative 1	426,491
Difference	-2,078
Percent Difference	0
Dry (22.5%)	
No Action Alternative	405,967
Alternative 1	403,012
Difference	-2,955
Percent Difference	-1
Critical (15%)	
No Action Alternative	316,344
Alternative 1	355,097
Difference	38,753
Percent Difference	12
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-3-2. Annual Mortality by Life Stage for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	169,230	2,282	0	0	0
Alternative 1	149,155	2,453	0	0	0
Difference	-20,075	171	0	0	0
Percent Difference ³	-12	7	0	0	0
Water Year Types²					
Wet (32.5%)					
No Action Alternative	54,929	2,217	0	0	0
Alternative 1	38,874	2,303	0	0	0
Difference	-16,055	86	0	0	0
Percent Difference	-29	4	0	0	0
Above Normal (12.5%)					
No Action Alternative	275,059	1,955	0	0	0
Alternative 1	256,999	2,360	0	0	0
Difference	-18,059	406	0	0	0
Percent Difference	-7	21	0	0	0
Below Normal (17.5%)					
No Action Alternative	108,811	2,619	0	0	0
Alternative 1	110,617	2,763	0	0	0
Difference	1,806	144	0	0	0
Percent Difference	2	5	0	0	0
Dry (22.5%)					
No Action Alternative	170,290	2,608	0	0	0
Alternative 1	175,971	2,682	0	0	0
Difference	5,681	73	0	0	0
Percent Difference	3	3	0	0	0
Critical (15%)					
No Action Alternative	397,589	1,814	0	0	0
Alternative 1	302,962	2,151	0	0	0
Difference	-94,627	337	0	0	0
Percent Difference	-24	19	0	0	0

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-3-3. Annual Mortality by Cause for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	167,192	4,321	171,512
Alternative 1	146,922	4,686	151,608
Difference	-20,270	366	-19,904
Percent Difference ³	-12	8	-12
Water Year Types²			
Wet (32.5%)			
No Action Alternative	53,038	4,108	57,146
Alternative 1	36,709	4,468	41,178
Difference	-16,329	360	-15,969
Percent Difference	-31	9	-28
Above Normal (12.5%)			
No Action Alternative	274,408	2,606	277,013
Alternative 1	256,534	2,826	259,360
Difference	-17,874	221	-17,653
Percent Difference	-7	8	-6
Below Normal (17.5%)			
No Action Alternative	107,177	4,253	111,431
Alternative 1	108,800	4,580	113,380
Difference	1,623	327	1,949
Percent Difference	2	8	2
Dry (22.5%)			
No Action Alternative	167,873	5,025	172,898
Alternative 1	173,420	5,232	178,652
Difference	5,547	207	5,754
Percent Difference	3	4	3
Critical (15%)			
No Action Alternative	394,171	5,232	399,403
Alternative 1	299,101	6,012	305,113
Difference	-95,070	780	-94,290
Percent Difference	-24	15	-24

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-4. Annual Mortality by Cause and Life Stage for Spring-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	47,267	2,039	119,924	1	2,282	0	0	171,512
Alternative 1	38,621	2,233	108,301	0	2,453	0	0	151,608
Difference	-8,646	194	-11,623	-1	172	0	0	-19,904
Percent Difference ³	-18	10	-10	-100	8	0	0	-12
Water Year Types²								
Wet (32.5%)								
No Action Alternative	340	1,893	52,697	2	2,215	0	0	57,146
Alternative 1	260	2,165	36,450	0	2,303	0	0	41,178
Difference	-80	272	-16,247	-2	88	0	0	-15,969
Percent Difference	-24	14	-31	-100	4	0	0	-28
Above Normal (12.5%)								
No Action Alternative	151,449	651	122,959	0	1,955	0	0	277,013
Alternative 1	99,868	466	156,666	0	2,360	0	0	259,360
Difference	-51,581	-185	33,707	0	406	0	0	-17,653
Percent Difference	-34	-28	27	0	21	0	0	-6
Below Normal (17.5%)								
No Action Alternative	63,840	1,634	43,337	0	2,619	0	0	111,431
Alternative 1	66,585	1,818	42,215	0	2,763	0	0	113,380
Difference	2,744	183	-1,122	0	144	0	0	1,949
Percent Difference	4	11	-3	0	5	0	0	2
Dry (22.5%)								
No Action Alternative	37,718	2,417	130,155	0	2,608	0	0	172,898
Alternative 1	34,417	2,551	139,003	0	2,682	0	0	178,652
Difference	-3,301	134	8,847	0	73	0	0	5,754
Percent Difference	-9	6	7	0	3	0	0	3
Critical (15%)								
No Action Alternative	57,112	3,419	337,059	0	1,814	0	0	399,403
Alternative 1	44,378	3,862	254,723	0	2,151	0	0	305,113
Difference	-12,734	443	-82,336	0	337	0	0	-94,290
Percent Difference	-22	13	-24	0	19	0	0	-24

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-5. Annual Mortality by All Factors for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	47,267	2,039	0	119,924	1	2,282	0	0	0	0	171,512
Alternative 1	38,621	2,233	0	108,301	0	2,453	0	0	0	0	151,608
Difference	-8,646	194	0	-11,623	-1	172	0	0	0	0	-19,904
Percent Difference ³	-18	10	0	-10	-100	8	0	0	0	0	-12
Water Year Types²											
Wet (32.5%)											
No Action Alternative	340	1,893	0	52,697	2	2,215	0	0	0	0	57,146
Alternative 1	260	2,165	0	36,450	0	2,303	0	0	0	0	41,178
Difference	-80	272	0	-16,247	-2	88	0	0	0	0	-15,969
Percent Difference	-24	14	0	-31	-100	4	0	0	0	0	-28
Above Normal (12.5%)											
No Action Alternative	151,449	651	0	122,959	0	1,955	0	0	0	0	277,013
Alternative 1	99,868	466	0	156,666	0	2,360	0	0	0	0	259,360
Difference	-51,581	-185	0	33,707	0	406	0	0	0	0	-17,653
Percent Difference	-34	-28	0	27	0	21	0	0	0	0	-6
Below Normal (17.5%)											
No Action Alternative	63,840	1,634	0	43,337	0	2,619	0	0	0	0	111,431
Alternative 1	66,585	1,818	0	42,215	0	2,763	0	0	0	0	113,380
Difference	2,744	183	0	-1,122	0	144	0	0	0	0	1,949
Percent Difference	4	11	0	-3	0	5	0	0	0	0	2
Dry (22.5%)											
No Action Alternative	37,718	2,417	0	130,155	0	2,608	0	0	0	0	172,898
Alternative 1	34,417	2,551	0	139,003	0	2,682	0	0	0	0	178,652
Difference	-3,301	134	0	8,847	0	73	0	0	0	0	5,754
Percent Difference	-9	6	0	7	0	3	0	0	0	0	3
Critical (15%)											
No Action Alternative	57,112	3,419	0	337,059	0	1,814	0	0	0	0	399,403
Alternative 1	44,378	3,862	0	254,723	0	2,151	0	0	0	0	305,113
Difference	-12,734	443	0	-82,336	0	337	0	0	0	0	-94,290
Percent Difference	-22	13	0	-24	0	19	0	0	0	0	-24

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-3-6. Annual Potential Production for Spring-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	402,980
Alternative 3	409,813
Difference	6,832
Percent Difference ³	2
Water Year Types²	
Wet (32.5%)	
No Action Alternative	442,676
Alternative 3	453,743
Difference	11,067
Percent Difference	2
Above Normal (12.5%)	
No Action Alternative	362,537
Alternative 3	368,403
Difference	5,866
Percent Difference	2
Below Normal (17.5%)	
No Action Alternative	428,569
Alternative 3	427,631
Difference	-938
Percent Difference	0
Dry (22.5%)	
No Action Alternative	405,967
Alternative 3	410,542
Difference	4,575
Percent Difference	1
Critical (15%)	
No Action Alternative	316,344
Alternative 3	327,260
Difference	10,915
Percent Difference	3
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-3-7. Annual Mortality by Life Stage for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	169,230	2,282	0	0	0
Alternative 3	150,290	2,435	0	0	0
Difference	-18,940	153	0	0	0
Percent Difference ³	-11	7	0	0	0
Water Year Types²					
Wet (32.5%)					
No Action Alternative	54,929	2,217	0	0	0
Alternative 3	29,787	2,271	0	0	0
Difference	-25,142	54	0	0	0
Percent Difference	-46	2	0	0	0
Above Normal (12.5%)					
No Action Alternative	275,059	1,955	0	0	0
Alternative 3	257,573	2,190	0	0	0
Difference	-17,485	236	0	0	0
Percent Difference	-6	12	0	0	0
Below Normal (17.5%)					
No Action Alternative	108,811	2,619	0	0	0
Alternative 3	107,671	2,858	0	0	0
Difference	-1,140	239	0	0	0
Percent Difference	-1	9	0	0	0
Dry (22.5%)					
No Action Alternative	170,290	2,608	0	0	0
Alternative 3	156,331	2,731	0	0	0
Difference	-13,959	123	0	0	0
Percent Difference	-8	5	0	0	0
Critical (15%)					
No Action Alternative	397,589	1,814	0	0	0
Alternative 3	362,639	2,060	0	0	0
Difference	-34,950	247	0	0	0
Percent Difference	-9	14	0	0	0

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-3-8. Annual Mortality by Cause for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	167,192	4,321	171,512
Alternative 3	148,223	4,502	152,726
Difference	-18,968	182	-18,786
Percent Difference ³	-11	4	-11
Water Year Types²			
Wet (32.5%)			
No Action Alternative	53,038	4,108	57,146
Alternative 3	27,591	4,467	32,057
Difference	-25,448	359	-25,089
Percent Difference	-48	9	-44
Above Normal (12.5%)			
No Action Alternative	274,408	2,606	277,013
Alternative 3	257,166	2,597	259,763
Difference	-17,242	-8	-17,250
Percent Difference	-6	0	-6
Below Normal (17.5%)			
No Action Alternative	107,177	4,253	111,431
Alternative 3	105,832	4,697	110,529
Difference	-1,345	444	-901
Percent Difference	-1	10	-1
Dry (22.5%)			
No Action Alternative	167,873	5,025	172,898
Alternative 3	154,048	5,014	159,062
Difference	-13,825	-11	-13,836
Percent Difference	-8	0	-8
Critical (15%)			
No Action Alternative	394,171	5,232	399,403
Alternative 3	359,528	5,172	364,700
Difference	-34,643	-60	-34,703
Percent Difference	-9	-1	-9

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-9. Annual Mortality by Cause and Life Stage for Spring-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	47,267	2,039	119,924	1	2,282	0	0	171,512
Alternative 3	37,164	2,067	111,060	0	2,435	0	0	152,726
Difference	-10,103	28	-8,864	-1	154	0	0	-18,786
Percent Difference ³	-21	1	-7	-100	7	0	0	-11
Water Year Types²								
Wet (32.5%)								
No Action Alternative	340	1,893	52,697	2	2,215	0	0	57,146
Alternative 3	189	2,196	27,402	0	2,271	0	0	32,057
Difference	-151	303	-25,295	-2	56	0	0	-25,089
Percent Difference	-44	16	-48	-100	3	0	0	-44
Above Normal (12.5%)								
No Action Alternative	151,449	651	122,959	0	1,955	0	0	277,013
Alternative 3	104,829	407	152,337	0	2,190	0	0	259,763
Difference	-46,620	-244	29,379	0	236	0	0	-17,250
Percent Difference	-31	-37	24	0	12	0	0	-6
Below Normal (17.5%)								
No Action Alternative	63,840	1,634	43,337	0	2,619	0	0	111,431
Alternative 3	62,085	1,839	43,747	0	2,858	0	0	110,529
Difference	-1,755	205	410	0	239	0	0	-901
Percent Difference	-3	13	1	0	9	0	0	-1
Dry (22.5%)								
No Action Alternative	37,718	2,417	130,155	0	2,608	0	0	172,898
Alternative 3	28,700	2,282	125,348	0	2,731	0	0	159,062
Difference	-9,018	-134	-4,807	0	123	0	0	-13,836
Percent Difference	-24	-6	-4	0	5	0	0	-8
Critical (15%)								
No Action Alternative	57,112	3,419	337,059	0	1,814	0	0	399,403
Alternative 3	44,510	3,112	315,018	0	2,060	0	0	364,700
Difference	-12,602	-307	-22,041	0	247	0	0	-34,703
Percent Difference	-22	-9	-7	0	14	0	0	-9

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-10. Annual Mortality by All Factors for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
No Action Alternative	47,267	2,039	0	119,924	1	2,282	0	0	0	0	171,512
Alternative 3	37,164	2,067	0	111,060	0	2,435	0	0	0	0	152,726
Difference	-10,103	28	0	-8,864	-1	154	0	0	0	0	-18,786
Percent Difference ³	-21	1	0	-7	-100	7	0	0	0	0	-11
Water Year Types²											
Wet (32.5%)											
No Action Alternative	340	1,893	0	52,697	2	2,215	0	0	0	0	57,146
Alternative 3	189	2,196	0	27,402	0	2,271	0	0	0	0	32,057
Difference	-151	303	0	-25,295	-2	56	0	0	0	0	-25,089
Percent Difference	-44	16	0	-48	-100	3	0	0	0	0	-44
Above Normal (12.5%)											
No Action Alternative	151,449	651	0	122,959	0	1,955	0	0	0	0	277,013
Alternative 3	104,829	407	0	152,337	0	2,190	0	0	0	0	259,763
Difference	-46,620	-244	0	29,379	0	236	0	0	0	0	-17,250
Percent Difference	-31	-37	0	24	0	12	0	0	0	0	-6
Below Normal (17.5%)											
No Action Alternative	63,840	1,634	0	43,337	0	2,619	0	0	0	0	111,431
Alternative 3	62,085	1,839	0	43,747	0	2,858	0	0	0	0	110,529
Difference	-1,755	205	0	410	0	239	0	0	0	0	-901
Percent Difference	-3	13	0	1	0	9	0	0	0	0	-1
Dry (22.5%)											
No Action Alternative	37,718	2,417	0	130,155	0	2,608	0	0	0	0	172,898
Alternative 3	28,700	2,282	0	125,348	0	2,731	0	0	0	0	159,062
Difference	-9,018	-134	0	-4,807	0	123	0	0	0	0	-13,836
Percent Difference	-24	-6	0	-4	0	5	0	0	0	0	-8
Critical (15%)											
No Action Alternative	57,112	3,419	0	337,059	0	1,814	0	0	0	0	399,403
Alternative 3	44,510	3,112	0	315,018	0	2,060	0	0	0	0	364,700
Difference	-12,602	-307	0	-22,041	0	247	0	0	0	0	-34,703
Percent Difference	-22	-9	0	-7	0	14	0	0	0	0	-9

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-3-11. Annual Potential Production for Spring-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	402,980
Alternative 5	401,678
Difference	-1,302
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	442,676
Alternative 5	441,971
Difference	-705
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	362,537
Alternative 5	363,460
Difference	923
Percent Difference	0
Below Normal (17.5%)	
No Action Alternative	428,569
Alternative 5	428,206
Difference	-363
Percent Difference	0
Dry (22.5%)	
No Action Alternative	405,967
Alternative 5	407,290
Difference	1,323
Percent Difference	0
Critical (15%)	
No Action Alternative	316,344
Alternative 5	306,861
Difference	-9,484
Percent Difference	-3
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-3-12. Annual Mortality by Life Stage for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	169,230	2,282	0	0	0
Alternative 5	171,978	2,371	0	0	0
Difference	2,748	89	0	0	0
Percent Difference ³	2	4	0	0	0
Water Year Types²					
Wet (32.5%)					
No Action Alternative	54,929	2,217	0	0	0
Alternative 5	57,192	2,203	0	0	0
Difference	2,263	-14	0	0	0
Percent Difference	4	-1	0	0	0
Above Normal (12.5%)					
No Action Alternative	275,059	1,955	0	0	0
Alternative 5	271,916	1,980	0	0	0
Difference	-3,143	26	0	0	0
Percent Difference	-1	1	0	0	0
Below Normal (17.5%)					
No Action Alternative	108,811	2,619	0	0	0
Alternative 5	108,195	2,925	0	0	0
Difference	-616	306	0	0	0
Percent Difference	-1	12	0	0	0
Dry (22.5%)					
No Action Alternative	170,290	2,608	0	0	0
Alternative 5	166,496	2,666	0	0	0
Difference	-3,794	57	0	0	0
Percent Difference	-2	2	0	0	0
Critical (15%)					
No Action Alternative	397,589	1,814	0	0	0
Alternative 5	420,039	1,972	0	0	0
Difference	22,449	159	0	0	0
Percent Difference	6	9	0	0	0

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-3-13. Annual Mortality by Cause for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	167,192	4,321	171,512
Alternative 5	170,196	4,153	174,349
Difference	3,004	-167	2,837
Percent Difference ³	2	-4	2
Water Year Types²			
Wet (32.5%)			
No Action Alternative	53,038	4,108	57,146
Alternative 5	55,390	4,005	59,395
Difference	2,351	-103	2,249
Percent Difference	4	-2	4
Above Normal (12.5%)			
No Action Alternative	274,408	2,606	277,013
Alternative 5	271,280	2,616	273,896
Difference	-3,128	11	-3,117
Percent Difference	-1	0	-1
Below Normal (17.5%)			
No Action Alternative	107,177	4,253	111,431
Alternative 5	106,681	4,439	111,120
Difference	-496	186	-310
Percent Difference	0	4	0
Dry (22.5%)			
No Action Alternative	167,873	5,025	172,898
Alternative 5	164,607	4,554	169,161
Difference	-3,266	-471	-3,737
Percent Difference	-2	-9	-2
Critical (15%)			
No Action Alternative	394,171	5,232	399,403
Alternative 5	417,191	4,820	422,011
Difference	23,020	-412	22,608
Percent Difference	6	-8	6

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-14. Annual Mortality by Cause and Life Stage for Spring-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Temperature	Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat				
Long-term									
Full Simulation Period¹									
No Action Alternative	47,267	2,039	119,924	1	2,282	0	0	171,512	
Alternative 5	44,327	1,783	125,868	0	2,371	0	0	174,349	
Difference	-2,940	-256	5,944	0	89	0	0	2,837	
Percent Difference ³	-6	-13	5	-52	4	0	0	2	
Water Year Types²									
Wet (32.5%)									
No Action Alternative	340	1,893	52,697	2	2,215	0	0	57,146	
Alternative 5	608	1,803	54,781	1	2,203	0	0	59,395	
Difference	268	-90	2,084	-1	-13	0	0	2,249	
Percent Difference	79	-5	4	-57	-1	0	0	4	
Above Normal (12.5%)									
No Action Alternative	151,449	651	122,959	0	1,955	0	0	277,013	
Alternative 5	125,685	636	145,595	0	1,980	0	0	273,896	
Difference	-25,764	-15	22,636	0	26	0	0	-3,117	
Percent Difference	-17	-2	18	0	1	0	0	-1	
Below Normal (17.5%)									
No Action Alternative	63,840	1,634	43,337	0	2,619	0	0	111,431	
Alternative 5	53,122	1,514	53,559	0	2,925	0	0	111,120	
Difference	-10,718	-120	10,222	0	306	0	0	-310	
Percent Difference	-17	-7	24	0	12	0	0	0	
Dry (22.5%)									
No Action Alternative	37,718	2,417	130,155	0	2,608	0	0	172,898	
Alternative 5	37,450	1,889	127,157	0	2,666	0	0	169,161	
Difference	-268	-528	-2,998	0	57	0	0	-3,737	
Percent Difference	-1	-22	-2	0	2	0	0	-2	
Critical (15%)									
No Action Alternative	57,112	3,419	337,059	0	1,814	0	0	399,403	
Alternative 5	71,310	2,848	345,881	0	1,972	0	0	422,011	
Difference	14,198	-571	8,822	0	158	0	0	22,608	
Percent Difference	25	-17	3	0	9	0	0	6	

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-15. Annual Mortality by All Factors for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super-imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	47,267	2,039	0	119,924	1	2,282	0	0	0	0	171,512
Alternative 5	44,327	1,783	0	125,868	0	2,371	0	0	0	0	174,349
Difference	-2,940	-256	0	5,944	0	89	0	0	0	0	2,837
Percent Difference ³	-6	-13	0	5	-52	4	0	0	0	0	2
Water Year Types²											
Wet (32.5%)											
No Action Alternative	340	1,893	0	52,697	2	2,215	0	0	0	0	57,146
Alternative 5	608	1,803	0	54,781	1	2,203	0	0	0	0	59,395
Difference	268	-90	0	2,084	-1	-13	0	0	0	0	2,249
Percent Difference	79	-5	0	4	-57	-1	0	0	0	0	4
Above Normal (12.5%)											
No Action Alternative	151,449	651	0	122,959	0	1,955	0	0	0	0	277,013
Alternative 5	125,685	636	0	145,595	0	1,980	0	0	0	0	273,896
Difference	-25,764	-15	0	22,636	0	26	0	0	0	0	-3,117
Percent Difference	-17	-2	0	18	0	1	0	0	0	0	-1
Below Normal (17.5%)											
No Action Alternative	63,840	1,634	0	43,337	0	2,619	0	0	0	0	111,431
Alternative 5	53,122	1,514	0	53,559	0	2,925	0	0	0	0	111,120
Difference	-10,718	-120	0	10,222	0	306	0	0	0	0	-310
Percent Difference	-17	-7	0	24	0	12	0	0	0	0	0
Dry (22.5%)											
No Action Alternative	37,718	2,417	0	130,155	0	2,608	0	0	0	0	172,898
Alternative 5	37,450	1,889	0	127,157	0	2,666	0	0	0	0	169,161
Difference	-268	-528	0	-2,998	0	57	0	0	0	0	-3,737
Percent Difference	-1	-22	0	-2	0	2	0	0	0	0	-2
Critical (15%)											
No Action Alternative	57,112	3,419	0	337,059	0	1,814	0	0	0	0	399,403
Alternative 5	71,310	2,848	0	345,881	0	1,972	0	0	0	0	422,011
Difference	14,198	-571	0	8,822	0	158	0	0	0	0	22,608
Percent Difference	25	-17	0	3	0	9	0	0	0	0	6

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-3-16. Annual Potential Production for Spring-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	410,722
No Action Alternative	402,980
Difference	-7,742
Percent Difference ³	-2
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	449,832
No Action Alternative	442,676
Difference	-7,156
Percent Difference	-2
Above Normal (12.5%)	
Second Basis of Comparison	367,591
No Action Alternative	362,537
Difference	-5,054
Percent Difference	-1
Below Normal (17.5%)	
Second Basis of Comparison	426,491
No Action Alternative	428,569
Difference	2,078
Percent Difference	0
Dry (22.5%)	
Second Basis of Comparison	403,012
No Action Alternative	405,967
Difference	2,955
Percent Difference	1
Critical (15%)	
Second Basis of Comparison	355,097
No Action Alternative	316,344
Difference	-38,753
Percent Difference	-11
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-3-17. Annual Mortality by Life Stage for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	149,155	2,453	0	0	0
No Action Alternative	169,230	2,282	0	0	0
Difference	20,075	-171	0	0	0
Percent Difference ³	13	-7	0	0	0
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	38,874	2,303	0	0	0
No Action Alternative	54,929	2,217	0	0	0
Difference	16,055	-86	0	0	0
Percent Difference	41	-4	0	0	0
Above Normal (12.5%)					
Second Basis of Comparison	256,999	2,360	0	0	0
No Action Alternative	275,059	1,955	0	0	0
Difference	18,059	-406	0	0	0
Percent Difference	7	-17	0	0	0
Below Normal (17.5%)					
Second Basis of Comparison	110,617	2,763	0	0	0
No Action Alternative	108,811	2,619	0	0	0
Difference	-1,806	-144	0	0	0
Percent Difference	-2	-5	0	0	0
Dry (22.5%)					
Second Basis of Comparison	175,971	2,682	0	0	0
No Action Alternative	170,290	2,608	0	0	0
Difference	-5,681	-73	0	0	0
Percent Difference	-3	-3	0	0	0
Critical (15%)					
Second Basis of Comparison	302,962	2,151	0	0	0
No Action Alternative	397,589	1,814	0	0	0
Difference	94,627	-337	0	0	0
Percent Difference	31	-16	0	0	0

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-3-18. Annual Mortality by Cause for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	146,922	4,686	151,608
No Action Alternative	167,192	4,321	171,512
Difference	20,270	-366	19,904
Percent Difference ³	14	-8	13
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	36,709	4,468	41,178
No Action Alternative	53,038	4,108	57,146
Difference	16,329	-360	15,969
Percent Difference	44	-8	39
Above Normal (12.5%)			
Second Basis of Comparison	256,534	2,826	259,360
No Action Alternative	274,408	2,606	277,013
Difference	17,874	-221	17,653
Percent Difference	7	-8	7
Below Normal (17.5%)			
Second Basis of Comparison	108,800	4,580	113,380
No Action Alternative	107,177	4,253	111,431
Difference	-1,623	-327	-1,949
Percent Difference	-1	-7	-2
Dry (22.5%)			
Second Basis of Comparison	173,420	5,232	178,652
No Action Alternative	167,873	5,025	172,898
Difference	-5,547	-207	-5,754
Percent Difference	-3	-4	-3
Critical (15%)			
Second Basis of Comparison	299,101	6,012	305,113
No Action Alternative	394,171	5,232	399,403
Difference	95,070	-780	94,290
Percent Difference	32	-13	31

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-19. Annual Mortality by Cause and Life Stage for Spring-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	38,621	2,233	108,301	0	2,453	0	0	151,608
No Action Alternative	47,267	2,039	119,924	1	2,282	0	0	171,512
Difference	8,646	-194	11,623	1	-172	0	0	19,904
Percent Difference ³	22	-9	11	0	-7	0	0	13
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	260	2,165	36,450	0	2,303	0	0	41,178
No Action Alternative	340	1,893	52,697	2	2,215	0	0	57,146
Difference	80	-272	16,247	2	-88	0	0	15,969
Percent Difference	31	-13	45	0	-4	0	0	39
Above Normal (12.5%)								
Second Basis of Comparison	99,868	466	156,666	0	2,360	0	0	259,360
No Action Alternative	151,449	651	122,959	0	1,955	0	0	277,013
Difference	51,581	185	-33,707	0	-406	0	0	17,653
Percent Difference	52	40	-22	0	-17	0	0	7
Below Normal (17.5%)								
Second Basis of Comparison	66,585	1,818	42,215	0	2,763	0	0	113,380
No Action Alternative	63,840	1,634	43,337	0	2,619	0	0	111,431
Difference	-2,744	-183	1,122	0	-144	0	0	-1,949
Percent Difference	-4	-10	3	0	-5	0	0	-2
Dry (22.5%)								
Second Basis of Comparison	34,417	2,551	139,003	0	2,682	0	0	178,652
No Action Alternative	37,718	2,417	130,155	0	2,608	0	0	172,898
Difference	3,301	-134	-8,847	0	-73	0	0	-5,754
Percent Difference	10	-5	-6	0	-3	0	0	-3
Critical (15%)								
Second Basis of Comparison	44,378	3,862	254,723	0	2,151	0	0	305,113
No Action Alternative	57,112	3,419	337,059	0	1,814	0	0	399,403
Difference	12,734	-443	82,336	0	-337	0	0	94,290
Percent Difference	29	-11	32	0	-16	0	0	31
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the Annual average ⁴ Mortality values do not include base mortality								

Table B-3-20. Annual Mortality by All Factors for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	38,621	2,233	0	108,301	0	2,453	0	0	0	0	151,608
No Action Alternative	47,267	2,039	0	119,924	1	2,282	0	0	0	0	171,512
Difference	8,646	-194	0	11,623	1	-172	0	0	0	0	19,904
Percent Difference ³	22	-9	0	11	0	-7	0	0	0	0	13
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	260	2,165	0	36,450	0	2,303	0	0	0	0	41,178
No Action Alternative	340	1,893	0	52,697	2	2,215	0	0	0	0	57,146
Difference	80	-272	0	16,247	2	-88	0	0	0	0	15,969
Percent Difference	31	-13	0	45	0	-4	0	0	0	0	39
Above Normal (12.5%)											
Second Basis of Comparison	99,868	466	0	156,666	0	2,360	0	0	0	0	259,360
No Action Alternative	151,449	651	0	122,959	0	1,955	0	0	0	0	277,013
Difference	51,581	185	0	-33,707	0	-406	0	0	0	0	17,653
Percent Difference	52	40	0	-22	0	-17	0	0	0	0	7
Below Normal (17.5%)											
Second Basis of Comparison	66,585	1,818	0	42,215	0	2,763	0	0	0	0	113,380
No Action Alternative	63,840	1,634	0	43,337	0	2,619	0	0	0	0	111,431
Difference	-2,744	-183	0	1,122	0	-144	0	0	0	0	-1,949
Percent Difference	-4	-10	0	3	0	-5	0	0	0	0	-2
Dry (22.5%)											
Second Basis of Comparison	34,417	2,551	0	139,003	0	2,682	0	0	0	0	178,652
No Action Alternative	37,718	2,417	0	130,155	0	2,608	0	0	0	0	172,898
Difference	3,301	-134	0	-8,847	0	-73	0	0	0	0	-5,754
Percent Difference	10	-5	0	-6	0	-3	0	0	0	0	-3
Critical (15%)											
Second Basis of Comparison	44,378	3,862	0	254,723	0	2,151	0	0	0	0	305,113
No Action Alternative	57,112	3,419	0	337,059	0	1,814	0	0	0	0	399,403
Difference	12,734	-443	0	82,336	0	-337	0	0	0	0	94,290
Percent Difference	29	-11	0	32	0	-16	0	0	0	0	31

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-21. Annual Potential Production for Spring-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	410,722
Alternative 3	409,813
Difference	-909
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	449,832
Alternative 3	453,743
Difference	3,911
Percent Difference	1
Above Normal (12.5%)	
Second Basis of Comparison	367,591
Alternative 3	368,403
Difference	812
Percent Difference	0
Below Normal (17.5%)	
Second Basis of Comparison	426,491
Alternative 3	427,631
Difference	1,140
Percent Difference	0
Dry (22.5%)	
Second Basis of Comparison	403,012
Alternative 3	410,542
Difference	7,530
Percent Difference	2
Critical (15%)	
Second Basis of Comparison	355,097
Alternative 3	327,260
Difference	-27,838
Percent Difference	-8
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-3-22. Annual Mortality by Life Stage for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	149,155	2,453	0	0	0
Alternative 3	150,290	2,435	0	0	0
Difference	1,135	-18	0	0	0
Percent Difference ³	1	-1	0	0	0
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	38,874	2,303	0	0	0
Alternative 3	29,787	2,271	0	0	0
Difference	-9,087	-33	0	0	0
Percent Difference	-23	-1	0	0	0
Above Normal (12.5%)					
Second Basis of Comparison	256,999	2,360	0	0	0
Alternative 3	257,573	2,190	0	0	0
Difference	574	-170	0	0	0
Percent Difference	0	-7	0	0	0
Below Normal (17.5%)					
Second Basis of Comparison	110,617	2,763	0	0	0
Alternative 3	107,671	2,858	0	0	0
Difference	-2,946	95	0	0	0
Percent Difference	-3	3	0	0	0
Dry (22.5%)					
Second Basis of Comparison	175,971	2,682	0	0	0
Alternative 3	156,331	2,731	0	0	0
Difference	-19,640	50	0	0	0
Percent Difference	-11	2	0	0	0
Critical (15%)					
Second Basis of Comparison	302,962	2,151	0	0	0
Alternative 3	362,639	2,060	0	0	0
Difference	59,677	-90	0	0	0
Percent Difference	20	-4	0	0	0

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-3-23. Annual Mortality by Cause for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	146,922	4,686	151,608
Alternative 3	148,223	4,502	152,726
Difference	1,302	-184	1,118
Percent Difference ³	1	-4	1
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	36,709	4,468	41,178
Alternative 3	27,591	4,467	32,057
Difference	-9,119	-1	-9,120
Percent Difference	-25	0	-22
Above Normal (12.5%)			
Second Basis of Comparison	256,534	2,826	259,360
Alternative 3	257,166	2,597	259,763
Difference	632	-229	404
Percent Difference	0	-8	0
Below Normal (17.5%)			
Second Basis of Comparison	108,800	4,580	113,380
Alternative 3	105,832	4,697	110,529
Difference	-2,968	117	-2,851
Percent Difference	-3	3	-3
Dry (22.5%)			
Second Basis of Comparison	173,420	5,232	178,652
Alternative 3	154,048	5,014	159,062
Difference	-19,372	-219	-19,590
Percent Difference	-11	-4	-11
Critical (15%)			
Second Basis of Comparison	299,101	6,012	305,113
Alternative 3	359,528	5,172	364,700
Difference	60,427	-840	59,587
Percent Difference	20	-14	20

¹ Based on the 90-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-24. Annual Mortality by Cause and Life Stage for Spring-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	38,621	2,233	108,301	0	2,453	0	0	151,608
Alternative 3	37,164	2,067	111,060	0	2,435	0	0	152,726
Difference	-1,457	-166	2,759	0	-18	0	0	1,118
Percent Difference ³	-4	-7	3	0	-1	0	0	1
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	260	2,165	36,450	0	2,303	0	0	41,178
Alternative 3	189	2,196	27,402	0	2,271	0	0	32,057
Difference	-71	31	-9,047	0	-33	0	0	-9,120
Percent Difference	-27	1	-25	0	-1	0	0	-22
Above Normal (12.5%)								
Second Basis of Comparison	99,868	466	156,666	0	2,360	0	0	259,360
Alternative 3	104,829	407	152,337	0	2,190	0	0	259,763
Difference	4,961	-59	-4,329	0	-170	0	0	404
Percent Difference	5	-13	-3	0	-7	0	0	0
Below Normal (17.5%)								
Second Basis of Comparison	66,585	1,818	42,215	0	2,763	0	0	113,380
Alternative 3	62,085	1,839	43,747	0	2,858	0	0	110,529
Difference	-4,500	22	1,532	0	95	0	0	-2,851
Percent Difference	-7	1	4	0	3	0	0	-3
Dry (22.5%)								
Second Basis of Comparison	34,417	2,551	139,003	0	2,682	0	0	178,652
Alternative 3	28,700	2,282	125,348	0	2,731	0	0	159,062
Difference	-5,717	-269	-13,654	0	50	0	0	-19,590
Percent Difference	-17	-11	-10	0	2	0	0	-11
Critical (15%)								
Second Basis of Comparison	44,378	3,862	254,723	0	2,151	0	0	305,113
Alternative 3	44,510	3,112	315,018	0	2,060	0	0	364,700
Difference	132	-750	60,295	0	-90	0	0	59,587
Percent Difference	0	-19	24	0	-4	0	0	20

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-25. Annual Mortality by All Factors for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	38,621	2,233	0	108,301	0	2,453	0	0	0	0	151,608
Alternative 3	37,164	2,067	0	111,060	0	2,435	0	0	0	0	152,726
Difference	-1,457	-166	0	2,759	0	-18	0	0	0	0	1,118
Percent Difference ³	-4	-7	0	3	0	-1	0	0	0	0	1
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	260	2,165	0	36,450	0	2,303	0	0	0	0	41,178
Alternative 3	189	2,196	0	27,402	0	2,271	0	0	0	0	32,057
Difference	-71	31	0	-9,047	0	-33	0	0	0	0	-9,120
Percent Difference	-27	1	0	-25	0	-1	0	0	0	0	-22
Above Normal (12.5%)											
Second Basis of Comparison	99,868	466	0	156,666	0	2,360	0	0	0	0	259,360
Alternative 3	104,829	407	0	152,337	0	2,190	0	0	0	0	259,763
Difference	4,961	-59	0	-4,329	0	-170	0	0	0	0	404
Percent Difference	5	-13	0	-3	0	-7	0	0	0	0	0
Below Normal (17.5%)											
Second Basis of Comparison	66,585	1,818	0	42,215	0	2,763	0	0	0	0	113,380
Alternative 3	62,085	1,839	0	43,747	0	2,858	0	0	0	0	110,529
Difference	-4,500	22	0	1,532	0	95	0	0	0	0	-2,851
Percent Difference	-7	1	0	4	0	3	0	0	0	0	-3
Dry (22.5%)											
Second Basis of Comparison	34,417	2,551	0	139,003	0	2,682	0	0	0	0	178,652
Alternative 3	28,700	2,282	0	125,348	0	2,731	0	0	0	0	159,062
Difference	-5,717	-269	0	-13,654	0	50	0	0	0	0	-19,590
Percent Difference	-17	-11	0	-10	0	2	0	0	0	0	-11
Critical (15%)											
Second Basis of Comparison	44,378	3,862	0	254,723	0	2,151	0	0	0	0	305,113
Alternative 3	44,510	3,112	0	315,018	0	2,060	0	0	0	0	364,700
Difference	132	-750	0	60,295	0	-90	0	0	0	0	59,587
Percent Difference	0	-19	0	24	0	-4	0	0	0	0	20

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-26. Annual Potential Production for Spring-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	410,722
Alternative 5	401,678
Difference	-9,044
Percent Difference ³	-2
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	449,832
Alternative 5	441,971
Difference	-7,862
Percent Difference	-2
Above Normal (12.5%)	
Second Basis of Comparison	367,591
Alternative 5	363,460
Difference	-4,131
Percent Difference	-1
Below Normal (17.5%)	
Second Basis of Comparison	426,491
Alternative 5	428,206
Difference	1,716
Percent Difference	0
Dry (22.5%)	
Second Basis of Comparison	403,012
Alternative 5	407,290
Difference	4,278
Percent Difference	1
Critical (15%)	
Second Basis of Comparison	355,097
Alternative 5	306,861
Difference	-48,237
Percent Difference	-14
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-3-27. Annual Mortality by Life Stage for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	149,155	2,453	0	0	0
Alternative 5	171,978	2,371	0	0	0
Difference	22,823	-82	0	0	0
Percent Difference ³	15	-3	0	0	0
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	38,874	2,303	0	0	0
Alternative 5	57,192	2,203	0	0	0
Difference	18,318	-100	0	0	0
Percent Difference	47	-4	0	0	0
Above Normal (12.5%)					
Second Basis of Comparison	256,999	2,360	0	0	0
Alternative 5	271,916	1,980	0	0	0
Difference	14,917	-380	0	0	0
Percent Difference	6	-16	0	0	0
Below Normal (17.5%)					
Second Basis of Comparison	110,617	2,763	0	0	0
Alternative 5	108,195	2,925	0	0	0
Difference	-2,422	163	0	0	0
Percent Difference	-2	6	0	0	0
Dry (22.5%)					
Second Basis of Comparison	175,971	2,682	0	0	0
Alternative 5	166,496	2,666	0	0	0
Difference	-9,475	-16	0	0	0
Percent Difference	-5	-1	0	0	0
Critical (15%)					
Second Basis of Comparison	302,962	2,151	0	0	0
Alternative 5	420,039	1,972	0	0	0
Difference	117,076	-179	0	0	0
Percent Difference	39	-8	0	0	0

1 Based on the 80-year simulation period
2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
3 Relative difference of the Annual average
4 Mortality values do not include base mortality
5 Eggs mortality includes pre-spawn mortality

Table B-3-28. Annual Mortality by Cause for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	146,922	4,686	151,608
Alternative 5	170,196	4,153	174,349
Difference	23,274	-533	22,742
Percent Difference ³	16	-11	15
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	36,709	4,468	41,178
Alternative 5	55,390	4,005	59,395
Difference	18,680	-463	18,217
Percent Difference	51	-10	44
Above Normal (12.5%)			
Second Basis of Comparison	256,534	2,826	259,360
Alternative 5	271,280	2,616	273,896
Difference	14,746	-210	14,536
Percent Difference	6	-7	6
Below Normal (17.5%)			
Second Basis of Comparison	108,800	4,580	113,380
Alternative 5	106,681	4,439	111,120
Difference	-2,119	-141	-2,260
Percent Difference	-2	-3	-2
Dry (22.5%)			
Second Basis of Comparison	173,420	5,232	178,652
Alternative 5	164,607	4,554	169,161
Difference	-8,813	-678	-9,491
Percent Difference	-5	-13	-5
Critical (15%)			
Second Basis of Comparison	299,101	6,012	305,113
Alternative 5	417,191	4,820	422,011
Difference	118,090	-1,192	116,898
Percent Difference	39	-20	38

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-3-29. Annual Mortality by Cause and Life Stage for Spring-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	38,621	2,233	108,301	0	2,453	0	0	151,608
Alternative 5	44,327	1,783	125,868	0	2,371	0	0	174,349
Difference	5,706	-450	17,567	0	-82	0	0	22,742
Percent Difference ³	15	-20	16	0	-3	0	0	15
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	260	2,165	36,450	0	2,303	0	0	41,178
Alternative 5	608	1,803	54,781	1	2,203	0	0	59,395
Difference	348	-362	18,331	1	-101	0	0	18,217
Percent Difference	134	-17	50	0	-4	0	0	44
Above Normal (12.5%)								
Second Basis of Comparison	99,868	466	156,666	0	2,360	0	0	259,360
Alternative 5	125,685	636	145,595	0	1,980	0	0	273,896
Difference	25,817	171	-11,071	0	-380	0	0	14,536
Percent Difference	26	37	-7	0	-16	0	0	6
Below Normal (17.5%)								
Second Basis of Comparison	66,585	1,818	42,215	0	2,763	0	0	113,380
Alternative 5	53,122	1,514	53,559	0	2,925	0	0	111,120
Difference	-13,463	-303	11,344	0	163	0	0	-2,260
Percent Difference	-20	-17	27	0	6	0	0	-2
Dry (22.5%)								
Second Basis of Comparison	34,417	2,551	139,003	0	2,682	0	0	178,652
Alternative 5	37,450	1,889	127,157	0	2,666	0	0	169,161
Difference	3,033	-662	-11,845	0	-16	0	0	-9,491
Percent Difference	9	-26	-9	0	-1	0	0	-5
Critical (15%)								
Second Basis of Comparison	44,378	3,862	254,723	0	2,151	0	0	305,113
Alternative 5	71,310	2,848	345,881	0	1,972	0	0	422,011
Difference	26,932	-1,013	91,158	0	-179	0	0	116,898
Percent Difference	61	-26	36	0	-8	0	0	38
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the Annual average ⁴ Mortality values do not include base mortality								

Table B-3-30. Annual Mortality by All Factors for Spring-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	38,621	2,233	0	108,301	0	2,453	0	0	0	0	151,608
Alternative 5	44,327	1,783	0	125,868	0	2,371	0	0	0	0	174,349
Difference	5,706	-450	0	17,567	0	-82	0	0	0	0	22,742
Percent Difference ³	15	-20	0	16	0	-3	0	0	0	0	15
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	260	2,165	0	36,450	0	2,303	0	0	0	0	41,178
Alternative 5	608	1,803	0	54,781	1	2,203	0	0	0	0	59,395
Difference	348	-362	0	18,331	1	-101	0	0	0	0	18,217
Percent Difference	134	-17	0	50	0	-4	0	0	0	0	44
Above Normal (12.5%)											
Second Basis of Comparison	99,868	466	0	156,666	0	2,360	0	0	0	0	259,360
Alternative 5	125,685	636	0	145,595	0	1,980	0	0	0	0	273,896
Difference	25,817	171	0	-11,071	0	-380	0	0	0	0	14,536
Percent Difference	26	37	0	-7	0	-16	0	0	0	0	6
Below Normal (17.5%)											
Second Basis of Comparison	66,585	1,818	0	42,215	0	2,763	0	0	0	0	113,380
Alternative 5	53,122	1,514	0	53,559	0	2,925	0	0	0	0	111,120
Difference	-13,463	-303	0	11,344	0	163	0	0	0	0	-2,260
Percent Difference	-20	-17	0	27	0	6	0	0	0	0	-2
Dry (22.5%)											
Second Basis of Comparison	34,417	2,551	0	139,003	0	2,682	0	0	0	0	178,652
Alternative 5	37,450	1,889	0	127,157	0	2,666	0	0	0	0	169,161
Difference	3,033	-662	0	-11,845	0	-16	0	0	0	0	-9,491
Percent Difference	9	-26	0	-9	0	-1	0	0	0	0	-5
Critical (15%)											
Second Basis of Comparison	44,378	3,862	0	254,723	0	2,151	0	0	0	0	305,113
Alternative 5	71,310	2,848	0	345,881	0	1,972	0	0	0	0	422,011
Difference	26,932	-1,013	0	91,158	0	-179	0	0	0	0	116,898
Percent Difference	61	-26	0	36	0	-8	0	0	0	0	38

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.


3 Relative difference of the Annual average

4 Mortality values do not include base mortality

1 **B.4. Winter-Run Chinook Salmon**


2

Figure B-4-1. Annual Potential Production for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-2. Annual Mortality for Winter-Run Chinook Salmon - Eggs




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-3. Annual Mortality for Winter-Run Chinook Salmon - Fry




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-4. Annual Mortality for Winter-Run Chinook Salmon - Pre-Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-5. Annual Mortality for Winter-Run Chinook Salmon - Immature Smolt




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-6. Annual Mortality for Winter-Run Chinook Salmon - Pre- & Immature Smolts




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-7. Annual Mortality for Winter-Run Chinook Salmon - All Lifestages




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-8. Incubation - Habitat based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-9. Super-imposition - Habitat based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-10. Fry - Habitat based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-11. Pre-smolt - Habitat based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-12. Immature Smolt - Habitat based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-13. Total Habitat based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-14. Pre-Spawn Mortality - Temperature based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-15. Eggs - Temperature based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-16. Fry - Temperature based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-17. Pre-smolt - Temperature based Annual Mortality for Winter-Run Chinook Salmon




Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-18. Immature Smolt - Temperature based Annual Mortality for Winter-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Figure B-4-19. Total Temperature based Annual Mortality for Winter-Run Chinook Salmon



Notes: 1) Exceedance probability is defined as the probability a given value will be exceeded in any one year. 2) All alternatives are simulated with projected hydrology and sea level at Year 2030 conditions. 3) Model results for Alternatives 1, 4, and Second Basis of Comparison are the same, therefore Alternatives 1 and 4 results are not presented. Qualitative differences, if applicable, are discussed in the text. 4) Model results for Alternative 2 and No Action Alternative are the same, therefore Alternative 2 results are not presented. Qualitative differences, if applicable, are discussed in the text.

Table B-4-1. Annual Potential Production for Winter-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	1,883,893
Alternative 1	1,885,400
Difference	1,507
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	1,952,705
Alternative 1	1,930,740
Difference	-21,965
Percent Difference	-1
Above Normal (12.5%)	
No Action Alternative	1,707,717
Alternative 1	1,746,928
Difference	39,211
Percent Difference	2
Below Normal (17.5%)	
No Action Alternative	1,863,415
Alternative 1	1,847,619
Difference	-15,795
Percent Difference	-1
Dry (22.5%)	
No Action Alternative	1,883,395
Alternative 1	1,894,107
Difference	10,712
Percent Difference	1
Critical (15%)	
No Action Alternative	1,906,250
Alternative 1	1,933,573
Difference	27,323
Percent Difference	1
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-4-2. Annual Mortality by Life Stage for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	222,517	196,405	26,961	138	27,099
Alternative 1	259,052	162,983	23,312	137	23,449
Difference	36,535	-33,421	-3,649	-2	-3,650
Percent Difference ³	16	-17	-14	-1	-13
Water Year Types²					
Wet (32.5%)					
No Action Alternative	90,910	197,835	1,943	54	1,997
Alternative 1	155,104	176,315	1,060	47	1,107
Difference	64,194	-21,520	-883	-7	-890
Percent Difference	71	-11	-45	-13	-45
Above Normal (12.5%)					
No Action Alternative	469,585	220,960	53,686	94	53,779
Alternative 1	438,691	167,899	63,706	103	63,808
Difference	-30,894	-53,061	10,020	9	10,029
Percent Difference	-7	-24	19	9	19
Below Normal (17.5%)					
No Action Alternative	275,022	176,292	19,822	61	19,884
Alternative 1	337,945	142,925	18,481	41	18,522
Difference	62,922	-33,367	-1,341	-21	-1,362
Percent Difference	23	-19	-7	-34	-7
Dry (22.5%)					
No Action Alternative	209,708	215,896	24,076	139	24,215
Alternative 1	240,069	172,393	22,611	143	22,755
Difference	30,361	-43,503	-1,465	4	-1,460
Percent Difference	14	-20	-6	3	-6
Critical (15%)					
No Action Alternative	259,734	167,072	71,553	447	72,000
Alternative 1	271,006	139,289	44,553	461	45,014
Difference	11,272	-27,783	-27,000	14	-26,985
Percent Difference	4	-17	-38	3	-37

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

⁵ Eggs mortality includes pre-spawn mortality

Table B-4-3. Annual Mortality by Cause for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	178,654	267,367	446,021
Alternative 1	149,945	295,539	445,484
Difference	-28,708	28,172	-537
Percent Difference ³	-16	11	0
Water Year Types²			
Wet (32.5%)			
No Action Alternative	3,522	287,219	290,741
Alternative 1	1,273	331,252	332,525
Difference	-2,249	44,034	41,785
Percent Difference	-64	15	14
Above Normal (12.5%)			
No Action Alternative	504,624	239,700	744,324
Alternative 1	388,548	281,850	670,398
Difference	-116,076	42,150	-73,926
Percent Difference	-23	18	-10
Below Normal (17.5%)			
No Action Alternative	212,903	258,295	471,198
Alternative 1	218,115	281,277	499,391
Difference	5,212	22,981	28,193
Percent Difference	2	9	6
Dry (22.5%)			
No Action Alternative	155,797	294,022	449,819
Alternative 1	134,348	300,869	435,217
Difference	-21,449	6,847	-14,602
Percent Difference	-14	2	-3
Critical (15%)			
No Action Alternative	280,793	218,012	498,805
Alternative 1	217,099	238,210	455,309
Difference	-63,694	20,198	-43,496
Percent Difference	-23	9	-9

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-4. Annual Mortality by Cause and Life Stage for Winter-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	0	93,980	128,537	24,093	172,312	26,023	1,076	446,021
Alternative 1	0	151,512	107,540	20,257	142,726	22,149	1,300	445,484
Difference	0	57,532	-20,997	-3,836	-29,585	-3,875	225	-537
Percent Difference ³	-36	61	-16	-16	-17	-15	21	0
Water Year Types²								
Wet (32.5%)								
No Action Alternative	0	88,673	2,236	182	197,652	1,103	893	290,741
Alternative 1	0	153,836	1,268	3	176,312	3	1,104	332,525
Difference	0	65,163	-969	-180	-21,340	-1,101	211	41,784
Percent Difference	0	73	-43	-98	-11	-100	24	14
Above Normal (12.5%)								
No Action Alternative	0	83,031	386,554	64,945	156,015	53,125	654	744,324
Alternative 1	0	169,913	268,778	56,974	110,925	62,797	1,012	670,398
Difference	0	86,882	-117,776	-7,972	-45,090	9,671	358	-73,926
Percent Difference	0	105	-30	-12	-29	18	55	-10
Below Normal (17.5%)								
No Action Alternative	0	101,792	173,231	20,940	155,352	18,732	1,152	471,198
Alternative 1	0	157,331	180,614	20,113	122,812	17,388	1,134	499,391
Difference	0	55,539	7,383	-827	-32,540	-1,344	-18	28,193
Percent Difference	0	55	4	-4	-21	-7	-2	6
Dry (22.5%)								
No Action Alternative	2	100,064	109,642	23,024	192,872	23,129	1,086	449,819
Alternative 1	1	148,149	91,919	21,162	151,231	21,266	1,488	435,217
Difference	0	48,085	-17,723	-1,862	-41,641	-1,863	402	-14,602
Percent Difference	-23	48	-16	-8	-22	-8	37	-3
Critical (15%)								
No Action Alternative	1	96,360	163,373	47,138	119,933	70,281	1,719	498,805
Alternative 1	0	129,397	141,609	32,354	106,935	43,136	1,878	455,309
Difference	-1	33,037	-21,764	-14,784	-12,999	-27,145	160	-43,496
Percent Difference	-100	34	-13	-31	-11	-39	9	-9

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality

Table B-4-5. Annual Mortality by All Factors for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	0	93,980	0	128,537	24,093	172,312	26,020	941	3	135	446,021
Alternative 1	0	151,512	0	107,540	20,257	142,726	22,146	1,167	3	134	445,484
Difference	0	57,532	0	-20,997	-3,836	-29,585	-3,875	226	0	-1	-537
Percent Difference ³	-36	61	0	-16	-16	-17	-15	24	-7	-1	0
Water Year Types²											
Wet (32.5%)											
No Action Alternative	0	88,673	0	2,236	182	197,652	1,101	842	3	51	290,741
Alternative 1	0	153,836	0	1,268	3	176,312	3	1,057	0	47	332,525
Difference	0	65,163	0	-969	-180	-21,340	-1,098	215	-3	-4	41,784
Percent Difference	0	73	0	-43	-98	-11	-100	26	-100	-8	14
Above Normal (12.5%)											
No Action Alternative	0	83,031	0	386,554	64,945	156,015	53,122	564	3	90	744,324
Alternative 1	0	169,913	0	268,778	56,974	110,925	62,779	926	17	85	670,398
Difference	0	86,882	0	-117,776	-7,972	-45,090	9,658	363	14	-5	-73,926
Percent Difference	0	105	0	-30	-12	-29	18	64	406	-6	-10
Below Normal (17.5%)											
No Action Alternative	0	101,792	0	173,231	20,940	155,352	18,732	1,091	0	61	471,198
Alternative 1	0	157,331	0	180,614	20,113	122,812	17,388	1,093	0	41	499,391
Difference	0	55,539	0	7,383	-827	-32,540	-1,344	3	0	-21	28,193
Percent Difference	0	55	0	4	-4	-21	-7	0	0	-34	6
Dry (22.5%)											
No Action Alternative	2	100,064	0	109,642	23,024	192,872	23,129	947	0	139	449,819
Alternative 1	1	148,149	0	91,919	21,162	151,231	21,264	1,348	3	141	435,217
Difference	0	48,085	0	-17,723	-1,862	-41,641	-1,865	401	3	2	-14,602
Percent Difference	-23	48	0	-16	-8	-22	-8	42	0	1	-3
Critical (15%)											
No Action Alternative	1	96,360	0	163,373	47,138	119,933	70,269	1,283	12	435	498,805
Alternative 1	0	129,397	0	141,609	32,354	106,935	43,135	1,418	1	460	455,309
Difference	-1	33,037	0	-21,764	-14,784	-12,999	-27,135	135	-11	25	-43,496
Percent Difference	-100	34	0	-13	-31	-11	-39	11	-90	6	-9

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-6. Annual Potential Production for Winter-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	1,883,893
Alternative 3	1,897,120
Difference	13,227
Percent Difference ³	1
Water Year Types²	
Wet (32.5%)	
No Action Alternative	1,952,705
Alternative 3	1,944,614
Difference	-8,091
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	1,707,717
Alternative 3	1,752,903
Difference	45,186
Percent Difference	3
Below Normal (17.5%)	
No Action Alternative	1,863,415
Alternative 3	1,840,343
Difference	-23,072
Percent Difference	-1
Dry (22.5%)	
No Action Alternative	1,883,395
Alternative 3	1,919,466
Difference	36,071
Percent Difference	2
Critical (15%)	
No Action Alternative	1,906,250
Alternative 3	1,947,116
Difference	40,866
Percent Difference	2
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-4-7. Annual Mortality by Life Stage for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	222,517	196,405	26,961	138	27,099
Alternative 3	237,813	165,266	21,803	140	21,943
Difference	15,296	-31,139	-5,158	2	-5,156
Percent Difference ³	7	-16	-19	1	-19
Water Year Types²					
Wet (32.5%)					
No Action Alternative	90,910	197,835	1,943	54	1,997
Alternative 3	131,631	174,265	1,188	34	1,222
Difference	40,721	-23,569	-755	-20	-774
Percent Difference	45	-12	-39	-37	-39
Above Normal (12.5%)					
No Action Alternative	469,585	220,960	53,686	94	53,779
Alternative 3	443,487	166,295	54,841	70	54,912
Difference	-26,098	-54,664	1,156	-23	1,133
Percent Difference	-6	-25	2	-25	2
Below Normal (17.5%)					
No Action Alternative	275,022	176,292	19,822	61	19,884
Alternative 3	324,721	159,309	20,994	55	21,049
Difference	49,699	-16,983	1,172	-6	1,166
Percent Difference	18	-10	6	-10	6
Dry (22.5%)					
No Action Alternative	209,708	215,896	24,076	139	24,215
Alternative 3	207,993	170,244	16,866	166	17,032
Difference	-1,715	-45,653	-7,210	27	-7,183
Percent Difference	-1	-21	-30	19	-30
Critical (15%)					
No Action Alternative	259,734	167,072	71,553	447	72,000
Alternative 3	239,816	144,393	47,286	490	47,776
Difference	-19,918	-22,679	-24,267	43	-24,224
Percent Difference	-8	-14	-34	10	-34

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-4-8. Annual Mortality by Cause for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	178,654	267,367	446,021
Alternative 3	142,827	282,195	425,022
Difference	-35,827	14,828	-20,999
Percent Difference ³	-20	6	-5
Water Year Types²			
Wet (32.5%)			
No Action Alternative	3,522	287,219	290,741
Alternative 3	1,126	305,992	307,118
Difference	-2,396	18,773	16,377
Percent Difference	-68	7	6
Above Normal (12.5%)			
No Action Alternative	504,624	239,700	744,324
Alternative 3	430,489	234,205	664,694
Difference	-74,135	-5,495	-79,630
Percent Difference	-15	-2	-11
Below Normal (17.5%)			
No Action Alternative	212,903	258,295	471,198
Alternative 3	210,138	294,942	505,080
Difference	-2,765	36,647	33,882
Percent Difference	-1	14	7
Dry (22.5%)			
No Action Alternative	155,797	294,022	449,819
Alternative 3	95,635	299,633	395,268
Difference	-60,162	5,611	-54,551
Percent Difference	-39	2	-12
Critical (15%)			
No Action Alternative	280,793	218,012	498,805
Alternative 3	202,386	229,599	431,984
Difference	-78,407	11,587	-66,821
Percent Difference	-28	5	-13

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-9. Annual Mortality by Cause and Life Stage for Winter-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	0	93,980	128,537	24,093	172,312	26,023	1,076	446,021
Alternative 3	0	135,049	102,763	19,523	145,743	20,541	1,402	425,022
Difference	0	41,070	-25,774	-4,571	-26,568	-5,482	326	-20,999
Percent Difference ³	-100	44	-20	-19	-15	-21	30	-5
Water Year Types²								
Wet (32.5%)								
No Action Alternative	0	88,673	2,236	182	197,652	1,103	893	290,741
Alternative 3	0	130,505	1,126	1	174,265	0	1,222	307,118
Difference	0	41,832	-1,111	-181	-23,388	-1,103	329	16,377
Percent Difference	0	47	-50	-100	-12	-100	37	6
Above Normal (12.5%)								
No Action Alternative	0	83,031	386,554	64,945	156,015	53,125	654	744,324
Alternative 3	0	119,969	323,517	52,929	113,366	54,043	869	664,694
Difference	0	36,938	-63,037	-12,016	-42,648	917	215	-79,630
Percent Difference	0	44	-16	-19	-27	2	33	-11
Below Normal (17.5%)								
No Action Alternative	0	101,792	173,231	20,940	155,352	18,732	1,152	471,198
Alternative 3	0	155,899	168,822	21,483	137,826	19,833	1,217	505,080
Difference	0	54,108	-4,409	542	-17,525	1,101	65	33,882
Percent Difference	0	53	-3	3	-11	6	6	7
Dry (22.5%)								
No Action Alternative	2	100,064	109,642	23,024	192,872	23,129	1,086	449,819
Alternative 3	0	146,046	61,947	18,345	151,898	15,343	1,689	395,268
Difference	-2	45,982	-47,695	-4,679	-40,974	-7,786	603	-54,551
Percent Difference	-100	46	-44	-20	-21	-34	55	-12
Critical (15%)								
No Action Alternative	1	96,360	163,373	47,138	119,933	70,281	1,719	498,805
Alternative 3	0	116,643	123,172	33,460	110,932	45,753	2,023	431,984
Difference	-1	20,283	-40,201	-13,678	-9,001	-24,528	305	-66,821
Percent Difference	-100	21	-25	-29	-8	-35	18	-13

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-10. Annual Mortality by All Factors for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
No Action Alternative	0	93,980	0	128,537	24,093	172,312	26,020	941	3	135	446,021
Alternative 3	0	135,049	0	102,763	19,523	145,743	20,536	1,267	5	135	425,022
Difference	0	41,070	0	-25,774	-4,571	-26,568	-5,484	326	2	0	-20,999
Percent Difference ³	-100	44	0	-20	-19	-15	-21	35	60	0	-5
Water Year Types²											
Wet (32.5%)											
No Action Alternative	0	88,673	0	2,236	182	197,652	1,101	842	3	51	290,741
Alternative 3	0	130,505	0	1,126	1	174,265	0	1,188	0	34	307,118
Difference	0	41,832	0	-1,111	-181	-23,388	-1,101	346	-3	-17	16,377
Percent Difference	0	47	0	-50	-100	-12	-100	41	-100	-33	6
Above Normal (12.5%)											
No Action Alternative	0	83,031	0	386,554	64,945	156,015	53,122	564	3	90	744,324
Alternative 3	0	119,969	0	323,517	52,929	113,366	54,043	799	0	70	664,694
Difference	0	36,938	0	-63,037	-12,016	-42,648	921	235	-3	-20	-79,630
Percent Difference	0	44	0	-16	-19	-27	2	42	-100	-22	-11
Below Normal (17.5%)											
No Action Alternative	0	101,792	0	173,231	20,940	155,352	18,732	1,091	0	61	471,198
Alternative 3	0	155,899	0	168,822	21,483	137,826	19,832	1,162	1	54	505,080
Difference	0	54,108	0	-4,409	542	-17,525	1,100	72	1	-7	33,882
Percent Difference	0	53	0	-3	3	-11	6	7	0	-11	7
Dry (22.5%)											
No Action Alternative	2	100,064	0	109,642	23,024	192,872	23,129	947	0	139	449,819
Alternative 3	0	146,046	0	61,947	18,345	151,898	15,343	1,523	0	166	395,268
Difference	-2	45,982	0	-47,695	-4,679	-40,974	-7,786	576	0	27	-54,551
Percent Difference	-100	46	0	-44	-20	-21	-34	61	0	19	-12
Critical (15%)											
No Action Alternative	1	96,360	0	163,373	47,138	119,933	70,269	1,283	12	435	498,805
Alternative 3	0	116,643	0	123,172	33,460	110,932	45,720	1,566	33	457	431,984
Difference	-1	20,283	0	-40,201	-13,678	-9,001	-24,549	283	21	22	-66,821
Percent Difference	-100	21	0	-25	-29	-8	-35	22	180	5	-13

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-4-11. Annual Potential Production for Winter-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
No Action Alternative	1,883,893
Alternative 5	1,883,178
Difference	-715
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
No Action Alternative	1,952,705
Alternative 5	1,943,241
Difference	-9,464
Percent Difference	0
Above Normal (12.5%)	
No Action Alternative	1,707,717
Alternative 5	1,698,809
Difference	-8,908
Percent Difference	-1
Below Normal (17.5%)	
No Action Alternative	1,863,415
Alternative 5	1,898,667
Difference	35,252
Percent Difference	2
Dry (22.5%)	
No Action Alternative	1,883,395
Alternative 5	1,876,977
Difference	-6,419
Percent Difference	0
Critical (15%)	
No Action Alternative	1,906,250
Alternative 5	1,897,912
Difference	-8,338
Percent Difference	0
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-4-12. Annual Mortality by Life Stage for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
No Action Alternative	222,517	196,405	26,961	138	27,099
Alternative 5	203,248	207,870	29,865	124	29,989
Difference	-19,269	11,465	2,904	-14	2,890
Percent Difference ³	-9	6	11	-10	11
Water Year Types²					
Wet (32.5%)					
No Action Alternative	90,910	197,835	1,943	54	1,997
Alternative 5	87,970	210,570	4,085	28	4,113
Difference	-2,939	12,735	2,142	-26	2,117
Percent Difference	-3	6	110	-48	106
Above Normal (12.5%)					
No Action Alternative	469,585	220,960	53,686	94	53,779
Alternative 5	464,585	236,533	52,336	89	52,425
Difference	-5,000	15,573	-1,349	-5	-1,354
Percent Difference	-1	7	-3	-5	-3
Below Normal (17.5%)					
No Action Alternative	275,022	176,292	19,822	61	19,884
Alternative 5	191,541	178,323	31,052	108	31,160
Difference	-83,481	2,031	11,229	47	11,276
Percent Difference	-30	1	57	76	57
Dry (22.5%)					
No Action Alternative	209,708	215,896	24,076	139	24,215
Alternative 5	200,255	234,855	20,690	134	20,824
Difference	-9,453	18,959	-3,386	-5	-3,391
Percent Difference	-5	9	-14	-3	-14
Critical (15%)					
No Action Alternative	259,734	167,072	71,553	447	72,000
Alternative 5	253,379	172,126	79,375	365	79,740
Difference	-6,354	5,055	7,822	-82	7,740
Percent Difference	-2	3	11	-18	11

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-4-13. Annual Mortality by Cause for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
No Action Alternative	178,654	267,367	446,021
Alternative 5	170,139	270,968	441,107
Difference	-8,515	3,601	-4,914
Percent Difference ³	-5	1	-1
Water Year Types²			
Wet (32.5%)			
No Action Alternative	3,522	287,219	290,741
Alternative 5	7,569	295,085	302,654
Difference	4,047	7,866	11,913
Percent Difference	115	3	4
Above Normal (12.5%)			
No Action Alternative	504,624	239,700	744,324
Alternative 5	499,928	253,615	753,543
Difference	-4,696	13,915	9,219
Percent Difference	-1	6	1
Below Normal (17.5%)			
No Action Alternative	212,903	258,295	471,198
Alternative 5	149,215	251,809	401,024
Difference	-63,688	-6,486	-70,174
Percent Difference	-30	-3	-15
Dry (22.5%)			
No Action Alternative	155,797	294,022	449,819
Alternative 5	146,764	309,170	455,934
Difference	-9,033	15,148	6,115
Percent Difference	-6	5	1
Critical (15%)			
No Action Alternative	280,793	218,012	498,805
Alternative 5	307,023	198,222	505,246
Difference	26,230	-19,790	6,441
Percent Difference	9	-9	1

¹ Based on the 80-year simulation period not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-14. Annual Mortality by Cause and Life Stage for Winter-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
No Action Alternative	0	93,980	128,537	24,093	172,312	26,023	1,076	446,021
Alternative 5	0	89,100	114,147	27,082	180,788	28,909	1,080	441,107
Difference	0	-4,880	-14,389	2,989	8,476	2,886	5	-4,914
Percent Difference ³	0	-5	-11	12	5	11	0	-1
Water Year Types²								
Wet (32.5%)								
No Action Alternative	0	88,673	2,236	182	197,652	1,103	893	290,741
Alternative 5	0	84,683	3,288	977	209,593	3,304	809	302,654
Difference	0	-3,991	1,051	795	11,941	2,201	-84	11,913
Percent Difference	0	-5	47	436	6	199	-9	4
Above Normal (12.5%)								
No Action Alternative	0	83,031	386,554	64,945	156,015	53,125	654	744,324
Alternative 5	0	80,569	384,016	64,143	172,390	51,769	656	753,543
Difference	0	-2,463	-2,538	-802	16,375	-1,356	2	9,219
Percent Difference	0	-3	-1	-1	10	-3	0	1
Below Normal (17.5%)								
No Action Alternative	0	101,792	173,231	20,940	155,352	18,732	1,152	471,198
Alternative 5	0	103,637	87,904	31,368	146,956	29,943	1,216	401,024
Difference	0	1,845	-85,326	10,427	-8,396	11,212	64	-70,174
Percent Difference	0	2	-49	50	-5	60	6	-15
Dry (22.5%)								
No Action Alternative	2	100,064	109,642	23,024	192,872	23,129	1,086	449,819
Alternative 5	2	94,247	106,007	21,110	213,744	19,645	1,179	455,934
Difference	0	-5,817	-3,635	-1,914	20,873	-3,484	93	6,115
Percent Difference	0	-6	-3	-8	11	-15	9	1
Critical (15%)								
No Action Alternative	1	96,360	163,373	47,138	119,933	70,281	1,719	498,805
Alternative 5	1	81,098	172,281	56,716	115,410	78,025	1,715	505,246
Difference	0	-15,262	8,908	9,578	-4,524	7,744	-4	6,441
Percent Difference	0	-16	5	20	-4	11	0	1

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-15. Annual Mortality by All Factors for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
No Action Alternative	0	93,980	0	128,537	24,093	172,312	26,020	941	3	135	446,021
Alternative 5	0	89,100	0	114,147	27,082	180,788	28,902	963	7	117	441,107
Difference	0	-4,880	0	-14,389	2,989	8,476	2,882	22	4	-18	-4,914
Percent Difference ³	0	-5	0	-11	12	5	11	2	118	-13	-1
Water Year Types²											
Wet (32.5%)											
No Action Alternative	0	88,673	0	2,236	182	197,652	1,101	842	3	51	290,741
Alternative 5	0	84,683	0	3,288	977	209,593	3,302	784	3	26	302,654
Difference	0	-3,991	0	1,051	795	11,941	2,201	-59	0	-25	11,913
Percent Difference	0	-5	0	47	436	6	200	-7	-8	-50	4
Above Normal (12.5%)											
No Action Alternative	0	83,031	0	386,554	64,945	156,015	53,122	564	3	90	744,324
Alternative 5	0	80,569	0	384,016	64,143	172,390	51,732	604	37	52	753,543
Difference	0	-2,463	0	-2,538	-802	16,375	-1,389	40	33	-38	9,219
Percent Difference	0	-3	0	-1	-1	10	-3	7	976	-42	1
Below Normal (17.5%)											
No Action Alternative	0	101,792	0	173,231	20,940	155,352	18,732	1,091	0	61	471,198
Alternative 5	0	103,637	0	87,904	31,368	146,956	29,943	1,108	0	108	401,024
Difference	0	1,845	0	-85,326	10,427	-8,396	11,212	18	0	47	-70,174
Percent Difference	0	2	0	-49	50	-5	60	2	0	76	-15
Dry (22.5%)											
No Action Alternative	2	100,064	0	109,642	23,024	192,872	23,129	947	0	139	449,819
Alternative 5	2	94,247	0	106,007	21,110	213,744	19,645	1,045	0	134	455,934
Difference	0	-5,817	0	-3,635	-1,914	20,873	-3,484	98	0	-5	6,115
Percent Difference	0	-6	0	-3	-8	11	-15	10	0	-3	1
Critical (15%)											
No Action Alternative	1	96,360	0	163,373	47,138	119,933	70,269	1,283	12	435	498,805
Alternative 5	1	81,098	0	172,281	56,716	115,410	78,016	1,359	9	356	505,246
Difference	0	-15,262	0	8,908	9,578	-4,524	7,747	75	-3	-79	6,441
Percent Difference	0	-16	0	5	20	-4	11	6	-22	-18	1

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-16. Annual Potential Production for Winter-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	1,885,400
No Action Alternative	1,883,893
Difference	-1,507
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	1,930,740
No Action Alternative	1,952,705
Difference	21,965
Percent Difference	1
Above Normal (12.5%)	
Second Basis of Comparison	1,746,928
No Action Alternative	1,707,717
Difference	-39,211
Percent Difference	-2
Below Normal (17.5%)	
Second Basis of Comparison	1,847,619
No Action Alternative	1,863,415
Difference	15,795
Percent Difference	1
Dry (22.5%)	
Second Basis of Comparison	1,894,107
No Action Alternative	1,883,395
Difference	-10,712
Percent Difference	-1
Critical (15%)	
Second Basis of Comparison	1,933,573
No Action Alternative	1,906,250
Difference	-27,323
Percent Difference	-1
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-4-17. Annual Mortality by Life Stage for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	259,052	162,983	23,312	137	23,449
No Action Alternative	222,517	196,405	26,961	138	27,099
Difference	-36,535	33,421	3,649	2	3,650
Percent Difference ³	-14	21	16	1	16
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	155,104	176,315	1,060	47	1,107
No Action Alternative	90,910	197,835	1,943	54	1,997
Difference	-64,194	21,520	883	7	890
Percent Difference	-41	12	83	15	80
Above Normal (12.5%)					
Second Basis of Comparison	438,691	167,899	63,706	103	63,808
No Action Alternative	469,585	220,960	53,686	94	53,779
Difference	30,894	53,061	-10,020	-9	-10,029
Percent Difference	7	32	-16	-8	-16
Below Normal (17.5%)					
Second Basis of Comparison	337,945	142,925	18,481	41	18,522
No Action Alternative	275,022	176,292	19,822	61	19,884
Difference	-62,922	33,367	1,341	21	1,362
Percent Difference	-19	23	7	50	7
Dry (22.5%)					
Second Basis of Comparison	240,069	172,393	22,611	143	22,755
No Action Alternative	209,708	215,896	24,076	139	24,215
Difference	-30,361	43,503	1,465	-4	1,460
Percent Difference	-13	25	6	-3	6
Critical (15%)					
Second Basis of Comparison	271,006	139,289	44,553	461	45,014
No Action Alternative	259,734	167,072	71,553	447	72,000
Difference	-11,272	27,783	27,000	-14	26,985
Percent Difference	-4	20	61	-3	60

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-4-18. Annual Mortality by Cause for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	149,945	295,539	445,484
No Action Alternative	178,654	267,367	446,021
Difference	28,708	-28,172	537
Percent Difference ³	19	-10	0
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	1,273	331,252	332,525
No Action Alternative	3,522	287,219	290,741
Difference	2,249	-44,034	-41,785
Percent Difference	177	-13	-13
Above Normal (12.5%)			
Second Basis of Comparison	388,548	281,850	670,398
No Action Alternative	504,624	239,700	744,324
Difference	116,076	-42,150	73,926
Percent Difference	30	-15	11
Below Normal (17.5%)			
Second Basis of Comparison	218,115	281,277	499,391
No Action Alternative	212,903	258,295	471,198
Difference	-5,212	-22,981	-28,193
Percent Difference	-2	-8	-6
Dry (22.5%)			
Second Basis of Comparison	134,348	300,869	435,217
No Action Alternative	155,797	294,022	449,819
Difference	21,449	-6,847	14,602
Percent Difference	16	-2	3
Critical (15%)			
Second Basis of Comparison	217,099	238,210	455,309
No Action Alternative	280,793	218,012	498,805
Difference	63,694	-20,198	43,496
Percent Difference	29	-8	10

¹ Based on the 90 year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-19. Annual Mortality by Cause and Life Stage for Winter-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	0	151,512	107,540	20,257	142,726	22,149	1,300	445,484
No Action Alternative	0	93,980	128,537	24,093	172,312	26,023	1,076	446,021
Difference	0	-57,532	20,997	3,836	29,585	3,875	-225	537
Percent Difference ³	57	-38	20	19	21	17	-17	0
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	0	153,836	1,268	3	176,312	3	1,104	332,525
No Action Alternative	0	88,673	2,236	182	197,652	1,103	893	290,741
Difference	0	-65,163	969	180	21,340	1,101	-211	-41,784
Percent Difference	0	-42	76	6,482	12	44,038	-19	-13
Above Normal (12.5%)								
Second Basis of Comparison	0	169,913	268,778	56,974	110,925	62,797	1,012	670,398
No Action Alternative	0	83,031	386,554	64,945	156,015	53,125	654	744,324
Difference	0	-86,882	117,776	7,972	45,090	-9,671	-358	73,926
Percent Difference	0	-51	44	14	41	-15	-35	11
Below Normal (17.5%)								
Second Basis of Comparison	0	157,331	180,614	20,113	122,812	17,388	1,134	499,391
No Action Alternative	0	101,792	173,231	20,940	155,352	18,732	1,152	471,198
Difference	0	-55,539	-7,383	827	32,540	1,344	18	-28,193
Percent Difference	0	-35	-4	4	26	8	2	-6
Dry (22.5%)								
Second Basis of Comparison	1	148,149	91,919	21,162	151,231	21,266	1,488	435,217
No Action Alternative	2	100,064	109,642	23,024	192,872	23,129	1,086	449,819
Difference	0	-48,085	17,723	1,862	41,641	1,863	-402	14,602
Percent Difference	30	-32	19	9	28	9	-27	3
Critical (15%)								
Second Basis of Comparison	0	129,397	141,609	32,354	106,935	43,136	1,878	455,309
No Action Alternative	1	96,360	163,373	47,138	119,933	70,281	1,719	498,805
Difference	1	-33,037	21,764	14,784	12,999	27,145	-160	43,496
Percent Difference	0	-26	15	46	12	63	-9	10
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the Annual average ⁴ Mortality values do not include base mortality								

Table B-4-20. Annual Mortality by All Factors for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	0	151,512	0	107,540	20,257	142,726	22,146	1,167	3	134	445,484
No Action Alternative	0	93,980	0	128,537	24,093	172,312	26,020	941	3	135	446,021
Difference	0	-57,532	0	20,997	3,836	29,585	3,875	-226	0	1	537
Percent Difference ³	57	-38	0	20	19	21	17	-19	8	1	0
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	0	153,836	0	1,268	3	176,312	3	1,057	0	47	332,525
No Action Alternative	0	88,673	0	2,236	182	197,652	1,101	842	3	51	290,741
Difference	0	-65,163	0	969	180	21,340	1,098	-215	3	4	-41,784
Percent Difference	0	-42	0	76	6,482	12	43,923	-20	0	9	-13
Above Normal (12.5%)											
Second Basis of Comparison	0	169,913	0	268,778	56,974	110,925	62,779	926	17	85	670,398
No Action Alternative	0	83,031	0	386,554	64,945	156,015	53,122	564	3	90	744,324
Difference	0	-86,882	0	117,776	7,972	45,090	-9,658	-363	-14	5	73,926
Percent Difference	0	-51	0	44	14	41	-15	-39	-80	6	11
Below Normal (17.5%)											
Second Basis of Comparison	0	157,331	0	180,614	20,113	122,812	17,388	1,093	0	41	499,391
No Action Alternative	0	101,792	0	173,231	20,940	155,352	18,732	1,091	0	61	471,198
Difference	0	-55,539	0	-7,383	827	32,540	1,344	-3	0	21	-28,193
Percent Difference	0	-35	0	-4	4	26	8	0	0	50	-6
Dry (22.5%)											
Second Basis of Comparison	1	148,149	0	91,919	21,162	151,231	21,264	1,348	3	141	435,217
No Action Alternative	2	100,064	0	109,642	23,024	192,872	23,129	947	0	139	449,819
Difference	0	-48,085	0	17,723	1,862	41,641	1,865	-401	-3	-2	14,602
Percent Difference	30	-32	0	19	9	28	9	-30	-100	-1	3
Critical (15%)											
Second Basis of Comparison	0	129,397	0	141,609	32,354	106,935	43,135	1,418	1	460	455,309
No Action Alternative	1	96,360	0	163,373	47,138	119,933	70,269	1,283	12	435	498,805
Difference	1	-33,037	0	21,764	14,784	12,999	27,135	-135	11	-25	43,496
Percent Difference	0	-26	0	15	46	12	63	-10	900	-5	10

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-21. Annual Potential Production for Winter-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	1,885,400
Alternative 3	1,897,120
Difference	11,720
Percent Difference ³	1
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	1,930,740
Alternative 3	1,944,614
Difference	13,874
Percent Difference	1
Above Normal (12.5%)	
Second Basis of Comparison	1,746,928
Alternative 3	1,752,903
Difference	5,975
Percent Difference	0
Below Normal (17.5%)	
Second Basis of Comparison	1,847,619
Alternative 3	1,840,343
Difference	-7,277
Percent Difference	0
Dry (22.5%)	
Second Basis of Comparison	1,894,107
Alternative 3	1,919,466
Difference	25,359
Percent Difference	1
Critical (15%)	
Second Basis of Comparison	1,933,573
Alternative 3	1,947,116
Difference	13,543
Percent Difference	1
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-4-22. Annual Mortality by Life Stage for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	259,052	162,983	23,312	137	23,449
Alternative 3	237,813	165,266	21,803	140	21,943
Difference	-21,239	2,283	-1,509	4	-1,506
Percent Difference ³	-8	1	-6	3	-6
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	155,104	176,315	1,060	47	1,107
Alternative 3	131,631	174,265	1,188	34	1,222
Difference	-23,473	-2,050	128	-13	116
Percent Difference	-15	-1	12	-28	10
Above Normal (12.5%)					
Second Basis of Comparison	438,691	167,899	63,706	103	63,808
Alternative 3	443,487	166,295	54,841	70	54,912
Difference	4,795	-1,603	-8,864	-32	-8,897
Percent Difference	1	-1	-14	-31	-14
Below Normal (17.5%)					
Second Basis of Comparison	337,945	142,925	18,481	41	18,522
Alternative 3	324,721	159,309	20,994	55	21,049
Difference	-13,223	16,384	2,513	14	2,527
Percent Difference	-4	11	14	35	14
Dry (22.5%)					
Second Basis of Comparison	240,069	172,393	22,611	143	22,755
Alternative 3	207,993	170,244	16,866	166	17,032
Difference	-32,076	-2,150	-5,745	22	-5,723
Percent Difference	-13	-1	-25	16	-25
Critical (15%)					
Second Basis of Comparison	271,006	139,289	44,553	461	45,014
Alternative 3	239,816	144,393	47,286	490	47,776
Difference	-31,190	5,104	2,733	29	2,762
Percent Difference	-12	4	6	6	6

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-4-23. Annual Mortality by Cause for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	149,945	295,539	445,484
Alternative 3	142,827	282,195	425,022
Difference	-7,118	-13,344	-20,462
Percent Difference ³	-5	-5	-5
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	1,273	331,252	332,525
Alternative 3	1,126	305,992	307,118
Difference	-147	-25,261	-25,407
Percent Difference	-12	-8	-8
Above Normal (12.5%)			
Second Basis of Comparison	388,548	281,850	670,398
Alternative 3	430,489	234,205	664,694
Difference	41,941	-47,645	-5,704
Percent Difference	11	-17	-1
Below Normal (17.5%)			
Second Basis of Comparison	218,115	281,277	499,391
Alternative 3	210,138	294,942	505,080
Difference	-7,977	13,666	5,688
Percent Difference	-4	5	1
Dry (22.5%)			
Second Basis of Comparison	134,348	300,869	435,217
Alternative 3	95,635	299,633	395,268
Difference	-38,713	-1,236	-39,949
Percent Difference	-29	0	-9
Critical (15%)			
Second Basis of Comparison	217,099	238,210	455,309
Alternative 3	202,386	229,599	431,984
Difference	-14,713	-8,612	-23,325
Percent Difference	-7	-4	-5

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-24. Annual Mortality by Cause and Life Stage for Winter-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)				Juvenile Habitat	Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature		
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	0	151,512	107,540	20,257	142,726	22,149	1,300	445,484
Alternative 3	0	135,049	102,763	19,523	145,743	20,541	1,402	425,022
Difference	0	-16,462	-4,776	-734	3,017	-1,607	102	-20,462
Percent Difference ³	-100	-11	-4	-4	2	-7	8	-5
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	0	153,836	1,268	3	176,312	3	1,104	332,525
Alternative 3	0	130,505	1,126	1	174,265	0	1,222	307,118
Difference	0	-23,331	-142	-2	-2,048	-3	118	-25,407
Percent Difference	0	-15	-11	-69	-1	-100	11	-8
Above Normal (12.5%)								
Second Basis of Comparison	0	169,913	268,778	56,974	110,925	62,797	1,012	670,398
Alternative 3	0	119,969	323,517	52,929	113,366	54,043	869	664,694
Difference	0	-49,944	54,739	-4,045	2,441	-8,754	-143	-5,704
Percent Difference	0	-29	20	-7	2	-14	-14	-1
Below Normal (17.5%)								
Second Basis of Comparison	0	157,331	180,614	20,113	122,812	17,388	1,134	499,391
Alternative 3	0	155,899	168,822	21,483	137,826	19,833	1,217	505,080
Difference	0	-1,432	-11,792	1,370	15,015	2,445	83	5,688
Percent Difference	0	-1	-7	7	12	14	7	1
Dry (22.5%)								
Second Basis of Comparison	1	148,149	91,919	21,162	151,231	21,266	1,488	435,217
Alternative 3	0	146,046	61,947	18,345	151,898	15,343	1,689	395,268
Difference	-1	-2,103	-29,972	-2,817	667	-5,923	200	-39,949
Percent Difference	-100	-1	-33	-13	0	-28	13	-9
Critical (15%)								
Second Basis of Comparison	0	129,397	141,609	32,354	106,935	43,136	1,878	455,309
Alternative 3	0	116,643	123,172	33,460	110,932	45,753	2,023	431,984
Difference	0	-12,754	-18,436	1,107	3,997	2,617	145	-23,325
Percent Difference	0	-10	-13	3	4	6	8	-5

1 Based on the 80-year simulation period

2 As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

3 Relative difference of the Annual average

4 Mortality values do not include base mortality

Table B-4-25. Annual Mortality by All Factors for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	Total
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	0	151,512	0	107,540	20,257	142,726	22,146	1,167	3	134	445,484
Alternative 3	0	135,049	0	102,763	19,523	145,743	20,536	1,267	5	135	425,022
Difference	0	-16,462	0	-4,776	-734	3,017	-1,609	100	2	2	-20,462
Percent Difference ³	-100	-11	0	-4	-4	2	-7	9	73	1	-5
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	0	153,836	0	1,268	3	176,312	3	1,057	0	47	332,525
Alternative 3	0	130,505	0	1,126	1	174,265	0	1,188	0	34	307,118
Difference	0	-23,331	0	-142	-2	-2,048	-3	131	0	-13	-25,407
Percent Difference	0	-15	0	-11	-69	-1	-100	12	0	-28	-8
Above Normal (12.5%)											
Second Basis of Comparison	0	169,913	0	268,778	56,974	110,925	62,779	926	17	85	670,398
Alternative 3	0	119,969	0	323,517	52,929	113,366	54,043	799	0	70	664,694
Difference	0	-49,944	0	54,739	-4,045	2,441	-8,737	-128	-17	-15	-5,704
Percent Difference	0	-29	0	20	-7	2	-14	-14	-100	-17	-1
Below Normal (17.5%)											
Second Basis of Comparison	0	157,331	0	180,614	20,113	122,812	17,388	1,093	0	41	499,391
Alternative 3	0	155,899	0	168,822	21,483	137,826	19,832	1,162	1	54	505,080
Difference	0	-1,432	0	-11,792	1,370	15,015	2,444	69	1	14	5,688
Percent Difference	0	-1	0	-7	7	12	14	6	0	34	1
Dry (22.5%)											
Second Basis of Comparison	1	148,149	0	91,919	21,162	151,231	21,264	1,348	3	141	435,217
Alternative 3	0	146,046	0	61,947	18,345	151,898	15,343	1,523	0	166	395,268
Difference	-1	-2,103	0	-29,972	-2,817	667	-5,921	176	-3	25	-39,949
Percent Difference	-100	-1	0	-33	-13	0	-28	13	-100	18	-9
Critical (15%)											
Second Basis of Comparison	0	129,397	0	141,609	32,354	106,935	43,135	1,418	1	460	455,309
Alternative 3	0	116,643	0	123,172	33,460	110,932	45,720	1,566	33	457	431,984
Difference	0	-12,754	0	-18,436	1,107	3,997	2,585	148	32	-3	-23,325
Percent Difference	0	-10	0	-13	3	4	6	10	2,700	-1	-5

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-26. Annual Potential Production for Winter-Run Chinook Salmon

Analysis Period	Annual Potential Production (# of Fish/year)
Long-term	
Full Simulation Period¹	
Second Basis of Comparison	1,885,400
Alternative 5	1,883,178
Difference	-2,222
Percent Difference ³	0
Water Year Types²	
Wet (32.5%)	
Second Basis of Comparison	1,930,740
Alternative 5	1,943,241
Difference	12,501
Percent Difference	1
Above Normal (12.5%)	
Second Basis of Comparison	1,746,928
Alternative 5	1,698,809
Difference	-48,120
Percent Difference	-3
Below Normal (17.5%)	
Second Basis of Comparison	1,847,619
Alternative 5	1,898,667
Difference	51,047
Percent Difference	3
Dry (22.5%)	
Second Basis of Comparison	1,894,107
Alternative 5	1,876,977
Difference	-17,130
Percent Difference	-1
Critical (15%)	
Second Basis of Comparison	1,933,573
Alternative 5	1,897,912
Difference	-35,661
Percent Difference	-2
¹ Based on the 80-year simulation period ² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD. ³ Relative difference of the annual average	

Table B-4-27. Annual Mortality by Life Stage for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)				
	Eggs	Fry	Pre-Smolt	Immature-Smolt	Juvenile (Pre & Immature Smolt)
Long-term					
Full Simulation Period¹					
Second Basis of Comparison	259,052	162,983	23,312	137	23,449
Alternative 5	203,248	207,870	29,865	124	29,989
Difference	-55,804	44,886	6,553	-12	6,540
Percent Difference ³	-22	28	28	-9	28
Water Year Types²					
Wet (32.5%)					
Second Basis of Comparison	155,104	176,315	1,060	47	1,107
Alternative 5	87,970	210,570	4,085	28	4,113
Difference	-67,133	34,255	3,025	-19	3,007
Percent Difference	-43	19	285	-40	272
Above Normal (12.5%)					
Second Basis of Comparison	438,691	167,899	63,706	103	63,808
Alternative 5	464,585	236,533	52,336	89	52,425
Difference	25,893	68,634	-11,369	-14	-11,383
Percent Difference	6	41	-18	-13	-18
Below Normal (17.5%)					
Second Basis of Comparison	337,945	142,925	18,481	41	18,522
Alternative 5	191,541	178,323	31,052	108	31,160
Difference	-146,403	35,399	12,571	67	12,638
Percent Difference	-43	25	68	165	68
Dry (22.5%)					
Second Basis of Comparison	240,069	172,393	22,611	143	22,755
Alternative 5	200,255	234,855	20,690	134	20,824
Difference	-39,814	62,462	-1,921	-9	-1,931
Percent Difference	-17	36	-8	-6	-8
Critical (15%)					
Second Basis of Comparison	271,006	139,289	44,553	461	45,014
Alternative 5	253,379	172,126	79,375	365	79,740
Difference	-17,627	32,838	34,822	-96	34,726
Percent Difference	-7	24	78	-21	77

¹ Based on the 80-year simulation period
² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.
³ Relative difference of the Annual average
⁴ Mortality values do not include base mortality
⁵ Eggs mortality includes pre-spawn mortality

Table B-4-28. Annual Mortality by Cause for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)		
	Temperature	Flow	Total
Long-term			
Full Simulation Period¹			
Second Basis of Comparison	149,945	295,539	445,484
Alternative 5	170,139	270,968	441,107
Difference	20,193	-24,571	-4,378
Percent Difference ³	13	-8	-1
Water Year Types²			
Wet (32.5%)			
Second Basis of Comparison	1,273	331,252	332,525
Alternative 5	7,569	295,085	302,654
Difference	6,296	-36,168	-29,872
Percent Difference	495	-11	-9
Above Normal (12.5%)			
Second Basis of Comparison	388,548	281,850	670,398
Alternative 5	499,928	253,615	753,543
Difference	111,380	-28,235	83,145
Percent Difference	29	-10	12
Below Normal (17.5%)			
Second Basis of Comparison	218,115	281,277	499,391
Alternative 5	149,215	251,809	401,024
Difference	-68,900	-29,468	-98,367
Percent Difference	-32	-10	-20
Dry (22.5%)			
Second Basis of Comparison	134,348	300,869	435,217
Alternative 5	146,764	309,170	455,934
Difference	12,416	8,302	20,717
Percent Difference	9	3	5
Critical (15%)			
Second Basis of Comparison	217,099	238,210	455,309
Alternative 5	307,023	198,222	505,246
Difference	89,925	-39,988	49,937
Percent Difference	41	-17	11

¹ Based on the 90-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-29. Annual Mortality by Cause and Life Stage for Winter-Run Chinook Salmon

Analysis Period	Pre-Spawn Mortality	Eggs Flow	Annual Mortality ⁴ (# of Fish/year)					Total
			Eggs - Temperature	Fry - Temperature	Fry - Habitat	Juvenile Temperature	Juvenile Habitat	
Long-term								
Full Simulation Period¹								
Second Basis of Comparison	0	151,512	107,540	20,257	142,726	22,149	1,300	445,484
Alternative 5	0	89,100	114,147	27,082	180,788	28,909	1,080	441,107
Difference	0	-62,412	6,608	6,825	38,061	6,761	-220	-4,378
Percent Difference ³	57	-41	6	34	27	31	-17	-1
Water Year Types²								
Wet (32.5%)								
Second Basis of Comparison	0	153,836	1,268	3	176,312	3	1,104	332,525
Alternative 5	0	84,683	3,288	977	209,593	3,304	809	302,654
Difference	0	-69,153	2,020	974	33,281	3,302	-295	-29,872
Percent Difference	0	-45	159	35,183	19	132,074	-27	-9
Above Normal (12.5%)								
Second Basis of Comparison	0	169,913	268,778	56,974	110,925	62,797	1,012	670,398
Alternative 5	0	80,569	384,016	64,143	172,390	51,769	656	753,543
Difference	0	-89,345	115,238	7,169	61,465	-11,028	-355	83,145
Percent Difference	0	-53	43	13	55	-18	-35	12
Below Normal (17.5%)								
Second Basis of Comparison	0	157,331	180,614	20,113	122,812	17,388	1,134	499,391
Alternative 5	0	103,637	87,904	31,368	146,956	29,943	1,216	401,024
Difference	0	-53,694	-92,710	11,254	24,144	12,556	82	-98,367
Percent Difference	0	-34	-51	56	20	72	7	-20
Dry (22.5%)								
Second Basis of Comparison	1	148,149	91,919	21,162	151,231	21,266	1,488	435,217
Alternative 5	2	94,247	106,007	21,110	213,744	19,645	1,179	455,934
Difference	0	-53,902	14,088	-52	62,514	-1,621	-309	20,717
Percent Difference	30	-36	15	0	41	-8	-21	5
Critical (15%)								
Second Basis of Comparison	0	129,397	141,609	32,354	106,935	43,136	1,878	455,309
Alternative 5	1	81,098	172,281	56,716	115,410	78,025	1,715	505,246
Difference	1	-48,299	30,672	24,363	8,475	34,889	-164	49,937
Percent Difference	0	-37	22	75	8	81	-9	11

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality

Table B-4-30. Annual Mortality by All Factors for Winter-Run Chinook Salmon

Analysis Period	Annual Mortality ⁴ (# of Fish/year)										Total
	Pre-Spawn Mortality	Incubation	Super- imposition	Eggs - Temperature	Fry - Temperature	Fry - Habitat	Pre-smolt - Temperature	Pre-smolt - Habitat	Smolt - Temperature	Smolt - Habitat	
Long-term											
Full Simulation Period¹											
Second Basis of Comparison	0	151,512	0	107,540	20,257	142,726	22,146	1,167	3	134	445,484
Alternative 5	0	89,100	0	114,147	27,082	180,788	28,902	963	7	117	441,107
Difference	0	-62,412	0	6,608	6,825	38,061	6,757	-204	4	-16	-4,378
Percent Difference ³	57	-41	0	6	34	27	31	-17	135	-12	-1
Water Year Types²											
Wet (32.5%)											
Second Basis of Comparison	0	153,836	0	1,268	3	176,312	3	1,057	0	47	332,525
Alternative 5	0	84,683	0	3,288	977	209,593	3,302	784	3	26	302,654
Difference	0	-69,153	0	2,020	974	33,281	3,299	-274	3	-21	-29,872
Percent Difference	0	-45	0	159	35,183	19	131,968	-26	0	-45	-9
Above Normal (12.5%)											
Second Basis of Comparison	0	169,913	0	268,778	56,974	110,925	62,779	926	17	85	670,398
Alternative 5	0	80,569	0	384,016	64,143	172,390	51,732	604	37	52	753,543
Difference	0	-89,345	0	115,238	7,169	61,465	-11,047	-322	19	-33	83,145
Percent Difference	0	-53	0	43	13	55	-18	-35	113	-39	12
Below Normal (17.5%)											
Second Basis of Comparison	0	157,331	0	180,614	20,113	122,812	17,388	1,093	0	41	499,391
Alternative 5	0	103,637	0	87,904	31,368	146,956	29,943	1,108	0	108	401,024
Difference	0	-53,694	0	-92,710	11,254	24,144	12,556	15	0	67	-98,367
Percent Difference	0	-34	0	-51	56	20	72	1	0	165	-20
Dry (22.5%)											
Second Basis of Comparison	1	148,149	0	91,919	21,162	151,231	21,264	1,348	3	141	435,217
Alternative 5	2	94,247	0	106,007	21,110	213,744	19,645	1,045	0	134	455,934
Difference	0	-53,902	0	14,088	-52	62,514	-1,619	-303	-3	-7	20,717
Percent Difference	30	-36	0	15	0	41	-8	-22	-100	-5	5
Critical (15%)											
Second Basis of Comparison	0	129,397	0	141,609	32,354	106,935	43,135	1,418	1	460	455,309
Alternative 5	1	81,098	0	172,281	56,716	115,410	78,016	1,359	9	356	505,246
Difference	1	-48,299	0	30,672	24,363	8,475	34,881	-60	8	-104	49,937
Percent Difference	0	-37	0	22	75	8	81	-4	679	-23	11

¹ Based on the 80-year simulation period

² As defined by the Sacramento Valley 40-30-30 Index Water Year Hydrologic Classification (SWRCB 1995). Water years may not correspond to the biological years in SALMOD.

³ Relative difference of the Annual average

⁴ Mortality values do not include base mortality